SUPPLEMENTARY TABLE 1: Comparison of baseline TAVI and SAVR populations

		TAVI (n=14195)	SAVR (n=91962)	p-value
Age, median [IQR], y		83.0 [77.0-86.0]	72.0 [63.9-78.1]	<0.001
	NK	128 (0.9%)	0 (0%)	
Sex, n (%)	Male	7595 (53.5%)	58718 (63.9%)	<0.001
	Female	6318 (44.5%)	33239 (36.1%)	
	NK	282 (0.2%)	5 (0%)	
BMI, n (%)	<30	9796 (69.0%)	59714 (63.8%)	<0.001
	>/=30	3554 (25.0%)	29548 (32.1%)	
	NK	845 (6.0%)	3700 (4.0%)	
DM, n (%)	No	10594 (74.6%)	74401 (80.9%)	<0.04
	Yes	3240 (22.8%)	13698 (14.8%)	
	NK	357 (2.5%)	3863 (4.2%)	
Creatinine, μmol/L, [IQR]		97.0 [78.0-123.0]	86.0[73.0-103.0]	<0.001
	NK	567 (4.4%)	36820 (40.0%)	
On dialysis, n (%)	Yes	158 (1.1%)	242 (0.26%)	<0.001
	No	9857 (69.4%)	54900 (59.7%)	
	NK	4180 (29.4%)	61507 (45.9%)	
Previous PCI, n (%)	Yes	2739 (19.3%)	84444 (91.8%)	<0.001
	No	11027 (77.7%)	4929 (5.4%)	
	NK	429 (3.0%)	2589 (2.8%)	
Smoking, n (%)	Yes	409 (2.9%)	84174 (91.5%)	<0.001
	No	6594 (46.5 %)	7068 (7.7%)	
	NK	7192 (50.7%)	720 (0.7%)	
Pulmonary disease, n (%)	Yes	3726 (26.2%)	14302 (15.5%)	<0.001
	No	9975 (70.3%)	77204 (40.0%)	
	NK	494 (3.4%)	456 (0.5%)	
History of AF, n (%)	Yes	3480 (24.5%)	11809 (12.8%)	<0.001
	No	9882 (69.6%)	74338 (80.8%)	
	NK	834 (5.9%)	6923 (5.2%)	

Previous cardiac surgery, n (%)	Yes	3593 (25.3%)	5815 (6.3%)	<0.001
	No	10453 (73.6%)	80029(87.0%)	
	NK	150 (1.1%)	8222 (8.9%)	
PPM, n (%)	Yes	2688 (18.9%)	1624 (1.7%)	<0.001
	No	10781 (75.9%)	84523 (91.9%)	
	NK	726 (5.1%)	5815 (6.3%)	
Logistic Euroscore, [IQR]		16.1 [10.7-25.3]	5.8 [3.3-10.1]	<0.001
	NK	5848 (41.2%)	890 (1.0%)	

AF = atrial fibrillation; BMI = body mass index; CABG = coronary artery bypass grafting; CI = confidence interval; IQR = interquartile range; NK = not known; PCI = percutaneous coronary intervention; PPM = permanent pacemaker; TAVI = transcatheter aortic valve implantation; SAVR = surgical aortic valve replacement