






Original research

Prognosis of patients with hypertrophic cardiomyopathy and low-normal left ventricular ejection fraction

You-Jung Choi,^{1,2} Hyung-Kwan Kim ,^{3,4} In-Chang Hwang,^{2,5} Chan Soon Park,³ Tae-Min Rhee,^{3,4} Hyun-Jung Lee,^{3,4} Jun-Bean Park ,^{3,4} Yeonyee Elizabeth Yoon ,^{4,5} Seung-Pyo Lee ,^{3,4} Goo-Yeong Cho ,^{2,4,5} Yong-Jin Kim^{2,3,4}

► Additional supplemental material is published online only. To view, please visit the journal online (<http://dx.doi.org/10.1136/heartjnl-2022-321853>).

For numbered affiliations see end of article.

Correspondence to

Professor Hyung-Kwan Kim, Department of Internal Medicine, Seoul National University Hospital, Seoul, 03080, Korea (the Republic of); cardiman73@gmail.com and Assistant professor In-Chang Hwang, Internal Medicine, Seoul National University Bundang Hospital, Seongnam-si, Gyeonggi-do, Korea (Republic of); inchang.hwang@gmail.com

Received 10 September 2022
Accepted 7 December 2022
Published Online First
29 December 2022



► <http://dx.doi.org/10.1136/heartjnl-2022-322218>



© Author(s) (or their employer(s)) 2023. No commercial re-use. See rights and permissions. Published by BMJ.

To cite: Choi Y-J, Kim H-K, Hwang I-C, et al. *Heart* 2023;**109**:771–778.

ABSTRACT

Objective To investigate whether low-normal left ventricular ejection fraction (LVEF) is associated with adverse outcomes in hypertrophic cardiomyopathy (HCM) and evaluate the incremental value of predictive power of LVEF in the conventional HCM sudden cardiac death (SCD)-risk model.

Methods This retrospective study included 1858 patients with HCM from two tertiary hospitals between 2008 and 2019. We classified LVEF into three categories: preserved ($\geq 60\%$), low normal ($50\%–60\%$) and reduced ($<50\%$); there were 1399, 415, and 44 patients with preserved, low-normal, and reduced LVEF, respectively. The primary outcome was a composite of SCD, ventricular tachycardia/fibrillation and appropriate implantable cardioverter-defibrillator shocks. Secondary outcomes were hospitalisation for heart failure (HHF), cardiovascular death and all-cause death.

Results During the median follow-up of 4.09 years, the primary outcomes occurred in 1.9%. HHF, cardiovascular death, and all-cause death occurred in 3.3%, 1.9%, and 5.3%, respectively. Reduced LVEF was an independent predictor of SCD/equivalent events (adjusted HR (aHR) 5.214, 95% CI 1.574 to 17.274, $p=0.007$), adding predictive value to the HCM risk-SCD model (net reclassification improvement 0.625). Compared with patients with HCM with preserved LVEF, those with low-normal and reduced LVEF had a higher risk of HHF (LVEF $50\%–60\%$, aHR 2.457, 95% CI 1.423 to 4.241, $p=0.001$; LVEF $<50\%$, aHR 7.937, 95% CI 3.315 to 19.002, $p<0.001$) and cardiovascular death (LVEF $50\%–60\%$, aHR 2.641, 95% CI 1.314 to 5.309, $p=0.006$; LVEF $<50\%$, aHR 5.405, 95% CI 1.530 to 19.092, $p=0.009$), whereas there was no significant association with all-cause death.

Conclusions Low-normal LVEF was an independent predictor of HHF and cardiovascular death in patients with HCM.

INTRODUCTION

Hypertrophic cardiomyopathy (HCM) is the most common inherited cardiomyopathy characterised by left ventricular (LV) hypertrophy in the absence of explainable abnormal loading conditions.¹ Although major advances in treating disease-related complications have made HCM a controllable disease with extended longevity,^{2,3} approximately 2%–5% of patients experience disease progression to end-stage HCM, defined as an LV ejection

WHAT IS ALREADY KNOWN ON THIS TOPIC

⇒ The 2020 American Heart Association/American College of Cardiology guideline for hypertrophic cardiomyopathy (HCM) recommends a left ventricular ejection fraction (LVEF) $<50\%$ as a predictor of sudden cardiac death (SCD) and an indication for an implantable cardioverter-defibrillator (class of recommendation IIa).

WHAT THIS STUDY ADDS

⇒ Low-normal left ventricular systolic function, in which LVEF is higher than 50% but not in the normal range (ie, LVEF of $50\%–60\%$), was a hazardous condition representing a higher risk of hospitalisation for heart failure and cardiovascular death in patients with HCM. Additionally, LVEF is an independent factor with incremental value for predicting SCD when added to the conventional HCM risk-SCD model.

HOW THIS STUDY MIGHT AFFECT RESEARCH, PRACTICE OR POLICY

⇒ This study provides supportive evidence for the need for more aggressive management and surveillance of patients with HCM with low-normal left ventricular systolic function, who were previously overlooked. Also, our findings support that LVEF should be considered in the HCM risk-SCD model.

fraction (LVEF) $<50\%$.⁴ Although rare, the development of LV systolic dysfunction is considered an unfavourable HCM clinical course with a substantial risk of morbidity and mortality.^{5–7}

The 2020 American Heart Association/American College of Cardiology (AHA/ACC) guideline for HCM recommends an LVEF $<50\%$ as a predictor of sudden cardiac death (SCD).⁸ However, little is known about the clinical implication of low-normal LVEF $50\%–60\%$ in HCM.⁹ Moreover, an LVEF $<50\%$ is only a class IIa recommendation for an implantable cardioverter-defibrillator (ICD) indication,^{4,8} and the additional predictive value of an LVEF $<50\%$ to the 5-year risk-SCD score endorsed by the 2014 European Society of Cardiology (ESC) guidelines has not yet been determined.^{8,10} Therefore, we

Table 1 Baseline characteristics of patients stratified based on left ventricular (LV) ejection fraction (LVEF)

Variable	Total N=1858	LV systolic function			P value
		LVEF ≥60% N=1399	LVEF 50%–60% N=415	LVEF <50% N=44	
Demographic data					
Age, years	60.0 (52.0–71.0)	63.0 (53.0–71.0)*	61.0 (51.0–72.0)	57.0 (45.5–66.0)†	0.010
Male, n (%)	1263 (68.0)	924 (66.0)‡	311 (74.9)†	28 (63.6)	0.002
Systolic blood pressure, mm Hg	129.0 (118.0–140.0)	129.0 (119.0–140.0)	128.0 (117.0–140.0)	122.0 (110.0–141.5)	0.223
Diastolic blood pressure, mm Hg	75.0 (69.0–82.0)	75.0 (69.0–81.0)‡	77.0 (70.0–84.0)†*	75.0 (65.0–80.75)‡	0.022
Body mass index, kg/m ²	24.8 (22.9–26.8)	24.8 (23.0–26.8)	24.8 (22.8–27.0)	23.6 (20.9–26.4)	0.196
Comorbidities, n (%)					
Hypertension	1031 (55.5)	778 (55.6)	235 (56.6)	18 (40.9)	0.135
Diabetes mellitus	387 (20.8)	289 (20.7)	90 (21.7)	8 (18.2)	0.820
Dyslipidaemia	670 (36.1)	516 (36.9)	137 (33.0)	17 (38.6)	0.329
End-stage renal disease	17 (1.0)	15 (1.1)	1 (0.2)	1 (2.3)	0.173
Atrial fibrillation	267 (14.4)	175 (12.5)†*	80 (19.3)†	12 (27.3)†	<0.001
Ischaemic stroke	212 (11.4)	167 (12.0)	38 (9.2)	7 (15.9)	0.182
Mitral regurgitation	40 (2.2)	30 (2.1)	9 (2.2)	1 (2.3)	0.998
Percutaneous coronary intervention	86 (4.6)	63 (4.5)	18 (4.3)	5 (11.4)	0.097
Beta-blocker	865 (46.6)	643 (46.0)	196 (47.2)	26 (59.1)	0.217
ICD implantation	54 (2.9)	38 (2.7)*	11 (2.7)*	5 (11.4)††	<0.001
Risk factors for SCD, n (%)					
Family history of SCD	133 (7.2)	95 (6.8)*	30 (7.2)*	8 (18.2)††	0.016
Unexplained syncope	188 (10.1)	141 (10.1)	46 (11.1)	1 (2.3)	0.181
Non-sustained VT (N=960)§	238 (24.8)	167 (23.1)	64 (30.3)	7 (29.2)	0.085
Echocardiographic data					
LV end-diastolic volume, mL	64.6 (52.7–80.0)	64.0 (52.4–78.9)†*	65.0 (52.0–82.4)†*	98.6 (79.5–115.0)††	<0.001
LV end-systolic volume, mL	23.0 (18.1–51.0)	21.6 (17.1–27.0)†*	28.0 (22.0–35.3)†*	53.8 (44.3–66.2)††	<0.001
LVEF, %	63.9 (60.0–67.4)	65.5 (63.0–68.6)†*	57.4 (55.7–58.7)†*	43.9 (40.8–47.4)††	<0.001
e' velocity, m/s	4.9 (4.0–6.0)	4.9 (4.0–6.0)*	4.9 (4.0–6.0)*	4.0 (3.1–4.5)††	<0.001
E/e' ratio	12.5 (9.8–16.2)	12.3 (9.6–16.1)*	12.1 (10.0–15.7)*	16.7 (14.1–19.5)††	<0.001
LA dimension, mm	43.3 (39.0–49.0)	43.0 (39.0–48.0)†*	44.5 (40.0–50.2)†*	51.5 (42.0–57.5)††	<0.001
LA volume index, mL/m ²	43.8 (34.5–56.5)	42.9 (34.0–55.4)†*	44.7 (36.2–59.1)†*	62.4 (39.4–79.9)††	<0.001
PASP, mm Hg	31.0 (26.2–36.6)	31.0 (26.2–36.0)	30.0 (25.8–35.0)	33.0 (28.8–39.4)	0.159
LV-MWT, mm	18.0 (16.0–20.0)	17.8 (16.0–20.0)	18.0 (16.0–20.0)	17.0 [15.0–19.7]	0.098
LV-MWT ≥30 mm	16 (0.9)	10 (0.7)	6 (1.4)	0 (0.0)	0.302
LVOTmaxPG, mm Hg	6.5 (4.0–14.6)	6.7 (4.3–18.3)*†	5.6 (3.6–10.2)†	4.0 (2.6–8.6)†	<0.001

*P<0.05 compared with the group with reduced LVEF.

†P<0.05 compared with the group with preserved LVEF.

‡P<0.05 compared with the group with low-normal LVEF.

§Among 1858 patients, 960 patients underwent 24-hour Holter monitoring.

E/e', ratio of peak early diastolic transmitral inflow velocity to e' velocity; e' velocity, early diastolic mitral annular velocity; ICD, implantable cardioverter-defibrillator; LA, left atrium; LV-MWT, maximal LV wall thickness; LVOTmaxPG, maximal LV outflow tract pressure gradient; PASP, pulmonary arterial systolic pressure; SCD, sudden cardiac death; VT, ventricular tachycardia.

aimed to investigate the prognostic value of LVEF 50%–60% in a large cohort of patients with HCM. Also, we attempted to confirm the prognostic value of LVEF <50% as an imaging biomarker of SCD/equivalent events to support the 2020 AHA/ACC guideline for HCM.⁸

METHODS

Study design and study population

This cohort study recruited 1936 consecutive patients diagnosed with HCM between 2008 and 2019 from two tertiary university hospitals. An end-diastolic maximal LV wall thickness (LV-MWT) ≥15 mm on echocardiography without abnormal loading conditions that sufficiently explain LV hypertrophy is generally considered diagnostic for HCM.^{8–10} Furthermore, LV-MWT ≥13 mm can be considered diagnostic in family members of patients with HCM⁸ and patients with an LVEF <50% were included regardless of their LV-MWT if they had prior objective evidence of HCM on echocardiography and/or cardiac MRI.¹¹ In this study, all recruited patients

met this diagnostic criterion after reviewing the electronic medical records. Based on exclusion criteria (online supplemental methods), a total of 1858 patients (median age, 60.0 years (52.0–71.0 years); 1263 men (68.0%)) were included in the final analysis (online supplemental figure 1).

Echocardiography and measurement of LVEF

Comprehensive transthoracic echocardiography was performed in a standard fashion using commercially available ultrasound machines (Vivid 7, GE Healthcare, Chicago, Illinois, USA; i33, Philips, Amsterdam, The Netherlands; Sequoia, Siemens Medical Solutions, Malvern, Pennsylvania, USA) according to current guidelines by expert sonographers and reviewed by an echocardiographer.^{12–13} LVEF was calculated using the biplane method of discs as recommended by the current guideline.¹⁴ LVEF measurement reproducibility was calculated in 20 randomly selected patients by demonstrating the intraclass correlation coefficient (ICC).




Figure 1 Annualised event rates of the primary outcomes. Annualised event rates of primary outcome including ventricular tachycardia/fibrillation, appropriate implantable cardioverter-defibrillator (ICD) shock and sudden cardiac death, in patients with hypertrophic cardiomyopathy are classified based on left ventricular ejection fraction (LVEF).

We classified patients into three groups based on their LVEF: preserved LVEF $\geq 60\%$, low-normal LVEF 50%–60% and reduced LVEF $< 50\%$.

Risk factors for SCD and the 5-year HCM risk-SCD score

Patients had undergone a comprehensive personal and family history and physical examination as part of the routine clinical care, two-dimensional transthoracic echocardiography and 24-hour Holter monitoring at baseline. The 5-year risk-SCD scores were calculated for each patient using the 5-year HCM risk-SCD model endorsed by the 2014 ESC guidelines (online supplemental methods).¹⁰ We classified patients into three categories: low risk ($< 4\%$), intermediate risk (4%–6%) and high risk ($\geq 6\%$) of SCD.¹⁰

Definition of clinical outcomes

The primary outcome was a composite of SCD and equivalent events, including documented ventricular tachycardia/fibrillation, appropriate ICD shocks and aborted SCD. Only sustained pulseless or unstable ventricular tachycardia or ventricular fibrillation was considered SCD equivalent events. The secondary outcomes were hospitalisation for heart failure (HHF), cardiovascular death and all-cause death. Detailed definitions of each outcome are described in online supplemental methods.¹⁵

Patients were censored when they underwent heart transplantation due to end-stage HCM. Each patient was followed from the date of the initial echocardiography until either the occurrence of any of the clinical outcomes, death from any cause, the end of the study follow-up (31 December 2020) or completion of 5 years of follow-up, whichever came first.

Potential incremental predictive value

We calculated the net reclassification improvement (NRI) and integrated discrimination improvement (IDI) by comparing the




Figure 2 Kaplan-Meier curves for the primary outcomes in patients with hypertrophic cardiomyopathy classified based on their left ventricular ejection fraction (LVEF). The Kaplan-Meier event-free survival curves for primary outcome, a composite of ventricular tachycardia/fibrillation, appropriate implantable cardioverter-defibrillator shocks and aborted sudden cardiac death.

Table 2 Multivariate Cox proportional regression analysis for the primary and secondary outcomes

Variable	Multivariate analysis*			Multivariable analysis†		
	HR	95% CI	P value	HR	95% CI	P value
Primary outcomes (SCD/equivalent events)						
LVEF, % (as a continuous variable)	0.954	0.917 to 0.993	0.020	–	–	–
Preserved LVEF ≥60%	–	–	–	1.000	Reference	–
Low-normal LVEF 50%–60%	–	–	–	1.838	0.886 to 3.812	0.102
Reduced LVEF <50%	–	–	–	5.214	1.574 to 17.274	0.007
Secondary outcomes						
Hospitalisation for HF						
LVEF, % (as a continuous variable)	0.940	0.912 to 0.970	<0.001	–	–	–
Preserved LVEF ≥60%	–	–	–	1.000	Reference	–
Low-normal LVEF 50%–60%	–	–	–	2.457	1.423 to 4.241	0.001
Reduced LVEF <50%	–	–	–	7.937	3.315 to 19.002	<0.001
Cardiovascular death						
LVEF, % (as a continuous variable)	0.936	0.901 to 0.973	<0.001	–	–	–
Preserved LVEF ≥60%	–	–	–	1.000	Reference	–
Low-normal LVEF 50%–60%	–	–	–	2.641	1.314 to 5.309	0.006
Reduced LVEF <50%	–	–	–	5.405	1.530 to 19.092	0.009
All-cause death						
LVEF, % (as a continuous variable)	0.967	0.937 to 0.999	0.044	–	–	–
Preserved LVEF ≥60%	–	–	–	1.000	Reference	–
Low-normal LVEF 50%–60%	–	–	–	1.524	0.942 to 2.466	0.086
Reduced LVEF <50%	–	–	–	2.349	0.834 to 6.613	0.106

The primary outcome was a composite of SCD and equivalent events, including documented ventricular tachycardia/fibrillation, appropriate implantable cardioverter-defibrillator shocks and aborted SCD.

*Adjusted for baseline variables with a $p < 0.1$ in the univariate analysis with LVEF as a continuous variable (%) (refer to online supplemental table 2).

†Adjusted for baseline variables with a $p < 0.1$ in the univariate analysis with LVEF as a categorical variable (LVEF ≥60%, LVEF 50%–60% and LVEF <50%) (refer to online supplemental table 2).

HF, heart failure; LVEF, left ventricular ejection fraction; SCD, sudden cardiac death.

predictive probabilities of the HCM risk-SCD risk category alone with the predictive probabilities of the HCM risk-SCD risk category endorsed by the 2014 ESC guidelines¹⁰ in combination with LVEF as a continuous or categorical variable, respectively. The conventional HCM risk-SCD model uses seven risk factors to calculate the risk-SCD score (online supplemental methods). Therefore, NRI and IDI were calculated only in patients with all seven risk factors (N=897).

Statistical analysis

Continuous variables were expressed as means±SD or medians (IQR), and categorical variables were expressed as frequencies (percentages). We compared the three LVEF groups using either the one-way analysis of variance or Kruskal-Wallis test for continuous variables and the X^2 test or Fisher's exact test for categorical variables. In the post hoc analysis, we used the Tukey honest significant difference test for pairwise comparisons.

Cumulative event-free survival was estimated using the Kaplan-Meier method and compared with the log-rank test. We performed pairwise post hoc analysis for a log-rank test using a Bonferroni adjustment for multiple comparisons. The proportional hazards assumption was checked using a statistical test based on Schoenfeld residuals and their plots. Time zero was defined as the date of the initial echocardiography. HR was calculated in the Cox proportional hazards model and presented as the 95% CI and p value. Variables that first achieved a $p < 0.1$ in the univariate Cox regression were assessed in a multivariable model. Two different multivariable Cox proportional hazards models were created, where LV systolic function was entered as

either a continuous (LVEF in %) or categorical variable (LVEF ≥60%, LVEF 50%–60% and LVEF <50%).

NRI and IDI were computed using the package 'Hmisc' in R programming based on the methods described by Pencina *et al.*¹⁶ Moreover, a competing risk analysis of HHF was performed with mortality as a competing risk.

All analyses used a two-sided p value, and a $p < 0.05$ was considered statistically significant. Statistical analyses were performed using IBM SPSS Statistics for Windows, V.25.0 (IBM Corp) and R software, V.4.2.2 (<http://www.R-project.org>); The R Foundation for Statistical Computing, Vienna, Austria).

RESULTS

Baseline characteristics

The median LVEF in the entire cohort of 1858 patients with HCM was 63.9% (60.0%–67.4%), among which 1399 (75.3%), 415 (22.3%), and 44 (2.4%) patients had a preserved LVEF of ≥60%, low-normal LVEF of 50%–60%, and reduced LVEF of <50%, respectively. The intraobserver and interobserver variability for LVEF measurement were good, with ICCs of 0.860 (95% CI 0.664 to 0.945) and 0.879 (95% CI 0.715 to 0.952), respectively.

The baseline demographic and clinical characteristics of the study population in the three LVEF groups are outlined in [table 1](#). The median age of the study population was 60.0 years (52.0–71.0 years), and the prevalence of hypertension was 55.5%. Patients with reduced LVEF were significantly younger than those with preserved LVEF (63.0 years (53.0–71.0 years) vs 57.0 years (45.5–66.0 years); $p < 0.001$). Atrial fibrillation




Figure 3 Annualised event rates of the secondary outcomes. Annualised event rates of secondary outcomes, such as hospitalisation for heart failure, cardiovascular death and all-cause death in patients with hypertrophic cardiomyopathy, are classified based on left ventricular ejection fraction (LVEF).

was more frequent in patients with low-normal and reduced LVEF than in those with preserved LVEF (19.3% and 27.3% vs 12.5%, $p < 0.001$). Compared with patients with preserved and low-normal LVEF, those with reduced LVEF had the highest prevalence of a family history of SCD (6.8% and 7.2% vs 18.2%, $p = 0.016$). Regarding echocardiographic parameters, patients with reduced LVEF had the largest LV volumes and largest left

atrial (LA) sizes, the lowest e' velocities and the highest E/e' ratios (all $p < 0.001$). The maximal LV outflow tract pressure gradient was smaller in patients with low-normal and reduced LVEF than in those with preserved LVEF ($p < 0.001$). During follow-up, there were two patients who underwent heart transplantation; one patient each in LVEF <50% and LVEF 50%–60% groups after HHF.




Figure 4 Kaplan-Meier curves for hospitalisation for heart failure in patients with hypertrophic cardiomyopathy classified based on their left ventricular ejection fraction (LVEF).




Figure 5 Kaplan-Meier curves for cardiovascular death in patients with hypertrophic cardiomyopathy classified based on their left ventricular ejection fraction (LVEF).




Figure 6 Kaplan-Meier curves for all-cause death in patients with hypertrophic cardiomyopathy classified based on their left ventricular ejection fraction (LVEF).

Primary outcome

During the median follow-up duration of 4.09 years (1.66–5.00 years), the primary outcome was attained in 35 (1.9%). Ventricular tachycardia/fibrillation occurred in 17 (0.9%), appropriate ICD shocks in 4 (0.2%) and SCD in 14 (0.7%). The annualised event rate of the primary outcome according to LVEF categories is shown in [figure 1](#) and online supplemental table 1.

Kaplan-Meier curves showed that the event-free survival probability for the primary outcome was significantly different among the three groups being 98.0%, 95.7%, and 87.0% in patients with preserved, low-normal, and reduced LVEF, respectively (log-rank test, $p < 0.001$, [figure 2](#)). In the univariate Cox analysis, SCD/equivalent events were associated with LVEF (%) and the LVEF categories, as were age, a family history of SCD, unexplained syncope and LA dimension (online supplemental table 2). The multivariable Cox analysis indicated that a higher LVEF (%) was associated with a significantly lower risk of SCD/equivalent events (adjusted HR 0.954, 95% CI 0.917 to 0.993, $p = 0.020$). In addition, a reduced LVEF was an independent predictor of SCD/equivalent events compared with preserved LVEF (adjusted HR 5.214, 95% CI 1.574 to 17.274, $p = 0.007$), whereas low-normal LVEF did not increase the risk of SCD/equivalent events compared with preserved LVEF ([table 2](#)). There was no effect modification by age on the association between LVEF and primary outcomes (online supplemental table 3).

Secondary outcomes

Overall, HHF occurred in 62 (3.3%), cardiovascular death in 36 (1.9%) and all-cause death in 98 (5.3%). The annualised event rates of each secondary outcome according to LVEF categories are shown in [figure 3](#) and online supplemental table 1. The reasons for cardiovascular death included SCD (N=14)/ventricular fibrillation (N=1), heart failure-related death (N=11) and ischaemic stroke-related death (n=6). Other causes include acute myocardial infarction and cardiovascular operation-related or procedure-related complications.

Kaplan-Meier curves showed a significant difference in the event-free survival probability between the three groups in terms of HHF (log-rank test, $p < 0.001$, [figure 4](#)) and cardiovascular

death (log-rank test, $p = 0.001$, [figure 5](#)), but not all-cause death (log-rank test, $p = 0.090$, [figure 6](#)). In the multivariable Cox analysis, both LVEF (%) and the LVEF categories were significantly associated with risks of HHF and cardiovascular death ([table 2](#)). There was no effect modification by age on the association between LVEF and secondary outcomes (online supplemental table 3). When mortality was considered a competing risk, LVEF was again significantly associated with the risk of HHF (online supplemental table 4).

Incremental value of LVEF in predicting SCD

The incremental predictive value of LVEF was assessed in the 897 patients for whom all seven risk factors can be identified. Among them, 681 (75.9%), 194 (21.6%), and 22 (2.5%) patients had preserved, low-normal, and reduced LVEF, respectively. The median 5-year HCM risk-SCD score was highest in patients with reduced LVEF (median score, 2.05% (1.40%–3.35%), 2.37% (1.52%–3.73%) and 2.55% (1.82%–3.52%) in patients with preserved, low-normal and reduced LVEF, respectively; $p = 0.019$), whereas the proportion of the three risk-SCD risk categories did not differ statistically between the three groups ($p = 0.109$) (online supplemental table 5). When added to the 5-year HCM risk-SCD category, LVEF as a continuous and categorical variable significantly improved the risk classification for SCD/equivalent events (LVEF (%): NRI 0.576 and IDI 0.041; the LVEF categories: NRI 0.625 and IDI 0.042) (online supplemental table 6).

DISCUSSION

This large cohort study of patients with HCM evaluated the clinical implications of low-normal LVEF in predicting adverse long-term clinical outcomes including SCD/equivalent events, HHF, cardiovascular death and all-cause death. The main findings are twofold: in patients with HCM, (1) low-normal LV systolic function, defined as an LVEF 50%–60%, was strongly associated with the risk of HHF and cardiovascular death, but not SCD/equivalent events; and (2) reduced LV systolic function, defined as an LVEF $< 50\%$, was an independent predictor for SCD/equivalent events, HHF and cardiovascular death, and provided an added predictive value for SCD/equivalent events to the 5-year HCM risk-SCD model.

Low-normal LV systolic function in HCM

HCM research on the prediction and prevention of SCD to date has focused on end-stage HCM with an LVEF $< 50\%$.^{6,7} Contrastingly, only limited attention has been paid to HCM with LVEF ranges from 50% to 60%. Olivotto *et al* proposed four clinical stages of HCM: non-hypertrophic HCM, classic HCM phenotype, adverse remodelling and overt dysfunction.¹⁷ Among these, the *adverse remodelling stage* represents worsening LV systolic function with relatively preserved clinical and haemodynamic balance. During this stage, several structural and functional features, including an LVEF in the low-normal range may coexist.^{10,18} However, all of these phenomena are not expected to be present in a single patient simultaneously, given that HCM is a heterogeneous disease, and all clinical and pathological manifestations are not always found concomitantly.¹⁷ Nevertheless, there has been no compelling evidence for the clinical and prognostic implications of low-normal LVEF in HCM.¹⁰ In this context, the novel findings of our study illustrate that the risks of HHF and cardiovascular death progressively increased from preserved, low-normal LVEF to end-stage HCM. This suggests that low-normal LVEF is not simply the preceding or transition

phase of end-stage HCM but needs to be considered an independent high-risk population requiring close clinical attention and monitoring.

In addition, despite no statistical significance in the multivariate analysis, SCD/ventricular fibrillation (15 of 36) accounted for a non-negligible fraction of cardiovascular deaths in the low-normal LVEF group. Therefore, large-scale longitudinal independent studies are necessary to establish knowledge of this unnoticed but unique population.

LV systolic dysfunction in HCM

LV systolic dysfunction in HCM is associated with substantial myocardial fibrosis and haemodynamic decompensation, potentially causing the fatal arrhythmia.^{19,20} Notably, end-stage HCM has a few clinical and histopathological characteristics, including HCM occurrence at a young age, heterogeneous patterns of LV remodelling and extensive myocardial fibrosis,^{4,21} all of which are associated with ventricular tachyarrhythmias.^{22,23} We also observed similar findings in the current study; that is, patients with HCM with LVEF <50% were the youngest and had the highest cases in family history of SCD.

Recently, Rowin *et al* demonstrated that end-stage HCM with an LVEF <50% is a significant risk for SCD, with no difference in the frequency of these events in patients with an LVEF <35%, and suggested an LVEF <50% as the threshold for recommending an ICD for primary prevention of SCD in HCM.⁶ In this regard, our study demonstrated that patients with HCM with an LVEF <50% had a fivefold higher risk of SCD/equivalent events than those with preserved LVEF. On the other hand, patients with low-normal LVEF were not at higher risk of SCD/equivalent events than those with preserved LVEF, supporting an LVEF <50% as an appropriate threshold for primary prevention ICDs. Moreover, we showed that a reduced LVEF can predict SCD/equivalent events and has incremental predictive value when added to the 5-year HCM risk-SCD prediction model. This is of value in that the 5-year HCM risk-SCD model does not consider LV systolic dysfunction.⁸ Hence, our findings do support the adoption of an LVEF <50% as a new indication for the primary prevention of SCD in the 2020 AHA/ACC guideline.⁸ Furthermore, the present study provides supportive evidence that LVEF is an additional risk factor that needs to be considered in the 5-year HCM-SCD risk score, especially for low and intermediate-risk patients.^{10,24}

LVEF as an easy-to-use and clinically useful prognostic factor in HCM

HCM is a highly heterogeneous disease with variable timings of disease onset and progression, phenotypes and clinical courses, such as normal longevity, progression to end-stage heart failure or SCD.^{25,26} Therefore, disease progression requiring medical management cannot be effectively predicted, necessitating an individualised approach and periodical follow-up. Although a few imaging markers were suggested as potential prognostic factors in patients with HCM,^{10,15} LVEF remains the most validated and commonly used indicator of LV systolic function because of its high reproducibility and ease of use.²⁷ We found that LVEF not only predicts SCD/equivalent events but also long-term clinical outcomes, such as HHF and cardiovascular death in HCM. Consequently, this easy-to-measure and well-validated imaging marker is an important tool for the clinical monitoring of patients with HCM.

Meanwhile, we did not observe an association between reduced LVEF and all-cause mortality in this study. We cannot

explain this comprehensively but assume that with the substantial improvement in HCM management, a 5-year follow-up might be too short to elucidate the impact of LV systolic dysfunction on all-cause mortality. Of note, a recently reported study demonstrated the supplemental role of LV global longitudinal strain for predicting SCD in HCM.²⁸ Therefore, further studies regarding the clinical implications of the LV global longitudinal strain are warranted to aid risk stratification in patients with HCM and low-normal LVEF.

Study limitations

First, this study was retrospectively analysed on pre-existing data of the prospective HCM registry. Accordingly, there could be missing values of variables to be used in the HCM-SCD risk model, leading to the possibility of selection bias. Second, our study cohort mainly consisted of Asian patients. Thus, the results of our study may not be generalisable given the potential racial and genetic differences among populations with HCM.^{29,30} Moreover, we did not provide genetic information for this population with HCM because only 198 patients underwent genetic testing. Fourth, despite our large HCM cohort, including almost 2000 patients, the number of patients with an LVEF <50% was relatively small, but statistical significance could be confirmed even with a wide CI. This might be partly explained by the fact that we excluded patients who required secondary prevention, who might be already in the burnout stage of HCM. Finally, we did not demonstrate serial changes in LVEF. We believe that this issue is of clinical interest and requires further studies.

CONCLUSIONS

In this large HCM cohort, we observed that patients with low-normal LVEF were at an increased risk of HHF and cardiovascular death compared with patients with preserved LVEF. In addition, LV systolic dysfunction, defined as an LVEF <50%, provided additional prognostic information above and beyond the 5-year HCM risk-SCD score.

Author affiliations

¹Internal Medicine, Korea University Guro Hospital, Seoul, Korea (the Republic of)

²Clinical Medical Sciences, Seoul National University College of Medicine, Seoul, Korea (the Republic of)

³Division of Cardiology, Department of Internal Medicine, Seoul National University Hospital, Seoul, Korea (the Republic of)

⁴Department of Internal Medicine, Seoul National University College of Medicine, Seoul, Korea (the Republic of)

⁵Division of Cardiology, Department of Internal Medicine, Seoul National University Bundang Hospital, Seongnam, Korea (the Republic of)

Contributors YJ Choi and HK Kim designed the study. HK Kim developed the theory and performed the computations. HKK, ICH, JBP, YEY, SPL, GYC, and YJK acquired data (patient recruitment). HK Kim, IC Hwang, and YJ Choi verified the analytical methods. HK Kim and IC Hwang encouraged YJ Choi to investigate and supervised the findings of this work. CS Park, TM Rhee, HJ Lee, JB Park, YE Yoon, SP Lee, GY Cho, and YJ Kim contributed to the interpretation of the results. YJ Choi, IC Hwang, and HK Kim wrote the manuscript. All authors approved the final version and are accountable for the integrity of the work. HKK and ICH accept full responsibility for the work and/or the conduct of the study, had access to the data, and controlled the decision to publish.

Funding This study was supported by a research grant from Seoul National University Research fund (no. 800-20210548) and partly by a research grant from Chong-Kun-Dang (no. 0620214840).

Competing interests G-YC is an International Editorial Advisory Board member for *Heart*.

Patient and public involvement Patients and/or the public were not involved in the design, or conduct, or reporting, or dissemination plans of this research.

Patient consent for publication Not required.

Ethics approval The study conformed to the principles of the Helsinki Declaration 1964 and its later amendments, and the Institutional Review Board of Seoul National University Hospital (no. H-2003-502-1107) and Seoul National University Bundang Hospital (no. B-2004/604-40) approved the study protocol. The requirement for written informed consent was waived due to the retrospective nature of the analyses.

Provenance and peer review Not commissioned; externally peer reviewed.

Data availability statement All data relevant to the study are included in the article or uploaded as supplementary information.

Supplemental material This content has been supplied by the author(s). It has not been vetted by BMJ Publishing Group Limited (BMJ) and may not have been peer-reviewed. Any opinions or recommendations discussed are solely those of the author(s) and are not endorsed by BMJ. BMJ disclaims all liability and responsibility arising from any reliance placed on the content. Where the content includes any translated material, BMJ does not warrant the accuracy and reliability of the translations (including but not limited to local regulations, clinical guidelines, terminology, drug names and drug dosages), and is not responsible for any error and/or omissions arising from translation and adaptation or otherwise.

ORCID iDs

Hyung-Kwan Kim <http://orcid.org/0000-0001-7950-2131>

Jun-Bean Park <http://orcid.org/0000-0003-4053-8713>

Yeonyee Elizabeth Yoon <http://orcid.org/0000-0002-8479-9889>

Seung-Pyo Lee <http://orcid.org/0000-0002-5502-3977>

Goo-Young Cho <http://orcid.org/0000-0002-7067-5535>

REFERENCES

- Watkins H, Ashrafian H, Redwood C. Inherited cardiomyopathies. *N Engl J Med* 2011;364:1643–56.
- Maron BJ, Ommen SR, Semsarian C, et al. Hypertrophic cardiomyopathy: present and future, with translation into contemporary cardiovascular medicine. *J Am Coll Cardiol* 2014;64:83–99.
- Moon I, Lee S-Y, Kim H-K, et al. Trends of the prevalence and incidence of hypertrophic cardiomyopathy in Korea: a nationwide population-based cohort study. *PLoS One* 2020;15:e0227012.
- Harris KM, Spirito P, Maron MS, et al. Prevalence, clinical profile, and significance of left ventricular remodeling in the end-stage phase of hypertrophic cardiomyopathy. *Circulation* 2006;114:216–25.
- Maron MS, Rowin EJ, Olivetto I, et al. Contemporary natural history and management of nonobstructive hypertrophic cardiomyopathy. *J Am Coll Cardiol* 2016;67:1399–409.
- Rowin EJ, Maron BJ, Carrick RT, et al. Outcomes in patients with hypertrophic cardiomyopathy and left ventricular systolic dysfunction. *J Am Coll Cardiol* 2020;75:3033–43.
- Marstrand P, Han L, Day SM, et al. Hypertrophic cardiomyopathy with left ventricular systolic dysfunction: insights from the share registry. *Circulation* 2020;141:1371–83.
- Ommen SR, Mital S, Burke MA, et al. 2020 AHA/ACC guideline for the diagnosis and treatment of patients with hypertrophic cardiomyopathy: a report of the American college of cardiology/American heart association joint committee on clinical practice guidelines. *J Am Coll Cardiol* 2020;76:e159–240.
- Olivetto I, Maron BJ, Appelbaum E, et al. Spectrum and clinical significance of systolic function and myocardial fibrosis assessed by cardiovascular magnetic resonance in hypertrophic cardiomyopathy. *Am J Cardiol* 2010;106:261–7.
- Authors/Task Force members, Elliott PM, Anastasakis A, et al. 2014 ESC guidelines on diagnosis and management of hypertrophic cardiomyopathy: the task force for the diagnosis and management of hypertrophic cardiomyopathy of the European society of cardiology (ESC). *Eur Heart J* 2014;35:2733–79.
- Maron MS. Clinical utility of cardiovascular magnetic resonance in hypertrophic cardiomyopathy. *J Cardiovasc Magn Reson* 2012;14:13.
- Mitchell C, Rahko PS, Blauwet LA, et al. Guidelines for performing a comprehensive transthoracic echocardiographic examination in adults: recommendations from the American society of echocardiography. *J Am Soc Echocardiogr* 2019;32:1–64.
- Nagueh SF, Bierig SM, Budoff MJ, et al. American society of echocardiography clinical recommendations for multimodality cardiovascular imaging of patients with hypertrophic cardiomyopathy: endorsed by the American Society of nuclear cardiology, Society for cardiovascular magnetic resonance, and society of cardiovascular computed tomography. *J Am Soc Echocardiogr* 2011;24:473–98.
- Lang RM, Badano LP, Mor-Avi V, et al. Recommendations for cardiac chamber quantification by echocardiography in adults: an update from the American society of echocardiography and the European association of cardiovascular imaging. *J Am Soc Echocardiogr* 2015;28:1–39.
- Choi Y-J, Kim H-K, Lee SC, et al. Validation of the hypertrophic cardiomyopathy risk-sudden cardiac death calculator in Asians. *Heart* 2019;105:1892–7.
- Pencina MJ, D'Agostino RB, D'Agostino RB, et al. Evaluating the added predictive ability of a new marker: from area under the ROC curve to reclassification and beyond. *Stat Med* 2008;27:157–72.
- Olivetto I, Cecchi F, Poggesi C, et al. Patterns of disease progression in hypertrophic cardiomyopathy: an individualized approach to clinical staging. *Circ Heart Fail* 2012;5:535–46.
- Melacini P, Basso C, Angelini A, et al. Clinicopathological profiles of progressive heart failure in hypertrophic cardiomyopathy. *Eur Heart J* 2010;31:2111–23.
- Raman B, Ariga R, Spartera M, et al. Progression of myocardial fibrosis in hypertrophic cardiomyopathy: mechanisms and clinical implications. *Eur Heart J Cardiovasc Imaging* 2019;20:157–67.
- Briasoulis A, Mallikethi-Reddy S, Palla M, et al. Myocardial fibrosis on cardiac magnetic resonance and cardiac outcomes in hypertrophic cardiomyopathy: a meta-analysis. *Heart* 2015;101:1406–11.
- Galati G, Leone O, Pasquale F, et al. Histological and Histometric characterization of myocardial fibrosis in end-stage hypertrophic cardiomyopathy: a clinical-pathological study of 30 explanted hearts. *Circ Heart Fail* 2016;9:e003090.
- Gulati A, Jabbour A, Ismail TF, et al. Association of fibrosis with mortality and sudden cardiac death in patients with nonischemic dilated cardiomyopathy. *JAMA* 2013;309:896–908.
- Nguyen M-N, Kiriazis H, Gao X-M, et al. Cardiac fibrosis and arrhythmogenesis. *Compr Physiol* 2017;7:1009–49.
- Zeppenfeld K, Tfelt-Hansen J, de Riva M, et al. 2022 ESC guidelines for the management of patients with ventricular arrhythmias and the prevention of sudden cardiac death. *Eur Heart J* 2022;43:3997–4126.
- Tuohy CV, Kaul S, Song HK, et al. Hypertrophic cardiomyopathy: the future of treatment. *Eur J Heart Fail* 2020;22:228–40.
- Lee H-J, Kim J, Chang S-A, et al. Major clinical issues in hypertrophic cardiomyopathy. *Korean Circ J* 2022;52:563–75.
- McDonagh TA, Metra M, Adamo M, et al. 2021 ESC guidelines for the diagnosis and treatment of acute and chronic heart failure. *Eur Heart J* 2021;42:3599–726.
- Lee HJ, Kim HK, Lee SC, et al. Supplementary role of left ventricular global longitudinal strain for predicting sudden cardiac death in hypertrophic cardiomyopathy. *Eur Heart J Cardiovasc Imaging* 2022;23:1108–16.
- Eberly LA, Day SM, Ashley EA, et al. Association of race with disease expression and clinical outcomes among patients with hypertrophic cardiomyopathy. *JAMA Cardiol* 2020;5:83–91.
- Kim HY, Park JE, Lee S-C, et al. Genotype-related clinical characteristics and myocardial fibrosis and their association with prognosis in hypertrophic cardiomyopathy. *J Clin Med* 2020;9:1671.