

S4. Characteristics of the study population subjects with missing data (n=1,180) compared with study population subjects with complete data from Rose Angina Questionnaire (n=28,974).

Characteristics	Compete data (n=28,974)		Missing data (n=1,180)			P*	
			Per protocol	>15% missing data		Per protocol	>15% missing data
Sociodemographics							
Age, median/(IQR)	57.4	(53.7-61.2)	57.6	(54.0-61.3)		0.210	
Female, n/(%)	14,944	(51.6)	564	(47.8)		0.011	
Civil status, living alone, n/(%)	7,505	(25.9)	172	(38.5)	(14.6)	<0.001	<0.001
Born outside of Sweden, n/(%)	4,573	(15.9)	199	(53.4)	(16.7)	<0.001	<0.001
Educational level, low, n/(%)	2,649	(9.1)	76	(18.1)	(6.4)	<0.001	<0.001
Unemployed, n/(%)	4,486	(15.4)	227	(51.2)	(19.2)	<0.001	
Poor economic status, n/(%)	758	(2.6)	31	(10.1)	(2.6)	<0.001	
Psychology							
High degree of general stress, n/(%)	6,042	(20.9)	125	(42.4)	(10.6)	<0.001	<0.001
Depressed, n/(%)	7,849	(27.0)	153	(52.6)	(13.0)	<0.001	<0.001
Lifestyle							
Smoking status							
Never, n/(%)	14,493	(50.0)	170	(41.3)	(14.4)	<0.001	<0.001
Alcohol consumption							
Low/Seldom, n/(%)	13,207	(45.6)	232	(70.5)	(19.7)	<0.001	<0.001
Physical activity							
-Self-estimation							
Never/Low (Grimby 1-2), n/(%)	2,351	(7.8)	91	(23.6)		<0.001	
-Accelerometry							
Sed, %of wear time, mean/(SD)	54	(10)	54	(11)		0.161	
Mod/vig, %of wear time, mean/(SD)	6	(3)	6	(4)		<0.001	
Physical status							
Body mass index, mean/(SD)	26.9	(4.5)	28.0	(4.9)		<0.001	
Waist circumference, mean/(SD)	94	(13)	97	(13)		<0.001	
Systolic blood pressure, mean/(SD)	126	(17)	128	(17)		<0.001	
Diastolic blood pressure, mean/(SD)	77	(11)	78	(10)		0.008	
Clinical chemistry							
Total cholesterol, mean/(SD)	5.5	(1.0)	5.5	(1.1)		<0.001	
HDL cholesterol, mean/(SD)	1.6	(0.5)	1.5	(0.5)		<0.001	
LDL cholesterol, mean/(SD)	3.4	(1.0)	3.5	(1.0)		0.541	
Triglycerides, mean/(SD)	1.2	(0.8)	1.5	(1.1)		<0.001	
Glucose, mean/(SD)	5.8	(1.1)	6.1	(1.7)		<0.001	
HbA1c, mean/(SD)	36	(6)	39	(10)		<0.001	
Hs-CRP, mean/(SD)	2.1	(4.0)	3.0	(9.1)		<0.001	
Estimated GFR, mean/(SD)	85	(12)	88	(13)		<0.001	
Cardiometabolic disease							
Treated hyperlipidemia, n/(%)	2255	(7.9)	31	(7.5)	(2.6)	0.730	<0.001
Treated hypertension, n/(%)	5647	(19.9)	84	(20.1)	(7.1)	0.885	<0.001
Treated diabetes mellitus, n/(%)	1046	(3.7)	25	(6.2)	(2.1)	0.008	<0.001
Previous myocardial infarction, n/(%)	460	(1.6)	9	(2.2)	(0.8)	0.389	<0.001
Angina pectoris, n/(%)	256	(0.9)	18	(4.3)	(1.5)	<0.001	<0.001
Previous revascularization, n/(%)	299	(1.1)	8	(1.9)	(0.7)	0.088	<0.001
Previous stroke, n/(%)	408	(1.4)	9	(2.2)	(0.8)	0.221	<0.001
Treated peripheral artery disease, n/(%)	86	(0.3)	3	(0.7)	(0.3)	0.128	<0.001
Heart failure, n/(%)	143	(0.5)	6	(1.4)	(0.5)	0.008	<0.001
Atrial fibrillation/flutter, n/(%)	521	(1.8)	13	(3.1)	(1.1)	0.054	<0.001
Pulmonary disease							
COPD, n/(%)	349	(1.2)	10	(2.4)	(0.8)	0.033	<0.001
OSAS, n/(%)	1260	(4.4)	25	(6.0)	(2.1)	0.127	<0.001

* Holm Bonferroni corrected $P < 0.04$ considered statistically significant