

Announcing the SKI M-mode to Real-time Conversion Program.

A unique opportunity for owners of the Ekoline 20A:

There'll never be a better
time to expand your M-mode
system to scanning.

Here's what we'll do to help.


We will exchange your Eko-
line 20A M-mode system for a
new factory assembled and calibrated Eko-
Sector I* two-dimensional scanning system
with a full one-year warranty at a price
which is bound to match your needs and

EKO LINE 20A
EKO LINE 20A
EKO SECTOR I
EKO SECTOR I
EKO SECTOR I

budget. And to make your pro-
cedures easier and faster, you
get at no extra charge our M-
mode Transducer Kit and 30°
High Resolution Probe Head.

**Take advantage of this
special offer today.**

For more details, call (7073) 24221, or
return this coupon.



SKI
a SmithKline company

SmithKline Instruments Co. Ltd.
Ridgeway
Welwyn Garden City
Hertfordshire, England AL7 2AA

Please contact me immediately with
complete information about your
M-mode to EkoSector I Conversion
Program. Please phone me at ()
to arrange a time.

Name _____

Hospital/Clinic _____

Street _____

City _____ State _____ Zip _____

Other SmithKline offices: United States—
Sunnyvale, California; Australia—
Brookvale, N.S.W.; Canada—Mississauga,
Ontario; Latin America/Far East—
Sunnyvale, California.

SLOW-TRASICOR[®] in

160 mg oxprenolol hydrochloride BP in a sustained-release formulation



The once daily treatment

Prescribing Notes:

Presentation Slow-Trasicor tablets each contain 160mg oxprenolol hydrochloride BP in a special sustained-release formulation available in simple daily reminder packs of 28 tablets. **Dosage in hypertension** Initially one or two tablets in the morning. **Dosage in angina** Initially one or two tablets in the morning. An evening dose may be beneficial in nocturnal angina. As with other beta blocking drugs, sudden withdrawal of treatment may induce severe and continuous angina. Patients should, therefore, be advised to avoid interruption of established therapy and if withdrawal becomes necessary it should be done gradually. **Side effects** Dizziness, drowsiness, headache, insomnia, excitement and gastro-intestinal disturbance may occur and, rarely, isolated cases of excessive bradycardia. Beta-blockers are occasionally associated with skin rashes and/or dry eyes. If any such reaction is suspected, treatment should be withdrawn gradually. **Precautions** If there is evidence of cardiac failure this must be controlled by digitalis and/or diuretics before and during Slow-Trasicor therapy. Should the pulse rate fall below 50 per minute, then treatment should be restarted at a lower dose. Caution should be observed when treating asthmatics, chronic bronchitis or other individuals where bronchospasm may be provoked. Slow-Trasicor should be given cautiously to patients with alcoholism, metabolic acidosis, during pregnancy or anaesthesia with ether or chloroform. Beta-blockers can mask symptoms of hypoglycaemia and also affect carbohydrate metabolism. It may therefore be necessary to readjust the insulin requirements in diabetic patients. Occasionally hypotension may occur with higher dose levels. **Contra-indications** Patients with atrio-ventricular block, marked bradyardia and cardiogenic shock. **Packs** Cartons of 28 Slow-Trasicor tablets consisting of two reminder calendar foils of 14. Basic NHS price: £6.66 PL0008 0130 PA28 7. Full prescribing information is available on request from CIBA Laboratories, Hertsford, West Sussex. R denotes registered trademark.

C I B A