

















'Inderal' LA
Full
24 hour
protection
from
a single
dose.



INDERAL LA


Propranolol hydrochloride BP
Once daily in hypertension and angina.

'Inderal' LA abridged prescribing information.

Presentation: Long-action capsules each containing 160mg of propranolol hydrochloride BP. **Dosage and Administration:** Oral, once daily. 1. **Hypertension** The starting dose is one capsule daily. If necessary, dosage may be increased to two capsules. 2. **Angina** An adequate response is usually obtained with one capsule daily. 3. **Children** 'Inderal' LA is not intended for use in children. **Contraindications, Warnings, etc:** 'Inderal' LA should not be used: 1. In the presence of second and third degree heart block. 2. If there is a history of bronchospasm. 3. After prolonged fasting. 4. In metabolic acidosis (eg in some diabetics). 5. With verapamil and neither drug should be administered within several days of discontinuing the other. **Precautions** 1. Special care should be taken in patients whose cardiac reserve is poor. 2. Bradycardia (usually less than 50-55 beats/min) indicates that dosage should not be further increased. 3. It is important that a beta blocking agent is not discontinued abruptly. 'Inderal' LA may be withdrawn by first substituting the equivalent dosage in 40mg tablets spread through the day and then gradually reducing the dose. 4. As with all other drugs, 'Inderal' should not be given in pregnancy unless its use is essential. 5. If 'Inderal' LA and clonidine are given concurrently the clonidine should not be discontinued until several days after the withdrawal of the beta blocker (see also prescribing information on clonidine). **Anaesthesia:** 'Inderal' may cause an altered response to stress and therefore it may be necessary to withdraw the drug before surgery; see the data sheet. **Adverse Reactions** 'Inderal' LA is usually well-tolerated. Minor side effects such as cold extremities, nausea, insomnia, lassitude and diarrhoea are usually transient, resolving on withdrawal of the drug. Isolated cases of paraesthesia of the hands have been reported. There have been reports of skin rashes and/or dry eyes associated with the use of beta blocking drugs. The reported incidence is small and in most cases the symptoms have cleared when the treatment was withdrawn. Discontinuation of the drug should be considered if any such reaction is not otherwise explicable. Cessation of therapy with a beta adrenergic blocker should be gradual. In the rare event of intolerance to 'Inderal' LA manifested as bradycardia and hypotension, the drug should be withdrawn and treatment instituted as below. **Overdosage:** Excessive bradycardia can be countered with atropine 1-2mg intravenously, followed, if necessary, by a beta receptor stimulant such as isoprenaline 25 micrograms intravenously or orciprenaline 0.5mg intravenously. Basic NHS cost £6.66 per pack of 28. P.L. 0029/0128.

'Inderal' is a trade mark for propranolol hydrochloride. 'Inderal' LA is a trade mark for propranolol hydrochloride in long-acting formulation.

Full prescribing information is available from:

 Imperial Chemical Industries Limited, Pharmaceuticals Division, Alderley House, Alderley Park, Macclesfield, Cheshire. 🌿🌿🌿

INDERAL LA

Propranolol hydrochloride BP

University of London

ROYAL POSTGRADUATE
MEDICAL SCHOOL

at the
HAMMERSMITH HOSPITAL

Diploma Course in Cardiology

*A one-year course commencing
on 5 October 1981*

This course is intended to provide intensive education, at postgraduate level to 15 students, by formal lectures every morning and seminars every afternoon, with practical, individual training in diagnosis, new methodologies and therapy for groups of two-four students.

CLINICAL APPROACH TO ADULT AND PAEDIATRIC DISEASES
ELECTROCARDIOGRAPHY, INCLUDING STRESS TESTING AND AMBULATORY MONITORING
PHONOCARDIOGRAPHY AND IMPULSE CARDIOGRAPHY
ECHOCARDIOGRAPHY
INTENSIVE CARDIAC CARE
CARDIAC CATHETERISATION AND ANGIOGRAPHY
NUCLEAR CARDIOLOGY
PACEMAKING AND CLINICAL ELECTROPHYSIOLOGY
CARDIAC ROENTENOLOGY
CARDIOVASCULAR PHARMACOLOGY
SURGICAL ASPECTS

Participation in Research Programmes is optional

The course for the diploma is open to graduates in medicine who will have at least six months specialist practice in cardiology before starting the course. Details and application forms available from:

School Office
Royal Postgraduate Medical School
Du Cane Road
London W12 0HS

Tel: 01 743 2030 ext 351