NO.2 1984
----SEQ: B34160000
BRITISH HEART JOURNAL

British Heart Journal

Amonthly journal of cardiology published in association with the British Cardiac Society

| Editorial: Cardiac hypertrophy in athletes D Oakley Cardiac structure and function in cyclists and runners. Comparative echocardiographic | 121 | Cross sectional echocardiographic assessment of left ventricular volume and ejection fraction in patients with tetralogy of Fallot. Comparison with biplane angiographic measurements I Oberhänsli, B Friedli | 191 |
|--------------------------------------------------------------------------------------------------------------------------------------------|-----|---------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|-----|
| study R Fagard, A Aubert, J Staessen, E V Eynde, L Vanhees, A Amery | 124 | Life threatening ventricular tachycardias in late survivors of surgically corrected tetralogy of Fallot A Dunnigan, MR Pritzker, D G Benditt, | |
| Physiological left ventricular hypertrophy L M Shapiro | 130 | D W Benson Jr | 198 |
| Computed tomography in patients with hypertrophic cardiomyopathy DL Stone, | | Clinical electrocardiographic studies of bifid T waves Y Watanabe, H Toda, M Nishimura | 207 |
| M C Petch, G I Verney, A K Dixon | 136 | Delayed recovery of left ventricular function after antithyroid treatment. Further evidence for | |
| Computer aided exercise electrocardiographic testing and coronary arteriography in patients with angina pectoris and with myocardial | | reversible abnormalities of contractility in hyperthyroidism J C Forfar, D M Matthews, A D Toft | 045 |
| infarction <i>J-E Angelhed, T I Bjurö, J Ejdebäck,</i> K Selin, D Schlossman, L S C Griffith, | | Effect of left atrial size on the oesophageal | 215 |
| R Bergstrand, A Vedin, C Wilhelmsson | 140 | transit of capsules KS Channer, J Bell, J P Virjee | 223 |
| Submaximal exercise testing early after myocardial infarction. Prognostic importance of exercise induced ST segment elevation | | Case reports | |
| ID Sullivan, D W Davies, E Sowton | 147 | Transient asymmetric ventricular septal hypertrophy in the newborn unassociated | |
| Angiographic features of the coronary arteries during intracoronary thrombolysis <i>P Terrosu</i> , | | with maternal diabetes F Cecchi, A Zuppiroli, A Manetti | 228 |
| G V Ibba, G M Contini, V Franceschino | 154 | Endocardial pacing in a patient with sinus node dysfunction and absent right superior | |
| Double outlet right ventricle. Cross sectional echocardiographic findings, their anatomical | | vena cava D R Ramsdale, R G Charles | 230 |
| explanation, and surgical relevance F J Macartney, M L Rigby, R H Anderson, J Stark, N H Silverman | 164 | Primary cardiac amyloidosis in a young man presenting with angina pectoris | 250 |
| Absent or occult pulmonary artery | | S Saltissi, P J Kertes, D G Julian | 233 |
| P Presbitero, C Bull, S G Haworth, M R de Leval | 178 | Coronary artery vein grafting in a case of Ehlers-Danlos syndrome | |
| Mitral valve abnormalities in patients with right ventricular pressure overload. Analysis by real | | A Redington, J McCue, S Lennox | 237 |
| time cross sectional echocardiography S Nagata, Y-D Park, N Nakanishi, | 100 | Notices | 239 |
| S Beppu, H Sakakibara, Y Nimura | 186 | Correspondence | 240 |