may require much closer links between coronary care units and cardiac x ray suites but it is difficult to separate such admissions to coronary care units from general medical admissions and to send them direct to off site regional cardiac centres. Currently I think that coronary angioplasty with off site surgical back up makes the best use of the facilities we have.

- 1 Richardson SG, Morton P, Murtagh JG, O'Keefe DB, Murphy P, Scott ME. Management of acute coronary occlusion during percutaneous transluminal coronary angioplasty: experience of complications in a hospital without on site facilities for cardiac surgery. *Br Med J* 1990;300:355-8.
- 2 Shaw TRD, Starkey IR, Essop AR, *et al.* Emergency coronary bypass surgery after vessel occlusion at angioplasty in centres without on-site cardiac surgery [Abstract]. *Br Heart J* 1989;61:438.
- 3 Barner HB, Lea JW, Naunheim KS, Stoney WS. Emergency coronary bypass not associated with pre-operative shock in failed angioplasty, after thrombolysis and for acute myocardial infarction. *Circulation* 1989; 79 (suppl I):152-9.
- 4 Golding LAR, Loop FD, Hollman JL, *et al.* Early results of emergency surgery after coronary angioplasty. *Circulation* 1986;74(suppl III):26-9.
- 5 Parsonnet V, Gielchinsky I, Hochberg MS, *et al.* Emergency surgery after failed angioplasty [Abstract]. *J Am Coll Cardiol* 1987;9:123A.
- 6 Talley JD, Weintraub WS, Anderson HV, *et al.* Late clinical outcome of coronary bypass surgery after failed angioplasty [Abstract]. *Circulation* 1987;76 (suppl IV):352.
- 7 ACC/AHA Task Force Report. Guidelines for percutaneous transluminal coronary angioplasty. *J Am Coll Cardiol* 1988;12:529-45.
- 8 Bourassa MG, Alderman EL, Bertrand M, *et al.* Report of the Joint International Society and Federation of Cardiology/World Health Organisation Task Force on Coronary Angioplasty. *Eur Heart J* 1988;9:1034-45.
- 9 Reul GJ, Cooley DA, Hallman GL, *et al.* Coronary artery bypass for unsuccessful percutaneous transluminal coronary angioplasty. *J Thorac Cardiovasc Surg* 1984;88: 685-94.
- 10 Killen DA, Hamaker WR, Reed WA. Coronary artery bypass following percutaneous transluminal coronary angioplasty. *Ann Thorac Surg* 1985;40:133-8.
- 11 O'Kane H, Cleland J, Gladstone D, Galvin I, Buckels J. Emergency operation after failed percutaneous angioplasty: how late? *Br Heart J* 1989;61:93-4.
- 12 Detre K, Holubkov R, Kelsey S, *et al.* Percutaneous transluminal coronary angioplasty in 1985-1986 and 1977-1981: The National Heart, Lung and Blood Registry. *N Engl J Med* 1988;318:265-70.

VIEWS FROM THE PAST *Ernest Henry Starling*

To cardiologists, Starling is best known for his law of the heart, which indicated that cardiac output is determined by diastolic fibre length. Here we see him as a young man at the threshold of his career. In 1899, at the age of 33, he succeeded Sharpey-Schafer as Jodrell professor of physiology at University College, London (Lewis collection).

DENNIS M KRIKLER