End diastolic flow velocity just beneath the aortic isthmus assessed by pulsed Doppler echocardiography: a new predictor of the aortic regurgitant fraction

Christophe Tribouilloy, Pierre Avinée, Wei Feng Shen, Jean-Luc Rey, Michel Slama, Jean-Philippe Lesbre

Abstract

End diastolic flow velocity just beneath the aortic isthmus was measured within 72 hours of cardiac catheterisation by pulsed Doppler echocardiography in 30 controls and 61 patients with aortic regurgitation. The end diastolic flow velocity was determined at the peak R wave on a simultaneously recorded electrocardiogram. In all controls there was no reverse flow at the end diastole beneath the aortic isthmus. In patients with aortic regurgitation the end diastolic flow velocity correlated well with the angiographic grade of regurgitation ($r = 0.81$) and regurgitant fraction ($r = 0.82$). The mean (SD) values were 6.3 (5.2), 12.2 (4.3), 22.1 (5.7), and 34.3 (9.3) cm/s for patients with regurgitant fraction of <20%, between 20% and 40%, between 41% and 60%, and >60%, respectively. An end diastolic flow velocity of >18 cm/s predicted a regurgitant fraction of $\geq 40\%$ with a sensitivity of 88.5% and a specificity of 96%.

The study suggests that the pulsed Doppler derived end diastolic flow velocity is a useful index in the routine non-invasive assessment of the severity of aortic regurgitation.

With the widespread application of both pulsed and continuous wave Doppler echocardiography¹⁻³ and colour flow imaging,⁴ several variables have been proposed as a measure of the severity of chronic aortic regurgitation. Clinical study showed pandiastolic flow in the descending aorta in patients with aortic regurgitation.⁵ We found that it was easy to measure the end diastolic flow velocity just beneath the aortic isthmus by pulsed Doppler echocardiography and that this method provides semiquantitative information on the degree of aortic regurgitation.⁶ In the present study we investigated further the usefulness of this measurement in estimating the aortic regurgitant fraction.

Patients and methods

PATIENTS

Seventy patients with chronic aortic regurgitation underwent cardiac catheterisation and Doppler echocardiography within 72 hours of

each other. Nine patients with poor echogenicity were excluded (pure aortic regurgitation in four and associated aortic stenosis in five). We studied the remaining 61 patients (43 men and 18 women, mean (SD) age 60 (17) years (range 22-84)). All patients were in stable sinus rhythm and had no angiographically significant mitral regurgitation. Forty two patients had pure aortic regurgitation and 19 had associated aortic stenosis. The cause of aortic regurgitation was rheumatic fever in 22, degenerative abnormality in 29, infective endocarditis in eight, post-traumatic lesion in one, and prosthetic valve dysfunction in one. All patients had had a clinical diagnosis of aortic regurgitation for more than six months.

Thirty patients without valve disease served as a control group. Cardiac catheterisation and angiography were performed because of atypical chest pain. All patients had normal electrocardiograms, chest x ray, and Doppler findings.

DOPPLER ECHOCARDIOGRAPHIC EXAMINATION

We used an ATL Mark 600 or a Hewlett-Packard 77020 system equipped with a 3 or 2.5 MHz pulsed Doppler transducer respectively. Patients were placed in a supine position with the head overextended and the shoulders raised. The transducer was positioned in the suprasternal notch and directed downwards to bring the ultrasound beam on to the thoracic aorta at a point between its horizontal and descending portions. The sample volume was placed in the proximal descending aorta just beneath the aortic isthmus and the origin of the left subclavian artery. The Doppler beam was aligned in parallel with the axis of aortic blood flow by showing a maximal systolic (negative) flow velocity curve. Doppler filters were adjusted to minimal values, and gains were reduced to obtain a contrasted but uninterrupted display of the model Doppler frequencies. Patients with chronic aortic regurgitation often show reverse flow velocities throughout diastole in the descending aorta and the flow velocity is relatively constant after mid diastole.⁷ The end diastolic flow velocity (cm/s) was measured at the peak of the R wave on a simultaneously recorded electrocardiogram (fig 1).

The reproducibility of this measurement using this method was tested in 20 patients with aortic regurgitation: the mean (SD) intra

Department of
Cardiology, South
Hospital, University of
Picardie, Amiens,
France
C Tribouilloy
P Avinée
W F Shen
J-L Rey
M Slama
J-P Lesbre

Correspondence to
Professor Jean-Philippe
Lesbre, Service de
Cardiologie B, Hôpital Sud,
80000 Amiens Cedex,
France.

Accepted for publication
13 September 1990

Figure 1 Measurement of end diastolic flow velocity by pulsed Doppler echocardiography. (A) Ultrasound beam aligned with the axis of the descending aorta with sample volume just beneath the aortic isthmus and junction with subclavian artery. (B) Flow velocity curve. Systolic flow velocity appears below the O line and diastolic flow velocity above it. The end diastolic flow velocity was measured at the peak of the R wave on a simultaneously recorded electrocardiogram.

Figure 2 Correlation between end diastolic flow velocity measured by pulsed Doppler method and angiographic grade of aortic regurgitation. Mean and standard deviations are shown for each angiographic grade. Some points represent more than one patient.

and inter observer variabilities were 4.4 (3.2)% and 5.6 (3.6)% respectively.

CARDIAC CATHETERISATION

Right and left heart catheterisations were performed within 72 hours of the Doppler echocardiographic studies. Cardiac output was determined by the thermodilution technique and total left ventricular output was measured from left ventriculography performed in the 30° right anterior oblique projection. Regurgitant fraction was calculated by the standard formula. Aortography was performed in 30° right anterior oblique projection with 40–60 ml of contrast media injected through a 7F pigtail catheter posi-

Table 1 Pulsed Doppler prediction of angiographic grade of aortic regurgitation by end diastolic flow velocity. The pulsed Doppler class accorded with the angiographic class in 39 of 61 patients and no patient was misclassified by more than one class

Angiographic grade	Doppler end diastolic flow velocity (cm/s)			
	0–9	> 9–18	> 18–30	> 30
I	5	3	0	0
II	3	13	2	0
III	0	4	16	2
IV	0	0	8	5

tioned 2–3 cm above the aortic valve. The degree of aortic regurgitation was graded on a scale of I to IV by the method of Sellers *et al.*⁸

STATISTICAL ANALYSIS

The results were expressed as mean (1 SD). We tested the relation between the end diastolic flow velocity and angiographic grades of regurgitation by the Spearman correlation analysis; for continuous data we used linear regression analysis.

Results

CONTROL GROUP

In all 30 controls the mean (1 SD) value of peak systolic flow velocity was 1.10 (0.20) m/s. Early diastolic flow beneath the aortic isthmus was detected with a velocity of <0.25 m/s and lasting <200 ms. There was no reverse flow during mid and late diastole.

AORTIC REGURGITATION

In 61 patients there was a good correlation between end diastolic flow velocity and the angiographic grade of aortic regurgitation (fig 2).

Mean (SD) end diastolic flow velocity in eight patients with grade I regurgitation was 5.3 (4.6) cm/s (end diastolic flow velocity was zero in four), for 18 patients with grade II regurgitation it was 11.8 (4.4) cm/s, for 22 patients with grade III regurgitation it was 21.4 (4.3) cm/s, and for 13 patients with grade IV regurgitation it was 30.4 (9.6) cm/s. Though there was some overlap in end diastolic flow velocity among individuals with each of the four grades of aortic regurgitation, this measurement was a good indication of the severity of aortic regurgitation. Thirty one of 35 patients with end diastolic flow velocity >18 cm/s had grade III or IV regurgitation, whereas 24 of 26 patients with end diastolic flow velocity <18 cm/s had grade I or II. Therefore, an end diastolic flow velocity >18 cm/s predicted severe aortic regurgitation (≥ grade III) with a sensitivity of 88.5% and a specificity of 92.3% (table 1).

The end diastolic flow velocity correlated

Figure 3 Correlation between end diastolic flow velocity measured by pulsed Doppler method and regurgitant fraction by haemodynamic method. Broken lines show 95% confidence interval and solid line is the regression line. Some points represent more than one patient.

Table 2 Pulsed Doppler prediction of angiographic regurgitant fraction by end diastolic flow velocity. The pulsed Doppler class accorded with the angiographic class in 46 of 61 patients and no patient was misclassified by more than one class

Regurgitant fraction (%)	Doppler end diastolic flow velocity (cm/s)			
	0-9	> 9-18	> 18-30	> 30
0-20	7	3	0	0
21-40	1	14	1	0
41-60	0	4	20	2
> 60	0	0	4	5

well with the regurgitant fraction measured by a combined haemodynamic and angiographic method (fig 3). Mean (SD) end diastolic flow velocity was 6.3 (5.2) cm/s in 10 patients with a regurgitant fraction < 20%, 12.2 (4.3) cm/s in 16 patients with a regurgitant fraction between 20-40%, 22.1 (5.7) cm/s in 26 patients with a regurgitant fraction between 41-60%, and 34.3 (9.3) cm/s in nine patients with a regurgitant fraction > 60%. The end diastolic flow velocity reliably classified the regurgitant fraction. Thirty one of 35 patients with end diastolic flow velocity > 18 cm/s had a regurgitant fraction > 40%. In contrast, 25 of 26 patients with end diastolic flow velocity < 18 cm/s had a regurgitant fraction of < 40%. Therefore, an end diastolic flow velocity of > 18 cm/s identified patients with regurgitant fraction > 40% with a sensitivity of 88.5% and a specificity of 96% (table 2).

Discussion

An excellent alignment can be achieved between the ultrasound beam and blood flow during Doppler assessment of flow velocity in the descending aorta and the flow velocity at the end of diastole measured just beneath the aortic isthmus is a good indicator of the severity of aortic regurgitation.

Previous studies of the flow profiles in the aorta showed an irregular flow velocity pattern in the ascending aorta with moderate or severe regurgitation.¹⁰ Because flow velocity across the lumen varies in diastole calculation of forward or reverse flow is unreliable. In contrast, flow in the descending aorta produces a much flatter velocity profile.⁹⁻¹¹ In most of our patients we found a narrow band of frequencies even when regurgitation was severe. Furthermore, movement during the cardiac cycle is much less in the aortic isthmus than the ascending aorta,¹¹ allowing better positioning of the sample volume. In our controls the early diastolic reverse flow detected in the descending aorta by the Doppler method may reflect compliance of the aortic wall, and it may be impossible to define a shape that takes account of the alteration in compliance and the diastolic reflux caused by aortic regurgitation.¹⁰ For these reasons, flow velocity during late diastole in the descending aorta has been investigated extensively to assess the severity of aortic regurgitation.

Boughner used continuous Doppler echocardiography to measure the flow velocity in the descending aorta; he found that the ratio

between the diastolic and systolic flow velocity-time integrals allowed measurement of aortic regurgitant fraction.⁵ Touche *et al* proposed a modified method using pulsed Doppler in which the flow velocity-time integral ratio was multiplied by the ratio of aortic diastolic area to systolic area obtained from an M mode echocardiogram.¹¹ They also reported a good correlation between this measurement and regurgitant fraction determined invasively. Diebold *et al* also reported that the ratio of the end diastolic to the peak systolic velocity can also be obtained from pulsed Doppler echocardiography and that it gives semiquantitative information about the severity of aortic regurgitation.¹⁰ Pulsed Doppler is better than continuous wave Doppler because it is not affected by the venous flow velocities during diastole because the site of the sample volume can be accurately located. These three methods of measuring the regurgitant fraction depend largely on determinations of systolic flow velocity. Thus they may not be reliable in the presence of associated aortic stenosis.¹⁰

A preliminary study from our laboratory showed that a simple measure of the end diastolic flow velocity gave an accurate semiquantitative measure of the severity of aortic regurgitation, and it can be easily determined in most patients with aortic regurgitation.⁶ The results of this study confirm our previous observations and further indicate the high sensitivity and specificity of an end diastolic flow velocity of > 18 cm/s in predicting moderate to severe (grade $\geq$ III) aortic regurgitation and identified patients with a regurgitant fraction of > 40%. This suggests that the end diastolic flow velocity may also be used to predict the aortic regurgitant fraction.

There are several potential limitations of this method. If aortic regurgitation is slight, velocities in the reverse flow will be low and may not be recorded if a filter with too high a cut off frequency is used. In patients with a ductus arteriosus, aortopulmonary fistula, aortic coarctation at the isthmus level, or aortic dissection, this method is not feasible because of a continuous flow component in these regions.¹¹ Because the accuracy of end diastolic flow velocity in measuring the severity of aortic regurgitation relies on a relatively constant reverse flow velocity throughout diastole, the regurgitant fraction may be overestimated during tachycardia (> 120 beats/min) and underestimated during bradycardia (< 50 beats/min).¹² In atrial fibrillation, when the diastole is prolonged, there may be no end diastolic flow. The compliance of the left ventricle in acute aortic regurgitation might well be quite different from that of patients with chronic aortic regurgitation. There may be equilibration of aortic and ventricular pressures before end diastole, which may lead to cessation of reverse flow before the proposed measurement.

The end diastolic flow velocity just beneath the aortic isthmus can be measured easily in most patients with chronic aortic regurgitation and is an accurate indicator of the regurgitant fraction. It may be useful in the routine assessment of patients with chronic aortic regurgita-

tion. The effect of various afterloads on this relation should be investigated.

- 1 Ciobanu M, Abbasi AS, Allen M, Herner A, Spellberg R. Pulsed Doppler echocardiography in the diagnosis and estimation of severity of aortic insufficiency. *Am J Cardiol* 1982;49:339-43.
- 2 Masuyama T, Kodama K, Kitabatake A, et al. Non invasive evaluation of aortic regurgitation by continuous wave Doppler echocardiography. *Circulation* 1986;73:460-6.
- 3 Teague S, Heinsimer J, Anderson J, et al. Quantification of aortic regurgitation utilizing continuous wave Doppler ultrasound. *J Am Coll Cardiol* 1986;3:592-9.
- 4 Perry GJ, Helmcke F, Nanda NC, Byard C, Soto B. Evaluation of aortic insufficiency by Doppler color flow mapping. *J Am Coll Cardiol* 1987;9:952-9.
- 5 Boughner DR. Assessment of aortic insufficiency by transcutaneous Doppler ultrasound. *Circulation* 1975;52:874-80.
- 6 Lesbre JP, Avinée P, Isorni C, Rey JL, Choquet D, Mertl C. Echo Doppler assessment of aortic insufficiency by a new pulsed index [Abstract]. *Eur Heart J* 1988;9(suppl 1):1-42.
- 7 Hatle L, Angelsen B. *Doppler ultrasound in cardiology: physical principles and clinical applications*. 2nd ed. Philadelphia: Lea and Febiger, 1985:97-292.
- 8 Sellers RD, Levy MJ, Amplatz K, Lillehei CW. Left retrograde cardioangiography in acquired cardiac disease: technic, indications and interpretations in 700 cases. *Am J Cardiol* 1964;14:437-47.
- 9 Schultz DL. Pressure and flow in large arteries. In: Bergel DH, ed. *Cardiovascular fluid dynamics*. London: Academic Press, 1972:287-97.
- 10 Diebold B, Peronneau P, Blanchard D, et al. Non-invasive quantification of aortic regurgitation by Doppler echocardiography. *Br Heart J* 1983;49:167-73.
- 11 Touche T, Prasquier R, Nitenberg A, De Zuttere D, Gourgon R. Assessment and follow up of patients with aortic regurgitation by an updated Doppler echocardiographic measurement of the regurgitant fraction in the aortic arch. *Circulation* 1985;72:819-24.
- 12 Avinée P, Rey JL, Isorni C, Darras B, Lesbre JP. Validité comparée des critères de quantification de l'insuffisance aortique en doppler pulsé et continu. *Arch Mal Coeur* 1988;81:895-901.