Propionibacterium acnes causing an aortic root abscess

S M Horner, M F Sturridge, R H Swanton

Abstract

A case of endocarditis caused by *Propionibacterium acnes* associated with an aortic root abscess is presented. This supports the current opinion that aortic root abscesses are not necessarily associated with microorganisms of high virulence.

(Br Heart J 1992;68:218-20)

Endocarditis caused by *Propionibacterium* is very rare despite the frequent presence of the bacterium in sebaceous glands and in the bypass filter during operation.¹ A case is presented in which endocarditis of a prosthetic aortic valve caused by *Propionibacterium* was associated with an aortic root abscess.

Case report

A 34 year old man was admitted after a cerebrovascular accident. At the age of 13 he had had rheumatic fever. In 1978 he had been admitted with pericarditis and at that time an aortic regurgitant murmur was heard. He had subsequently progressed well until 1986 when he noticed that he was short of breath on exertion and more tired than usual. On examination he had the signs of aortic regurgitation. Cardiac catheterisation showed severe aortic regurgitation, a normal aortic root, and a dilated ventricle with good function. He underwent aortic valve replacement with a Starr-Edwards prosthesis as the native valve showed the signs of rheumatic damage. He made a good recovery.

On 19 July 1989 he was admitted to hospital with numbness and weakness in his right arm and leg which had come on suddenly. He was smoking 10 cigarettes a day and taking warfarin. On examination he looked well and was afebrile. Auscultation of his heart showed only the signs of a normally functioning aortic valve prosthesis. He had mild expressive dysphasia and decreased sensation over the right side of his face, right arm, and right leg. Muscle power in all muscle groups of the right arm and leg was decreased, and the tendon reflexes were all brisk, and the right plantar response was upgoing. His international normalised ratio was 2.4, haemoglobin 14.1 g/l, white cell count $15.7 \times 10^9/l$, and erythrocyte sedimentation rate 27 mm/h. A computerised tomographic scan of his brain showed a haemorrhage in the left occipital region surrounded by oedema (fig 1).

Blood cultures were incubated for three weeks without terminal subculture. Two sets

of blood cultures produced a growth of *Propionibacterium acnes* in the anaerobic bottles after six days of incubation. Another two sets yielded a similar organism after two weeks of incubation. A midstream urine culture was negative. The chest x ray film and an abdominal ultrasound showed no abnormalities. No complement fixing antibodies were detected for *Coxiella* species or *Chlamydia psittaci*. An autoantibody screen was negative.

The *Propionibacterium acnes* isolated was sensitive to penicillin, gentamicin, vancomycin, teicoplanin, and rifampicin. The minimum inhibitory concentrations were 0.03 mg/l, 1 mg/l, 0.25 mg/l, 0.06 mg/l, and 0.004 mg/l respectively. Minimum bactericidal concentrations were 0.12 mg/l, 16 mg/l, 0.5 mg/l, 0.25 mg/l, and 0.5 mg/l. The patient was treated with benzyl penicillin (2.4 g every four hours) and gentamicin was added when his fever failed to resolve within five days. The serum bactericidal dilution was 1/4 at peak and 1/2 at trough. His fever did not respond so antibiotic treatment was changed to vancomycin (1 g) and rifampicin (300 mg) twice daily giving a serum bactericidal dilution of 1/16 at peak and 1/8 at trough. His symptoms still did not improve, suggesting that a root abscess might be present. An early diastolic murmur was heard and echocardiography suggested the presence of an aortic abscess in showing thickening of the anterior wall of the aortic root and aortic regurgitation. Aortography confirmed an aortic root abscess

Figure 1 Computerised tomographic head scan showing a large occipital haemorrhage.

Department of
Cardiology, Middlesex
Hospital, Mortimer
Street, London
S M Horner
R H Swanton

Department of
Cardiothoracic
Surgery, Middlesex
Hospital, Mortimer
Street, London
M F Sturridge

Correspondence to
Dr S M Horner,
Department of Cardiology,
Middlesex Hospital,
Mortimer Street, London
W1N 8AA.

Figure 2 Aortogram showing an aortic root abscess.

(fig 2). The patient developed the "red man" response to vancomycin and the antibiotics were changed to teicoplanin (400 mg) once daily. But increasing pruritus forced a further change in antibiotic treatment to benzyl penicillin and gentamicin. In view of his recent intracerebral bleed, digital subtraction angiography of his cerebral arteries was undertaken and was normal.

He underwent replacement of the aortic valve and repair of an aortic root abscess; one suture caused complete heart block and was removed but first degree heart block developed on tightening the other sutures. After the operation he developed left bundle branch block and first degree heart block and thus a permanent pacing system was inserted. This was initially a VVI system but when he became dependent on the pacemaker he had symptoms of dyspnoea and tiredness so the pacemaker was replaced by a dual chamber system. This left him symptom free. He continued to receive benzyl penicillin for six weeks after surgery and he has made a complete recovery.

Discussion

Propionibacterium acnes is often considered to be a contaminant² but it has been found to be the principal pathogen in meningitis,³ scalp pustules,⁴ osteomyelitis,⁵ central nervous system shunt infection,⁶ endophthalmitis⁷ and septic arthritis.² *Propionibacterium acnes* is also a rare cause of infective endocarditis.⁸⁻¹⁰ It has been suggested that aortic root abscesses are caused by bacteria that are particularly virulent^{11,12} although this is not borne out by later studies^{13,14} or this case. *Propionibacterium acnes* is a slow growing anaerobic Gram positive coryneform. It has a pathogenic role in acne vulgaris and is found in sebaceous glands, thus cleaning the skin does not prevent operative contamination. Indeed *Propionibacterium* species can often be isolated from the blood of patients undergoing

cardiopulmonary bypass who are given flu-cloxacillin and tobramycin prophylaxis (37%) but rarely from those receiving teicoplanin (2%).¹ It is thus surprising that infection by *Propionibacterium* species is so rarely documented and it has been suggested that this is due to the lack of good laboratory anaerobic conditions and prolonged culture.⁸ *Propionibacterium acnes* is normally sensitive to penicillin, tetracycline, and teicoplanin.

In infective endocarditis, spread of infection to surrounding structures is more common with aortic valves. This may result in an aortic root abscess, annular erosion of the aortic root, mycotic aneurysm, or involvement of the aorticomitral apparatus. Aortic root abscess is more common with prosthetic valve endocarditis occurring in 52% compared with 15% of cases of native valve endocarditis.¹³ Cross sectional echocardiography is helpful in identifying aortic root abscesses but few patients (< 5%) have the obvious finding of an echo free abscess cavity and thus the diagnosis rests on associated findings such as the wall of the aorta being more than 10 mm thick, prosthetic valve rocking, sinus of Valsalva aneurysm, or perivalvular density in the septum of 14 mm or more. When these factors are considered, the positive and negative predictive values are 86% and 87% respectively.¹⁵ Transoesophageal echocardiography is particularly helpful in diagnosing this complication of infective endocarditis¹⁶ and magnetic resonance imaging may also clearly show an aortic root abscess.¹⁷ The role of cardiac catheterisation in patients with infective endocarditis is debatable but most would advise its use before cardiac surgery¹⁸⁻²⁰ although the rationale for this, resting as it does upon diagnostic accuracy, is being constantly eroded as non-invasive diagnostic techniques improve. Interestingly cardiac catheterisation may be much safer than is often thought. In one study of 35 patients with infective endocarditis undergoing cardiac catheterisation there was not one embolic episode despite vegetations being seen echocardiographically on 15 of the 21 valves crossed by the catheter.²⁰ Similar experience has also been reported in another study.²¹ We would not advocate crossing infected valves but think that aortography can be safely used in endocarditis of the aortic valve.

An aortic root abscess itself does not absolutely indicate the necessity for cardiac surgery,¹³ and these abscesses have occasionally healed with medical treatment alone.¹⁷ Some studies, however, show an improved survival rate with operative intervention in infective endocarditis in general especially when associated with a prosthetic valve.^{18,22} If surgery is performed mortality is significantly higher in those with an aortic root abscess (13.6% compared with 2.2% for those without this complication).¹³

This case suggests that a blood culture growing *Propionibacterium acnes* should not be dismissed as a contaminant without first considering the clinical circumstances. In this patient *Propionibacterium acnes* caused an

aortic root abscess. This supports the current opinion that an aortic root abscess is not necessarily associated with more virulent microorganisms.

1 Wilson APR, Felmingham D, Gruneberg RN, Treasure T, Sturridge MF. Contamination of blood during cardiopulmonary bypass: the effect of antibiotic prophylaxis. *J Hosp Infect* 1988;12:35-42.

2 Yocum RC, McArthur J, Petty BG, Diehl AM, Moench TR. Septic arthritis caused by *Propionibacterium acnes*. *JAMA* 1982;248(14):1740-1.

3 Schlesinger JJ, Ross AL. *Propionibacterium acnes* meningitis in a previously normal adult. *Arch Intern Med* 1977;137:921-3.

4 Maibach HI. Scalp pustules due to *Corynebacterium acnes*. *Arch Dermatol* 1967;96:453-5.

5 Newman JH, Mitchell RG. Diptheroid infection of the cervical spine. *Acta Orthop Scand* 1975;46:67-70.

6 Skinner PR, Taylor AJ, Coakham H. *Propionibacteria* as a cause of shunt and neurosurgical infections. *J Clin Pathol* 1978;31:1085-90.

7 Friedman E, Peyman GA, May DR. Endophthalmitis caused by *propionibacterium acnes*. *Can J Ophthalmol* 1978;13:50-2.

8 O'Neill TM, Hone R, Blake S. Prosthetic valve endocarditis caused by *Propionibacterium acnes*. *BMJ* 1988;296:1444.

9 Lewis JF, Abramson JH. Endocarditis due to *Propionibacterium acnes*. *Am J Clin Pathol* 1980;74:690-2.

10 Wilson WR, Martin WJ, Widowske CJ, et al. Anaerobic bacteraemia. *Mayo Clin Proc* 1972;47:639-87.

11 Sheldon WH, Golden A. Abscess of the valve rings of the heart, a frequent but not well recognised complication of acute bacterial endocarditis. *Circulation* 1951;iv:1-12.

12 Utley JR, Mills J. Annular erosion and pericarditis-complications of endocarditis of the aortic root. *J Thorac Cardiovasc Surg* 1972;64:76-81.

13 John RM, Pugsley W, Treasure T, Sturridge MF, Swanton RH. Aortic root complications of infective endocarditis— influence on surgical outcome. *Eur Heart J* 1991;12:241-8.

14 Arnett EN, Roberts WC. Valve ring abscess in active infective endocarditis. *Circulation* 1976;54:140-5.

15 Ellis SG, Goldstein J, Popp RL. Detection of endocarditis-associated perivalvular abscess by two dimensional echocardiography. *J Am Coll Cardiol* 1985;5:647-53.

16 Daniel WG, Schroder E, Mugge A, Lichten PR. Transoesophageal echocardiography in infective endocarditis. *Am J Cardiac Imaging* 1988;2:78-85.

17 Jeang MK, Fuentes F, Gately A, Byrnes J, Lewis M. Aortic root abscess: initial experience using magnetic resonance imaging. *Chest* 1986;89:613-5.

18 Lau JM, Guinn GA, Beall AC Jr, et al. Operative techniques in infective endocarditis. *Annals Thorac Surg* 1981;32:351-6.

19 Cheitlin MD, Mills J. Infective endocarditis. Is cardiac catheterisation usually needed before cardiac surgery? *Chest* 1984;86:4-5.

20 Welton DE, Young JB, Raizner AE, et al. Value and safety of cardiac catheterisation during active infective endocarditis. *Am J Cardiol* 1979;44:1306-10.

21 Mills J, Abbott J, Utley JR, Ryan C. Role of cardiac catheterisation in infective endocarditis. *Chest* 1977;72:576-82.

22 Nihoyannopoulos P, Oakley CM, Exadactlos N, Ribeiro P, Westaby S, Foale RA. Duration of symptoms and the effects of a more aggressive surgical policy: two factors affecting prognosis of infective endocarditis. *Eur Heart J* 1985;6:380-90.

Br Heart J: first published as 10.1136/hrt.68.8.218 on 1 August 1992. Downloaded from <http://heart.bmj.com/> on April 26, 2023 by guest. Protected by copyright.