*International roundup***Primary angioplasty in myocardial infarction***An introductory overview*

Outcome after myocardial infarction can be improved by prompt reperfusion of the infarct-related coronary artery by thrombolysis or by primary angioplasty. Despite the uncertain role for coronary angioplasty as an early intervention after successful thrombolytic therapy, it seems that primary angioplasty can open acutely occluded arteries as effectively as does thrombolysis, making it a rational first intervention for acute infarction in patients in whom thrombolytic therapy is relatively contraindicated. Indeed, several recent studies have suggested that primary angioplasty during the early hours of myocardial infarction may be even more likely than thrombolysis to establish stable reperfusion. These observations raise important questions that are central to the future of coronary care. Problems for discussion and resolution extend beyond the scientific conclusions of ongoing studies to the regional logistic, political, and ethical implications of these findings.

Emerging data strongly support a role for acute angioplasty where it is available and accessible, but the extent and limitations of this role require further clarification. Patient populations with infarction are generally more heterogeneous than those included in selected clinical trials. Are we now at a point where acute angioplasty represents optimal care for some patients with evolving myocardial infarction? Precisely which patients are these, and what additional studies are needed to identify them? What are the end points that govern the overall value of these methods? Is angioplasty directly competitive with thrombolysis as a primary intervention, or are these techniques likely to be complementary? And if they are complementary, which subgroups of patients might be most effectively treated with each method?

Thrombolysis can be undertaken in centres that provide relatively little technical and laboratory support. Angioplasty requires not only an experienced operator

but also continuous staffing of an accessible catheterisation facility. However favourable primary angioplasty may be when performed by enthusiasts during focused trials in large clinical centres, limitations and problems with the method undoubtedly will emerge with growing experience in suboptimal settings, particularly during awkward hours. What will be the comparative value of these methods when they are applied under average conditions, with average technical support? What will be the relative consequences of these methods for the subsequent course and duration of hospital stay, including further use of laboratory resources? Are the ultimate logistics of primary angioplasty more closely defined by the community hospital than by the university centre?

Economic, logistic, and ethical issues arise in parallel with scientific debate. If primary angioplasty is shown to confer important benefit on some patients with acute myocardial infarction, what reasons might limit its general availability? How much would it cost to make these procedures widely accessible? What would be the consequences of widespread adoption of these evolving coronary care strategies for physician training in cardiology, distribution of consultants, technical staffing and support, and laboratory use? Are these medical, political, or ethical problems? Who should decide whether new approaches are appropriate, who should decide when they should be implemented, and who should decide if they are worth the cost?

PAUL KLIGFIELD

*Division of Cardiology, Department of Medicine,
The New York Hospital—Cornell Medical Center,
525 East 68th Street,
New York, New York 10021,
USA*

Primary angioplasty is the most effective treatment for an acute myocardial infarction*A view from the Netherlands*

Over the past decades, great efforts have been made to assess the optimal approach to patients with acute myocardial infarction. Although the use of aspirin,¹ β adrenergic blockers,² and angiotensin converting enzyme inhibitors³ has significantly reduced mortality in these patients, attention has focused on the restoration of normal blood flow in the infarct-related coronary artery. Coronary artery bypass grafting was the only accepted revascularisation therapy in the 1970s. Although data from small studies suggested that the results of early reperfusion by means of bypass grafting were promising,^{4,5} this approach was never tested in large scale studies. In the 1980s intravenous thrombolytic therapy became widely used, with an estimated average reduction of early mortality of 20–30%.^{1,3,6,7} Different thrombolytic regimens do result in small differences in clinical outcome⁸ but all are associated with only modest reperfusion rates. Normal flow (TIMI 3) in the infarct-related artery 90 minutes after the start of thrombolytic therapy was found

in 29–54%, and 5–7 days later in 51–58%.^{7,8} After several major trials showed that angioplasty performed immediately after thrombolytic therapy conferred no additional benefit^{9,10} interest in the role of angioplasty as a treatment for acute infarction dwindled. Thrombolytic therapy followed by “watchful waiting” became an almost universally accepted approach.¹¹ The simultaneous publication of three trials in which this approach was compared with primary coronary angioplasty (angioplasty without prior or concomitant administration of thrombolytic drugs) has rekindled the debate on the role of angioplasty.^{12–14}

PATENCY OF THE INFARCT-RELATED ARTERY

Primary angioplasty performed by experienced operators restores normal (TIMI 3 flow) blood flow in more than 90% of patients.^{12–15} Reocclusion rates after angioplasty are low.¹³ This compares favourably with the 50–70% of patients in whom normal flow is achieved after thrombolytic therapy.^{8,13}