- 4 Langber JJ, Leon A, Borganelli M, Kalbfleisch SJ, El-Atassi R, Calkins H, *et al.* A randomized, prospective comparison of anterior and posterior approaches to radiofrequency catheter ablation of atrioventricular nodal tachycardia. *Circulation* 1993;87:1551-6.
- 5 Haissaguerre M, Gaita F, Fischer F, Commenges D, Montserrat P, d'Ivernois C, *et al.* Elimination of atrioventricular nodal reentrant tachycardia using discrete slow potentials to guide application of radiofrequency energy. *Circulation* 1992;85:2162-75.
- 6 Chen S-A, Chiang C, Tsang W-P, Hsia C-P, Wang D-C, Yeh C-T, *et al.* Selective radiofrequency catheter ablation of fast and slow pathway in 100 patients with atrioventricular nodal reentrant tachycardia. *Am Heart J* 1993;125:1-10.
- 7 Denes P, Wu D, Dhingra RC, Chuquimia R, Rosen KM. Demonstration of dual AV nodal pathways in patients with supraventricular tachycardia. *Circulation* 1973;48:549-55.
- 8 Kocovic DZ, Harada T, Shea JB, Soroff D, Freidman PL. Alterations of heart rate and of heart rate variability after radiofrequency catheter ablation of supraventricular tachycardia. Delineation of parasympathetic pathways in the human heart. *Circulation* 1993;88:1671-81.

IMAGES IN CARDIOLOGY

Pericardial cyst

This chest x ray shows an incidental finding of a smooth round opacity in the left cardiophrenic angle. Pericardial (spring-water) cysts are a rare developmental anomaly most commonly found in the cardiophrenic angles, and more frequently on the right. Occasionally they are located superiorly in the mediastinum. They usually do not cause symptoms or signs and are invariably benign. Computed tomographic scanning is a useful method of distinguishing between cysts and solid tumours.

M K DAVIES