

before any clinical component of their course; recent recommendations that all applicants to medical school should be screened, and immunised if appropriate, have far-reaching implications and have provoked controversy.^{17,18}

In conclusion, among invasive cardiologists in the United Kingdom there is clear scope for improvement in vaccine uptake, particularly in completion of the immunisation regimen. Screening of patients, as recommended, should also be performed. Careful practice to avoid needle stick injury should continue and immunisation should also be offered to nursing and other ancillary staff caring for patients undergoing invasive cardiac procedures.

We thank Dr J Heptonstall, Consultant Medical Microbiologist, Public Health Laboratory Service Communicable Disease Surveillance Centre, London for invaluable advice before the preparation of this report. We also thank all those who completed and returned the questionnaire.

- 1 Recommendations of the Advisory Group on Hepatitis. Protecting health care workers and patients from hepatitis B. HMSO, London 1993.
- 2 Zuckermann A. Who should be immunised against hepatitis B? *BMJ* 1984;289:1243.
- 3 Recommendations of the Expert Advisory Group on AIDS. Guidance for clinical health care workers: protection against infection with HIV/hepatitis B. HMSO, London 1990.
- 4 Centers for Disease Control guidelines for prevention of HIV and hepatitis B viruses to health care and public safety workers. *MMWR* 1989;38(suppl 56, 4-5):31-3.
- 5 Department of Health. Immunisation against infectious diseases. London: HMSO, 1992:110-9.
- 6 Polakoff S. Acute viral hepatitis B, reported to the Public Health Laboratory Service. *J Infect* 1990;20:163-8.
- 7 Heptonstall J. Outbreaks of hepatitis B virus infection associated with infected surgical staff. *Communicable Disease Report* 1991;1(8):R81-5.
- 8 Cartwright A. Professionals as poor responders: variations in and effects of response rates to questionnaires 1961-77. *BMJ* 1978;ii:1419-21.
- 9 Editorial. Few physicians vaccinated against hepatitis B. *Wis Med J* 1989;88:14.
- 10 Kinnersley P. Attitudes of general practitioners towards their vaccination against hepatitis B. *BMJ* 1990;300:238.
- 11 Burden AD, Whorwell PJ. Poor uptake of hepatitis B immunisation amongst hospital-based health care staff. *Postgrad Med J* 1991;67:256-8.
- 12 Buss PW, McCabe M, Verrier Jones ER. Attitudes of paediatricians to HIV and hepatitis B virus infection. *Arch Dis Child* 1991;66:961-5.
- 13 Berridge DC, Galea MH, Evans DF, Pugh S, Hopkinson BR, Makin GS. Hepatitis B immunisation in vascular surgeons. *Br J Surg* 1990;77:585-6.
- 14 Williams JR, Flowerdew ADS. Uptake of immunisation against hepatitis B among surgeons in Wessex Regional Health Authority. *BMJ* 1990;301:154.
- 15 Le F, Porteous MJ. Operating practices of and precautions taken by orthopaedic surgeons to avoid infection with HIV and hepatitis B virus during surgery. *BMJ* 1990;301:167-9.
- 16 Lever AML. Treatment of the chronic hepatitis B virus carrier state. *J Infect* 1988;16:221-9.
- 17 Lever AML. Editorial: Hepatitis B and medical student admission. *BMJ* 1994;308:870-1.
- 18 Kingman S. Hepatitis B status must be known for medical school. *BMJ* 1994;308:876.

CORRECTIONS

Failure of "effective" treatment for heart failure to improve normal customary activity. *J T Walsh, R Andrews, A Evans, A J Cowley (Br Heart J 1995;74:373-6).*

Exercise ventilation after balloon dilatation of the mitral valve. *A P Banning, N P Lewis, J S Elborn, R J C Hall (Br Heart J 1995;74:386-9).*

We regret that the incorrect volume number was given in the strap lines and at the foot of the abstracts of these papers. The correct volume number is 74.

Permanent pacemaker practice at a Scottish district general hospital between 1987 and 1994. *J G Doherty, F Dawson, F Kerr*

We regret that an error appeared in the second sentence of the discussion on page 477 of this article (*Br Heart J* 1995;73:475-8) which should have read, "Our average implantation rate over the 79 months (130/million population/year) compares with the United Kingdom national 1989 value of 148/million population/year."⁵

NOTICES

The 1996 Annual General Meeting of the **British Cardiac Society** will take place at the Scottish Exhibition & Conference Centre, Glasgow from 7 to 9 May.

The **Fourth Annual Molecular Symposium at UCL Medical School on Ischaemic Preconditioning and Adaptation to Ischaemia** will take place on 12 December 1995 in London. For further information please contact Jan Wenley, Symposium Administrator, Department of Molecular Pathology, 46 Cleveland Street, London W1P 6DB (tel: +44 171 380 9343; fax: +44 171 387 3310).