- 8 Gibbs JL, Rothman MT, Rees MR, Parsons JM, Blackburn ME, Ruiz CE. Stenting of arterial duct: a new approach to palliation for pulmonary atresia. *Br Heart J* 1992;67:240-5.
- 9 Ruiz CE, Gamra H, Zhang HP, Garcia EJ, Boucek MM. Brief report: stenting of the ductus arteriosus as a bridge to cardiac transplantation in infants with the hypoplastic left-heart syndrome. *N Engl J Med* 1993;328:1605-8.
- 10 Rosenthal E, Qureshi S, Tynan M. Percutaneous pulmonary valvotomy and arterial duct stenting in neonates with right ventricular hypoplasia. *Am J Cardiol* 1994;74:304-6.
- 11 Jonas RA, Freed MO, Mayer JE, Castaneda AR. Long-term follow-up of patients with synthetic right heart conduits. *Circulation* 1985;72(suppl II):II-77-83.
- 12 McGoon DC, Danielson GK, Puga FT, Ritter DG, Mair DD, Ilstrup DM. Late results after extracardiac conduit repair for congenital cardiac defects. *Am J Cardiol* 1982;49:1741-9.
- 13 Razzouk AJ, Williams WG, Cleveland DC, et al. Surgical connections from ventricle to pulmonary artery: comparison of four types of valved implants. *Circulation* 1992;86(suppl II):II-154-8.
- 14 Zeevi B, Keane JF, Perry SB, Lock JE. Balloon dilation of postoperative right ventricular outflow obstruction. *J Am Coll Cardiol* 1989;14:401-8.
- 15 Ensing GJ, Hagler DJ, Seward JB, Julsrud PR, Mair DD. Caveats of balloon dilation of conduits and conduit valves. *J Am Coll Cardiol* 1989;14:367-400.
- 16 Morrow WR, Palmaz JC, Tio FO, Ehler WJ, VanDellen AF, Mullins CE. Re-expansion of balloon-expandable stents after growth. *J Am Coll Cardiol* 1993;22:2007-13.

IMAGES IN CARDIOLOGY

Extreme pulsus alternans presenting as 2:1 electromechanical dissociation

A 12 year old boy presented with pulmonary oedema and severe hypertension and was found to be in end stage renal failure. The echocardiogram showed a dilated but well contracting left ventricle with mild hypertrophy. He was ventilated for 24 hours, but after haemodialysis was started he was able to be extubated and became ambulant. During his fourth dialysis cycle, 5 days after admission, he became dizzy, hypotensive, and bradycardic. Auscultation and an electrocardiographic monitoring lead revealed a heart rate that was double his pulse rate. The echocardiogram showed that the aortic valve only opened every second electrocardiographic complex (left panel b)

and a Doppler signal in the descending aorta was only present every second electrocardiographic complex (left panel c). Ventricular function was considerably depressed and an M mode tracing showed alternate strong and weak contractions (arrows) (left panel a). The mitral, tricuspid, and pulmonary valves opened with every electrocardiographic complex. Serum calcium was 1.55 mmol/l. After calcium was replaced and an infusion of dobutamine (5 μ g/kg/min) was started he initially developed palpable pulsus alternans (middle panel a, b, c) and then full pulses (right panel a, b, c).

ERIC ROSENTHAL

In all three panels (left, middle, right) a is an M mode tracing of left ventricle, b is an M mode tracing of aortic valve, and c is a Doppler tracing from the descending aorta. All traces are at the same paper speed except for the left panel c which is at half speed. Time markers are 0.2 s. Depth markers are 1 cm in a and b and 20 cm/s in c.