


- failure. *J Nucl Med* 1992;33:471-7.
- 26 Wakasugi S, Wada A, Hasegawa Y, Nakano S, Shibata N. Detection of abnormal cardiac adrenergic neuron activity in adriamycin-induced cardiomyopathy with iodine-123-metaiodobenzylguanidine. *J Nucl Med* 1992;33:208-14.
- 27 Nakajima K, Bunko H, Taki J, Shimizu M, Muramori A, Hisada K. Quantitative analysis of <sup>123</sup>I-meta-iodobenzylguanidine (MIBG) uptake in hypertrophic cardiomyopathy. *Am Heart J* 1990;119:1329-37.
- 28 Stanton MS, Tuli MM, Radtke NL, Heger JJ, Miles WM, Mock BH, et al. Regional sympathetic denervation after myocardial infarction in humans detected noninvasively using I-123-metaiodobenzylguanidine. *J Am Coll Cardiol* 1989;14:1519-26.
- 29 Dae MW, De Marco T, Botvinick EH, O'Connell JW, Hattner RS, Huberty HP, et al. Scintigraphic assessment of MIBG uptake in globally denervated human and canine hearts—implications for clinical studies. *J Nucl Med* 1992;33:1444-50.
- 30 Hellman G, Hertting G, Peskar B. Uptake kinetics and metabolism of 7-<sup>3</sup>H-dopamine in isolated perfused rat heart. *Br J Pharmacol* 1971;41:256-69.
- 31 Merlet P, Delforge J, Syrota A, Angevin E, Mazière B, Crouzel C, et al. Positron emission tomography with <sup>11</sup>C CGP-12177 to assess  $\beta$ -adrenergic receptor concentration in idiopathic dilated cardiomyopathy. *Circulation* 1993;87:1169-78.
- 32 Delforge J, Syrota A, Lançon J-P, Nakajima K, Loc'h C, Janier M, et al. Cardiac beta-adrenergic receptor density measured in vivo using PET, CGP 12177, and a new graphical method. *J Nucl Med* 1991;32:739-48.

## IMAGES IN CARDIOLOGY



### Prolapse of an aortic dissection flap imaged by transoesophageal echocardiography



This 66 year old farmer collapsed in a field while tending his sheep. On admission to hospital he complained of interscapular pain and was noted to have an early diastolic murmur. Here we present images obtained at subsequent transoesophageal echocardiography. These three sequential transverse views (A, B, and C) demonstrate a dilated aortic root and a proximal dissection flap prolapsing during diastole from aorta to left ventricle through the aortic valve.

The aortic valve was excised and the dissection resected down to the level of the valve. A 29 mm St Jude valved conduit was implanted, and the coronary arteries were anastomosed on aortic buttons. He left hospital at 14 days and was well six months later.

TREVOR RICHENS  
TERENCE LEWIS  
PETER MILLS



AC, aortic cusp; DF, dissection flap; FL, false lumen; TL, true lumen; LV, left ventricle; LA, left atrium; AMVL, anterior mitral valve leaflet.