



# The evidence is stacked in its favour

**Established efficacy in both  
hypertension and angina**

**A reliable choice for good  
tolerability in both young and  
elderly patients<sup>1</sup>**

**More consistent compliance  
than nifedipine retard<sup>2</sup>**

**ABBREVIATED PRESCRIBING INFORMATION FOR ISTIN™ (AMLODIPINE): UK. PRESENTATION:** TABLETS CONTAINING 5MG OR 10MG AMLODIPINE. **INDICATIONS:** FIRST-LINE TREATMENT OF HYPERTENSION AND MYOCARDIAL ISCHAEMIA ASSOCIATED WITH STABLE ANGINA PECTORIS OR VASOSPASTIC (PRINZMETAL'S OR VARIANT) ANGINA. **DOSAGE:** FOR HYPERTENSION AND ANGINA, INITIAL DOSAGE 5MG ORALLY ONCE DAILY WHICH MAY BE INCREASED TO A MAXIMUM DAILY DOSAGE OF 10MG. **USE IN CHILDREN:** NOT RECOMMENDED. **USE IN THE ELDERLY:** NORMAL DOSAGE. **USE IN RENAL IMPAIRMENT:** NORMAL DOSAGE. **USE IN HEPATIC IMPAIRMENT:** DOSAGE RECOMMENDATIONS HAVE NOT BEEN ESTABLISHED; USE WITH CAUTION. **CONTRA-INDICATIONS:** KNOWN SENSITIVITY TO DIHYDROPYRIDINES. **WARNINGS AND PRECAUTIONS:** PREGNANCY AND LACTATION: ISTIN SHOULD NOT BE ADMINISTERED DURING PREGNANCY OR LACTATION, OR TO WOMEN OF CHILD-BEARING POTENTIAL UNLESS EFFECTIVE CONTRACEPTION IS USED. **SIDE-EFFECTS:** OEDEMA, HEADACHE, FLUSHING, DIZZINESS, NAUSEA, PALPITATIONS, FATIGUE, ABDOMINAL PAIN AND SOMNOLENCE. LESS COMMONLY, PRURITUS, DYSPNOEA, ASTHENIA, MUSCLE CRAMPS AND DYSPEPSIA. RASH, AND RARELY ERYTHEMA MULTIFORME HAVE BEEN OBSERVED. AS WITH OTHER CALCIUM CHANNEL BLOCKERS, THE FOLLOWING, WHICH CANNOT BE DISTINGUISHED FROM THE NATURAL HISTORY OF THE UNDERLYING DISEASE HAVE BEEN RARELY REPORTED: MYOCARDIAL INFARCTION AND CHEST PAIN. **FURTHER INFORMATION:** STUDIES HAVE SHOWN THAT ISTIN DID NOT LEAD TO CLINICAL

istin™  
AMLODIPINE




#### DETERIORATION

IN NYHA CLASS II-III HEART FAILURE. STUDIES HAVE NOT BEEN PERFORMED IN PATIENTS WITH CLASS IV HEART FAILURE. **LEGAL CATEGORY:** POM. **PACKAGE QUANTITIES AND BASIC NHS COST:** 5MG TABLETS CALENDAR PACK OF 28 £11.85 (PL 0057/0297); 10MG TABLETS CALENDAR PACK OF 28 £17.70 (PL 0057/0298). **FURTHER INFORMATION ON REQUEST. PFIZER LIMITED,** RAMSGATE ROAD, SANDWICH, KENT CT13 9NJ. **REFERENCES:** 1. CROSS BW ET AL. BR J CLIN PRACT, 1993, 47(5): 237-240. 2. DETRY JR. CLIN CARDIOL, 1994, 17 (SUPPL III): 12-16.



# Which of these devices has been proven effective in reducing restenosis?

Recently, more than 15 new devices, including a variety of stents, atherectomy catheters, and ablative lasers, have undergone clinical investigation. But only one, the PALMAZ-SCHATZ™ balloon-expandable STENT, has been proven capable of reducing the rate of restenosis.<sup>1-3</sup>



*Johnson & Johnson*

**INTERVENTIONAL SYSTEMS**

Opening the Way in Interventional Medicine

1. Spaedy TJ, Wilensky RL. Coronary stenting. *ACC Curr J Rev* 1994; 6:59-62.

2. Fischman DL, Leon MB, Bain DS, et al. A randomized comparison of coronary-stent placement and balloon angioplasty in the treatment of coronary artery disease. *N Engl J Med* 1994; 331:496-501.

3. Serruys PW, de Jaegere P, Kiemeneij F, et al. A comparison of balloon-expandable-stent implantation with balloon angioplasty in patients with coronary artery disease. *N Engl J Med* 1994; 331:489-495.

See package insert for full product information.

PALMAZ-SCHATZ is a registered trademark of Johnson & Johnson Interventional Systems Co.

©1995 by Johnson & Johnson Interventional Systems Co. All rights reserved.  
PML101



NOW AVAILABLE FOR ANGINA



MORE ADVANCED THAN ADALAT RETARD  
HAS REDUCED SIDE-EFFECTS<sup>1</sup>  
A SMOOTHER PLASMA PROFILE<sup>2</sup>  
AND COMES IN A ONCE-DAILY DOSE

NOW ONCE DAILY

# WORKS A 24 HOUR DAY IN ANGINA

*Tildiem*<sup>®</sup> **LA**  
*diltiazem 200mg & 300mg*

24 HOUR CONTROL OF ANGINA • CONTROLS HEART RATE • WELL TOLERATED

Tildiem<sup>®</sup> LA200/Tildiem<sup>®</sup> LA300 Abbreviated Prescribing Information (refer to data sheet for full prescribing information). Presentation: Capsules each containing 200mg or 300mg diltiazem in a modified (extended) release formulation. Indications: Tildiem<sup>®</sup> LA200 and Tildiem<sup>®</sup> LA300 are indicated for angina pectoris and mild to moderate hypertension. Dosage and Administration: Tildiem LA200 and Tildiem LA300 capsules should not be chewed but swallowed whole with water, ideally before or during a meal. The usual adult starting dose is Tildiem LA300 once daily. This dose may be titrated up to a maximum of 500mg o.d. (one LA300 capsule and one LA200 capsule). Recommended starting dose in the elderly and patients with impaired hepatic or renal function is Tildiem LA200 once daily. This dose may be increased to one capsule of Tildiem LA300 daily if clinically indicated. Heart rate should be monitored and dose should not be increased if this falls below 50 beats per minute. Contraindications: Pregnancy, women of child-

Lorex Synthelabo

bearing potential, marked bradycardia, sick sinus syndrome, left ventricular failure with stasis, second or third degree AV block in the absence of a functioning pacemaker, concomitant use with dantrolene infusion. Warnings and Precautions: Caution in patients with mild bradycardia, reduced left ventricular function, first degree AV block, prolonged PR interval, and during concomitant use with alpha-blockers, beta-blockers or other drugs known to induce bradycardia. (Refer to data sheet for full information.) Side Effects: Headache, malaise, ankle oedema, hot flushes, gastrointestinal disturbances, skin rash, asthma, fatigue and palpitations. Basic NHS Cost: Tildiem LA200 28 capsules £11.10. Tildiem LA300 28 capsules £11.80. Product Licence Numbers: Tildiem LA200 4969/0016. Tildiem LA300 4969/0014. Legal Category: POM. Tildiem and Lorex Synthelabo are trade marks. Further information is available from Lorex Synthelabo Ltd, Lunar House, Fieldhouse Lane, Globe Park, Marlow, Bucks. SL7 1LW. Date of preparation: January 1996. Code no: TIL 180.





## ZOCOR® (simvastatin, MSD)

### ABRIDGED PRODUCT INFORMATION

Refer to Data Sheet before prescribing.

#### PRESENTATION

Peach, oval-shaped, film-coated tablets, marked 'ZOCOR 10' on one side, containing 10 mg simvastatin, MSD.

Tan, oval-shaped, film-coated tablets, marked 'ZOCOR 20' on one side, containing 20 mg simvastatin, MSD.

#### INDICATIONS

Primary hypercholesterolaemia unresponsive to diet and other non-pharmacological measures.

In patients with coronary heart disease and a plasma cholesterol level of 5.5 mmol/l or greater, to

- reduce risk of mortality
- reduce risk of coronary death and non-fatal myocardial infarction
- reduce risk for undergoing myocardial revascularising procedures (CABG and PTCA)
- slow the progression of coronary atherosclerosis, including reducing development of new lesions and new total occlusions.

#### DOSAGE AND ADMINISTRATION

##### *Hypercholesterolaemia*

Initially 10 mg *nocte*; dose range 10-40 mg once daily *nocte*.

Maximum therapeutic response occurs within four to six weeks. Consider dose reduction if total serum cholesterol level falls below 3.6 mmol/l or if LDL cholesterol falls below 1.94 mmol/l. (See Data Sheet for full dosage instructions.) A standard cholesterol-lowering diet should be continued.

##### *Coronary heart disease*

Starting dose 20 mg day *nocte*. Adjustment of dose as above.

*Concomitant therapy:* 'Zocor' is effective alone or in combination with bile-acid sequestrants. In patients taking immunosuppressants concomitantly with 'Zocor', the maximum recommended dosage is 10 mg day (see below).

*Impaired renal function:* In patients with severe renal insufficiency (creatinine clearance < 30 ml/min), dosages above 10 mg day should be carefully considered and, if deemed necessary, implemented cautiously.

*Elderly patients:* Modification of dose should not be necessary.

*Children:* Studies to show safety and efficacy have not been done.

#### CONTRA-INDICATIONS

Hypersensitivity to this product; active liver disease or unexplained persistent elevations of serum transaminases; porphyria; pregnancy and breast-feeding; women of childbearing potential unless adequately protected by non-hormonal methods.

#### PRECAUTIONS

*Homozygous familial hypercholesterolaemia:* 'Zocor' is unlikely to be effective.

*Hypertriglyceridaemia:* 'Zocor' is not indicated where hypertriglyceridaemia is the abnormality of most concern.

*Hepatic effects:* Initial and periodic liver-function monitoring recommended. Discontinue if persistent enzyme elevations occur particularly if they rise to three times the upper limit of normal. Caution in patients with a history of liver disease and/or alcoholism.

*Muscle effects:* Clinically insignificant transient mild elevations of creatine phosphokinase have been seen. Therapy with HMG-CoA reductase inhibitors has rarely been associated with myopathy (< 0.1%). Myopathy should be considered in any patient with marked elevations of creatine phosphokinase (CPK) levels (> 10 times the upper limit of normal) or with diffuse myalgias, muscle tenderness and such marked elevations of CPK levels. The patient should be asked to promptly report unexplained muscle pain, tenderness or weakness. The risk of myopathy with HMG-CoA reductase inhibitors is known to be increased by concomitant immunosuppressive therapy including cyclosporine, by concomitant therapy with a fibric acid derivative or lipid-lowering doses of nicotinic acid, and believed to be enhanced by itraconazole. There have been rare reports of severe rhabdomyolysis with secondary acute renal failure. Therefore, the benefits and risks of using simvastatin concomitantly with immunosuppressive o



fibrate drugs, lipid-lowering doses of nicotinic acid, or itraconazole and other systemic azole antifungal derivatives should be carefully considered.

**Pregnancy:** Contra-indicated. One month should elapse between ending therapy with 'Zocor' and planned conception.

**Pediatric use:** Safety and effectiveness in children have not been established.

**Drug interactions:** Care should be taken in patients on concomitant lipid-lowering therapy, particularly fibrates or nicotinic acid derivatives or itraconazole or immunosuppressive therapies, as they are at increased risk of myopathy. In two clinical studies, 'Zocor' modestly potentiated the anticoagulant effect of warfarin; patients taking coumarin derivatives should have their prothrombin time determined prior to therapy with 'Zocor' and monitored as usual. Slight elevation in digoxin levels has been seen when co-administered with 'Zocor'.

#### SIDE EFFECTS

side effects reported most frequently in controlled clinical trials: abdominal pain, constipation, flatulence, asthenia, and headache. Rarely, myopathy. Side effects reported either in long-term extension studies or in marketed use: nausea, diarrhoea, rash, dyspepsia, pruritus, alopecia, dizziness, muscle cramps, myalgia, pancreatitis, xerostomia, peripheral neuropathy, vomiting, and anaemia. Rarely, rhabdomyolysis and hepatitis jaundice occurred. An apparent

hypersensitivity syndrome has been reported rarely which has included some of the following features: angioedema, lupus-like syndrome, polymyalgia rheumatica, vasculitis, thrombocytopenia, eosinophilia, ESR increased, arthritis, arthralgia, urticaria, fever, flushing, dyspnoea, and malaise. Marked and persistent increased serum transaminases have been reported infrequently. Elevated alkaline phosphatase and  $\gamma$ -glutamyl transpeptidase have been reported. Liver-function test abnormalities have generally been mild and transient. Increases in CPK (muscle derived) have been reported. Side effects reported but where a causal relationship to 'Zocor' is not established: depression, erythema multiforme including Stevens-Johnson syndrome, leucopenia, and purpura.

#### PACKAGE QUANTITIES AND BASIC NHS COST

10 mg tablets, £18.29 for 28-tablet calendar pack  
20 mg tablets, £31.09 for 28-tablet calendar pack

#### Product licence numbers:

10 mg tablets, 0025 0241; 20 mg tablets, 0025 0242

**Product licence holder:** Merck Sharp & Dohme Limited, Hertford Road, Hoddesdon, Hertfordshire, EN11 9BU.

POXI Date of review: August 1995.

\* denotes registered trademark of Merck & Co., Inc., Whitehouse Station, NJ, USA.

© Merck Sharp & Dohme Limited 1995. All rights reserved.

# ZOCOR®

(simvastatin, MSD)

**Improving survival in  
post-MI and angina patients**



Merck Sharp & Dohme Limited  
Hertford Road, Hoddesdon, Hertfordshire, EN11 9BU

08-96 ZOCOR 95 Gb 70195 J A

It's easy to be the leader  
in cardiac imaging.  
We just follow you.



Listening to cardiologists puts us way ahead in meeting your clinical and economic needs. That's why more cath labs use Philips X-ray equipment than any other kind. Why hospitals are improving productivity with "swing labs" and CD-Medical digital archiving. And why we're working with leading research institutions to develop practical MR cardiac perfusion and coronary artery imaging.

Why are we always looking ahead?  
Because you are.

*Let's make things better.*



**PHILIPS**




#### Modalim® Prescribing Information

**Presentation** White, capsule-shaped tablets embossed MODALIM on one side with a breakline on the other, each containing 100mg ciprofibrate. **Uses:** For the treatment of primary hyperlipidaemia resistant to appropriate dietary management, including hypercholesterolaemia, hypertriglyceridaemia and combined hyperlipidaemia. In the Fredrickson classification, this includes types IIa, IIb, III and IV. **Dosage Adults:** One tablet (100mg ciprofibrate) per day. **Elderly patients:** As for adults but see precautions and warnings. **Use in impaired renal function:** In moderate renal impairment it is recommended that dosage be reduced to one tablet every other day. Patients should be carefully monitored. MODALIM should not be used in severe renal impairment. **Use in children:** Not recommended since safety and efficacy in children have not been established. **Contra-indications:** Severe hepatic impairment, severe renal impairment, pregnancy and lactation. **Use in Pregnancy and Lactation:** There is no evidence that ciprofibrate is teratogenic, but there were signs of toxicity at high doses in teratogenicity tests in animals, and ciprofibrate has been shown to be excreted in breast milk in rats. In the absence of data on its use in human pregnancy or lactation, Modalim is contraindicated during pregnancy and in nursing mothers.

**Precautions:** The daily dose should not exceed 100mg; doses of 200mg or more have been associated with a high risk of muscle related side effects. Use with caution in patients with impaired renal or hepatic function. If, after several months therapy, serum lipid concentrations are not satisfactorily controlled, additional or different therapeutic measures should be considered. **Interactions:** Ciprofibrate is highly protein bound and therefore likely to displace other drugs from plasma protein binding sites. MODALIM has been shown to potentiate the effect of warfarin indicating that concomitant oral anticoagulant therapy should be given at reduced dosage and adjusted according to prothrombin time. Although there are no specific data, it is likely that ciprofibrate will also potentiate the action of oral hypoglycaemic agents and its action may be affected by oral contraceptives. As with other fibrates, the concomitant use of Modalim with HMG-CoA reductase inhibitors, or other fibrates, may predispose patients to myopathy. **Side effects:** There have been occasional reports of headache, vertigo, rashes and gastrointestinal symptoms including nausea, vomiting, diarrhoea and dyspepsia. Generally these side effects were mild to moderate in nature and occurred early on, becoming less frequent as treatment progressed. Isolated cases of pneumonitis have been reported. As with other drugs of this class, a low incidence of myalgia, elevation of serum creatine phosphokinase, impotence, hair loss and rare cases of rhabdomyolysis, have been reported. Dizziness, drowsiness or tiredness have only rarely been reported in association with MODALIM. It is therefore unlikely to affect ability to drive or to use machinery. Abnormal liver function tests have been observed occasionally. Periodic liver function tests are recommended. MODALIM should be halted if liver enzyme abnormalities persist. **NHS Price** £13.35 per pack of 25 tablets. **Legal Category:** POM. PL11723/06/90. Modalim is a registered trademark.

Modalim is a registered trademark.  
Further information is available from:  
Sanofi Winthrop Ltd, One Onslow Street,  
Guildford, Surrey, GU1 4YS  
Telephone: 01483 505515  
Fax: 01483 35432  
Date of Preparation: December 1995


# MIXED HYPERLIPIDAEMIA - A GREATER RISK OF CHD THAN RAISED CHOLESTEROL ALONE



**Modalim**   
ciprofibrate

LEADING THE CONTROL OF MIXED HYPERLIPIDAEMIA

# A NEW BEGINNING IN CORONARY MEDICINE



## The PALMAZ-SCHATZ™ balloon-expandable STENT introduces a new era for selected patients with symptomatic ischemic heart disease

### REDUCES INCIDENCE OF RESTENOSIS

Implantation of the PALMAZ-SCHATZ stent reduces the incidence of restenosis compared to angioplasty alone.<sup>1,2</sup>

### INCREASES SUCCESS RATE OF ANGIOPLASTY

The procedural success rate of angioplasty performed with the PALMAZ-SCHATZ stent is higher than that of angioplasty alone.<sup>2</sup>

### EXTENSIVE CLINICAL EXPERIENCE

Over 75,000 PALMAZ-SCHATZ stents have been successfully implanted in patients worldwide.

### YIELDS HIGHER RATES OF EVENT-FREE SURVIVAL

In patients treated with the PALMAZ-SCHATZ coronary stent, 87% with *de novo* lesions survived event-free for one year after implantation.<sup>3</sup>

### NEW SPIRAL STENT OFFERS IMPROVED STRENGTH

The newest PALMAZ-SCHATZ stent incorporates a spiral articulation for improved radial strength.

To learn more about the PALMAZ-SCHATZ stent and training programs for stent implantation, contact your JJIS representative.

<sup>1</sup>Serruys PW, de Jaegere P, Kiemeneij F, et al. A comparison of balloon-expandable stent implantation with balloon angioplasty in patients with coronary artery disease. *N Engl J Med* 1994; 331:489-495.

<sup>2</sup>Fischman DL, Leon MB, Raim DS, et al. A randomized comparison of coronary stent placement and balloon angioplasty in the treatment of coronary artery disease. *N Engl J Med* 1994; 331:496-501.

<sup>3</sup>Savage MP, Fischman DL, Schatz RA, et al. Long-term angiographic and clinical outcome after implantation of a balloon-expandable stent in the native coronary circulation. *JACC* 1994; 24:1207-1212.

**Johnson & Johnson**  
**INTERVENTIONAL SYSTEMS**

a Johnson & Johnson company

# SIEMENS



BS EN ISO 9001 : 1994  
Certificate No. FS 32798

## Compact • Customised • Cost efficient COROSKOP Classic

**A system that adapts to your changing requirements.**

As a member of the T.O.P.-Line family your COROSKOP® Classic system is designed with the future in mind, and its modular structure supports system upgrades that fit the realities of your budget. The use of international interface standards opens your cath. lab. to the world of digital image transfer, which today means cost effective, leading edge, CD-M technology. COROSKOP will easily adapt to the technologies of the future.

### MULTIMAP


intelligent image presentation.



**MONOSCREEN** mapping allows the simultaneous display of the reference image and live fluoroscopy on the same monitor. Picture-in-picture and variable image sizing ensure maximum flexibility.

**Image quality - no doubt about it.**

HICOR® digital image processing always delivers outstanding image quality.



Time

*Optimal patient access and expanded application range are the result of the new 3-way C-arm positioning.*

For further information on the **COROSKOP Classic**, please contact:

Siemens plc  
Medical Engineering  
Siemens House  
Oldbury, Bracknell  
Berkshire, RG12 8FZ  
Telephone: 01344 396317  
Fax: 01344 396337



**Siemens UPTIME Services**  
your partner for performance

# Quality around the World

## A worldwide reputation for precision and excellence

Relied upon by leading surgeons, Swann-Morton blades are as worldwide as their reputation for quality.

Alongside their classic surgical range, there are the "Major" orthopaedic blades, "Minor" disposables, and special blades for cervical biopsy and myringotomy. Most recent is the completely new "Fine" range.


Extensively researched, all blades are produced under strict quality control by precise automation, and inspected at every stage. All products are sterilised by Swann-Morton (Services) Ltd.

Swann-Morton always have the blade for the task, uniformly correct in shape, size and weight, and durably sharp – that's why they are famous worldwide!

**INFORMATION  
LINE  
0114 234 4223**

# Swann-Morton

Swann-Morton Limited,  
Owlerton Green, Sheffield S6 2BJ, England.  
Tel: 0114 234 4231. Fax: 0114 231 4966



H0039/SP

BS EN ISO 9001 1994  
Certificate No. FM 14713

Every  
is a



## MANAGEMENT FOR DOCTORS

Jenny Simpson, *Chief Executive, British Association of Medical Managers*  
Richard Smith, *Editor, BMJ*


Every doctor needs to know how to manage, but few receive formal training. This user friendly guide fills the gap with practical advice on all the aspects of management that doctors need to deal with.

- Written by a team of doctors and business academics
- Uses case studies to show examples of good and bad practice
- Invaluable for both hospital doctors and GPs

Topics covered include: working in teams and leadership; getting the best from people; managing change; making finance work for you

*Buy your copy now and find out what management  
has to offer you*


UK £12.95;  
Overseas £14.00  
(BMA members  
£11.95; £13.00)  
ISBN 0 7279 0858 8  
200 pages  
February 1995



To order phone our credit card  
hotline on 0171 383 6185/6245  
(fax 0171 383 6662) Visa, Mastercard, Amex  
accepted or write to: BMJ Publishing Group,  
PO Box 295, London WC1H 9TE enclosing  
your cheque made payable to BMJ.

Also available from medical  
booksellers, including the BMJ  
Bookshop at BMA House.


**BMJ**  
Publishing  
Group



*Technology for lasting clinical success.*

**acuson**






# WANT TO GET YOUR PAPERS PUBLISHED?

## Read this first!

**How to Write a Paper** tells you all you need to know to get a paper accepted for publication. Edited by George M Hall, with contributions from editors of international journals such as the *BMJ*, *British Journal of Anaesthesia* and *Cardiovascular Research*, it explains in refreshingly direct style what journal editors are looking for in a good paper.

### Contents cover:

- How to approach the various sections of a scientific paper – abstract, introduction, methods, results, discussion, and references
- How to write case reports and reviews
- The publishing process – the role of the editor, assessor, publisher, house style and electronic publishing



ISBN 07279 0822 7 126pp 1994  
UK £8.95; Overseas £11.00  
BMA Members: £8.45; Overseas £10.50

**Available from:** BMJ Publishing Group, PO Box 295, London WC1H 9TE (tel: 071 383 6185/6245), medical booksellers or the BMJ bookshop in BMA House.



University of Alberta  
Edmonton

## Director of Division of Cardiology, Department of Medicine

Applications are invited for the position of Director, Division of Cardiology, Department of Medicine at the University of Alberta. The Division currently consists of 13 geographic full time cardiologists located at the University of Alberta Hospital site, with a full range of sub-specialties and diagnostic and therapeutic services. A similar number of part-time clinical faculty in cardiology exist in the region and an active cardiovascular surgical division is present on site, including a cardiac transplantation program. Successful cardiovascular research programs exist in fundamental, clinical and community based research and there is a fully approved postgraduate educational program in cardiology. Significant opportunities to recruit exist and are facilitated by the support of the Alberta Heritage Foundation for Medical Research. Interested applicants should hold an MD and fellowship in the Royal College of Physicians and Surgeons of Canada in Cardiology or equivalent and have demonstrated leadership and clinical expertise along with scholarly accomplishment in research and teaching.

Academic rank and remuneration for this senior position will be commensurate with qualifications and experience. Deadline for applications is 1 May 1996. Please send curriculum vitae and the names and addresses of three references to:

**Dr. P.W. Armstrong**  
Chair, Department of Medicine  
2F1.30 Walter C. Mackenzie  
Health Sciences Centre  
University of Alberta  
Edmonton, Alberta, Canada, T6G 2R7

*The University of Alberta is committed to the principle of equity in employment. As an employer we welcome diversity in the workplace and encourage applications from all qualified women and men, including Aboriginal peoples, persons with disabilities, and members of visible minorities.*

In post menopausal  
hypertension  
why treat her like a man?

A new once daily ACE

# PERDIX<sup>®</sup>

Moexipril

Heart disease is the single largest killer of women in the UK.<sup>1</sup> Hypertension is found frequently in post menopausal women. It has been shown that Perdix controls hypertension and is metabolically neutral in post menopausal women treated with HRT.<sup>2</sup>

**Perdix<sup>®</sup> ▼ 7.5mg and 15mg Tablets. Prescribing Information.**

**Refer to Summary of Product Characteristics before prescribing. Presentation:** Tablets containing 7.5mg and 15mg moexipril hydrochloride. **Uses:** Treatment of hypertension as monotherapy. Second line therapy for the treatment of hypertension in combination with diuretics or calcium antagonists. **Dosage and Administration:** *Untreated Patients:* in patients with uncomplicated essential hypertension the recommended initial dose is 7.5mg once a day. Adjust dosage according to response. Usual dosage range is 15 to 30mg per day as a single daily dose. Doses over 30mg have been used, but do not appear to give a greater effect. If blood pressure is not controlled with Perdix alone, a low dose of a diuretic may be added. *Diuretic treated patients:* symptomatic hypotension may occur occasionally following the initial dose of Perdix. Discontinue diuretic 2-3 days before starting Perdix to reduce the likelihood of hypotension. Adjust dosage of Perdix according to response. Resume diuretic later if required. *Nifedipine treated patients:* initial dose of 3.75mg recommended. *Elderly:* initial dose of 3.75mg followed by titration to optimal response. *Children:* not recommended. *Renal failure:* if creatinine clearance  $\leq 40\text{ml/min}$ , initial dosage should be 3.75mg. *Hepatic cirrhosis:* initial dosage of 3.75mg is recommended. *Afro-Caribbean patients:* may show a reduced therapeutic response. **Contra-indications:** Hypersensitivity to moexipril hydrochloride. History of angioedema following treatment with ACE inhibitors. Pregnancy and lactation. **Special warnings and precautions for use:** **Warnings:** Angioedema: angioedema involving the extremities, face, lips, mucous membranes, tongue, glottis or larynx has been reported in patients treated with ACE inhibitors. Discontinue treatment with Perdix and institute appropriate therapy immediately. **Hypotension:** Perdix can cause symptomatic hypotension, most commonly in volume and/or salt-depleted patients. Correct before initiating therapy with Perdix. **Neutropenia/agranulocytosis:** agranulocytosis and bone marrow depression may result particularly in patients with renal impairment and a collagen-vascular disease. **Precautions:** Changes in renal function may be anticipated in susceptible individuals. Increases in blood urea nitrogen and serum creatinine may occur in hypertensive patients on diuretic therapy and more commonly those with renal artery stenosis in a solitary kidney or bilateral renal artery stenosis. Dosage reduction of Perdix and/or discontinuation of the diuretic may be required. Hyperkalaemia occurs rarely. Risk factors

include renal insufficiency, diabetes mellitus, and concomitant use of potassium-sparing diuretics, potassium supplements, and/or potassium-containing salt substitutes. Patients with hepatic cirrhosis may develop elevated plasma levels of moexipril hydrochloride. In patients undergoing surgery or during anaesthesia with agents that produce hypotension, Perdix will block the angiotensin II formation that could otherwise occur secondary to compensatory renin release. **Interactions:** Combination with diuretics or other antihypertensive agents may have a potentiating effect. Potassium loss caused by thiazide diuretics may be attenuated. Concurrent use of potassium supplements or potassium sparing diuretics may lead to elevated serum potassium. Increased serum lithium levels and symptoms of lithium toxicity have been reported in patients receiving ACE inhibitors during lithium therapy. **Side effects:** include cough, headache, dizziness, fatigue, flushing, and rash. Less commonly, symptomatic hypotension, postural hypotension, syncope, chest pain, angina/myocardial infarction, palpitations, rhythm disturbances and cerebrovascular accident. Increases in serum creatinine levels. Abdominal pain, dyspepsia, constipation, nausea, vomiting, diarrhoea, appetite/weight change, dry mouth, pancreatitis, hepatitis. Upper respiratory infection, pharyngitis, sinusitis/rhinitis, bronchospasm, dyspnoea. Renal insufficiency. Hypersensitivity reactions, drowsiness, sleep disturbances, nervousness, mood changes, anxiety. Also angioedema, taste disturbances, tinnitus, sweating, flu syndrome, malaise, arthralgia, myalgia. **Pharmaceutical precautions:** Store in a dry place below 25°C. **Legal category:** POM. **Package quantities and prices:** Perdix 7.5mg: calendar packs of 28 tablets £8.50; Perdix 15mg: calendar packs of 28 tablets £9.80. **Product licence numbers:** Perdix 7.5mg – 4438/0033. Perdix 15mg – 4438/0034. **Product licence holder:** Schwarz Pharma Ltd., Schwarz House, East Street, Chesham, Bucks. HP5 1DG. Telephone: 01494 772071. Fax: 01494 773934. **Date of preparation:** September 1995 (389). **Further information is available from the licence holder:** Schwarz Pharma Limited, East Street, Chesham, Bucks. HP5 1DG. **References:** 1. British Heart Foundation, 1995. 2. Data on file 02.

SCHWARZ



PERDIX<sup>®</sup>  
Moexipril