

- report and the review of the literature. *Kyobu Geka* 1993; 46:283-6.
- 25 Kurosawa H, Wagenaar SS, Becker AE. Sudden death in a youth. A case of quadricuspid aortic valve with isolation of origin of left coronary artery. *Br Heart J* 1981;46: 211-15.
  - 26 Peretz DJ, Changfoot GH, Gourlay RH. Four-cusped aortic valve with significant hemodynamic abnormality. *Am J Cardiol* 1969;23:291-3.
  - 27 Wyatt JP, Goldenberg H. Supernumerary aortic cusps with multiple fenestrations and with displacement of the left coronary orifice. *Arch Pathol* 1948;45:784.
  - 28 Kim H-S, McBride RA, Titus JL. Quadricuspid aortic valve and single coronary ostium. *Arch Pathol Lab Med* 1988;112:842-4.
  - 29 Feizi OF, Farrer-Brown G, Emanuel R. Familial study of hypertrophic cardiomyopathy and congenital aortic valve disease. *Am J Cardiol* 1978;41:956-64.
  - 30 Iglesias A, Oliver J, Munoz JE, Nunez L. Quadricuspid aortic valve associated with fibromuscular subaortic stenosis and aortic regurgitation treated by conservative surgery. *Chest* 1981;80:327-8.
  - 31 Fericola DJ, Mann JM, Roberts WC. Congenitally quadricuspid aortic valve: analysis of six necropsy patients. *Am J Cardiol* 1989;63:136-8.
  - 32 Karam R, Lever HM, Healey BP. Hypertensive hypertrophic cardiomyopathy or hypertrophic cardiomyopathy with hypertension? A study of 78 patients. *J Am Coll Cardiol* 1989;13:580-4.
  - 33 McRonald RE, Dean DC. Congenital quadricuspid aortic valve. *Am J Cardiol* 1966;18:761-3.
  - 34 Nalbantgil I, Cagatay G. Quadricuspid aortic valve [letter]. *Chest* 1975;67:623-4.
  - 35 Stassano P, Iorio D, Di Lello F, Tecchia LB, Spampinato N. Aortic insufficiency due to quadricuspid valve. *Minerva Cardioangiol* 1982;30:7-9.
  - 36 Kosach GA, Evzhanov A, Serbin VI, Zubkova GA, Tadzhibaeva ZA. Quadricuspid aortic valve. *Kardiologiya* 1982;22:113-14.
  - 37 Luisi VS, Pasque A, Verunelli F, Aliboni M, Urbano U, Reginato E. Quadricuspid aortic valve. *J Cardiovasc Surg Torino* 1984;25:252-4.
  - 38 Matsumoto M, Miki S, Kusuhara K, Ueda Y, Ohkita Y, Tahata T, Komeda M. Quadricuspid aortic valve associated with severe aortic regurgitation. *Jpn Circ J* 1985;49:190-1.
  - 39 Subramanian R, Olson LJ, Edwards WD. Surgical pathology of combined aortic stenosis and insufficiency: a study of 213 cases. *Mayo Clin Proc* 1985;60:247-54.
  - 40 Waller BF, Taliencio CP, Dickos DK, Howard J, Adlam JH, Jolly W. Rare or unusual causes of chronic, isolated, pure aortic regurgitation. *Clin Cardiol* 1990;13:577-81.
  - 41 Koizumi S, Arai S, Kato T. A case of quadricuspid aortic valve with aortic regurgitation. *Kyobu Geka* 1990;43: 505-8.
  - 42 Fischler D, Fitzmaurice M, Ratliff NB. Quadricuspid aortic valve. *Am J Cardiovasc Pathol* 1990;3:91-4.
  - 43 Shioi K, Nagata Y, Kato S, Kato R, Tsuchioka H. A case of quadricuspid aortic valve associated with coronary arterial lesion. *Nippon Kyobu Geka Gakkai Zasshi* 1991;39: 1072-6.
  - 44 Aoyagi S, Kawara T, Yasunaga H, Kosuga K, Oishi K. Congenital quadricuspid aortic valve associated with aortic regurgitation. *Thorac Cardiovasc Surg* 1992;40:225-6.
  - 45 Irisawa T, Yoshiya K, Yokosawa T, Iwamatsu T, Arai K, Aoki T. A case of quadricuspid aortic valve associated with mitral regurgitation. *Kyobu Geka* 1993;46:618-21.

## IMAGES IN CARDIOLOGY

### Non-invasive detection of quadricuspid aortic valve

Transthoracic echocardiography of a 66 year old Japanese woman with a history of cardiac murmur from early childhood showed large regurgitant blood flow from the aortic valve into the left ventricle that continued until the end of the diastolic phase.

Cine magnetic resonance imaging (MRI) revealed unexpected results: the aortic valve was quadricuspid (normal sized right cusp, slightly smaller left cusp, and half sized two non-coronary cusps without enlargement of annulus). The images of the end diastolic phase (left panel) and the mid-systolic phase

(right panel) showed that these four cusps could not close completely even at the end of the diastolic phase resulting in the formation of a "regurgitant window" (approximately 8 × 8 mm). MRI also showed that her pulmonary valve was tricuspid. This non-invasive examination clearly demonstrated both the presence of a quadricuspid aortic valve and the relation between structure and function of this rare aortic valve anomaly.

KOUJI KAJINAMI  
NOBORU TAKEKOSHI  
HIROSHI MABUCHI

