Left atrial spontaneous echo contrast in patients with permanent pacemakers

Tsung-Ming Lee, Sheng-Fang Su, Kuo-Liong Chien, Yueh-Juh Lin, Wen-Jone Chen, Ming-Fong Chen, Chiau-Suong Liao, Yuan-Teh Lee

Abstract

Objective—To determine the relations between left atrial appendage function, spontaneous echo contrast, and thromboembolism in patients with different modes of permanent pacemakers.

Patients and methods—88 patients with pacemaker implantation and 25 healthy controls in sinus rhythm had transoesophageal echocardiographic examination of the left atrial appendage. Left atrial size, appendage area, peak filling and emptying velocities of the atrial appendage, and the presence or absence of spontaneous echo contrast and thromboembolism were determined. The results in 63 patients with ventricular pacing (group 1, subdivided into subgroup 1A: 42 patients with sinus rhythm, and subgroup 1B: 21 patients with atrial fibrillation) were compared with those in 25 patients with synchronous pacing (group 2), and 25 normal control subjects (group 3).

Results—Patients with ventricular pacing had two distinct appendage flow patterns: well defined biphasic filling and emptying waves in subgroup 1A, and irregular very low filling and emptying waves in subgroup 1B. The ejection fraction of the left atrial appendage in subgroup 1A was significantly better than that in subgroup 1B (mean (SD) 40.6(12.0)% *v* 7.6(5.0)%, $P < 0.0001$). The spontaneous echo contrast was observed in 90% of subgroup 1B patients but in only 19% in subgroup 1A ($P < 0.05$) and was not found in groups 2 and 3 ($P < 0.0001$). There was a trend for increased prevalence of spontaneous echo contrast in subgroup 1A *v* group 2 ($P = 0.053$). Thrombi were detected in two cases, and cardiogenic embolism occurred in one case in subgroup 1B. All patients with spontaneous echo contrast had ventricular pacing. Multivariate analysis showed that atrial fibrillation was associated with occurrence of spontaneous echo contrast in patients with ventricular pacing ($P = 0.005$).

Conclusions—The left atrial appendage ejection fraction was lower with ventricular pacing than with synchronous pacing. With ventricular pacing there was a trend towards increased prevalence of left atrial spontaneous echo contrast in patients in sinus rhythm, and a significantly increased prevalence in patients with atrial fibrillation.

(Heart 1997;78:262-267)

Keywords: echocardiography; left atrial appendage; pacemaker; spontaneous echo contrast

It is well known that patients with ventricular pacing have an increased incidence of thromboembolism.¹⁻³ In recent years synchronous pacing modes of pacemaker (AAI (R), DDD (R), VDD) have been developed to offer alternative options to VVI (R) mode. It is believed that normal atrioventricular conduction is not only physiologically relevant but also clinically important. In comparison with synchronous pacing, ventricular pacing has the disadvantage of the absence of—or random occurrence of—atrial contraction to ventricular filling; thus atrial fibrillation may be induced.⁴⁻⁶ Although the association of atrial fibrillation with thromboembolism is well established,⁷⁻⁹ the mechanism of thromboembolism in patients with ventricular pacing is still not clear.

Spontaneous echo contrast of the left atrium and the left atrial appendage can be demonstrated by transoesophageal echocardiography. The appearance of spontaneous atrial echo contrast is believed to be associated with left atrial appendage thrombus, with increased risks of thromboembolism.^{10,11} The purpose of our study was (1) to assess the effect of synchronous versus ventricular pacing on the function of the left atrial appendage; and (2) to investigate the correlation between the incidence of left atrial spontaneous echo contrast phenomenon and different pacing modes.

Methods

PATIENTS

During a six month period, 92 patients with permanent pacemaker underwent transoesophageal echocardiographic examination. All patients were in sinus rhythm at the time of the pacemaker implantation. There was no pacemaker syndrome and no patients were taking oral anticoagulants. Patients with concomitant valvar disease were not included. Four patients were excluded because of the presence of persistent spontaneous rhythm at the time of echocardiographic study. Only paced beats were included for analysis at the time of echocardiographic study. The remaining 88 patients, with a mean age of 66.6 (SD 11.3) years (range 30 to 88; 50 men, 38 women), were prospectively studied. All

Department of
Internal Medicine,
Cardiology Section,
National Taiwan
University Hospital,
Taipei, Taiwan
T-M Lee
K-L Chien
Y-J Lin
W-J Chen
M-F Chen
C-S Liao
Y-T Lee

Department of Clinical
Pharmacy, National
Cheng Kung
University, Tainan,
Taiwan
S-F Su

Correspondence to:
Dr T-M Lee, Department of
Internal Medicine,
Cardiology Section, National
Taiwan University Hospital,
7 Chung-Shan S Rd, Taipei,
Taiwan 10002, ROC.
email: tmlee@
ha.mc.ntu.edu.tw

Accepted for publication
8 May 1997

Figure 1 Biplane longitudinal transoesophageal echocardiography in a VVI patient. The maximum (A) and minimum (B) areas of appendage were measured by tracing a line starting from the top of the limbus of the left upper pulmonary vein along the entire endocardial border to the anteriormost portion (arrowhead) of the mitral annulus. LA, left atrium.

patients underwent detailed electrophysiological study before pacemaker implantation. Indications for pacemaker implantation included sick sinus syndrome ($n = 67$) and complete atrioventricular block ($n = 21$). A VVI(R) pacemaker was implanted in 63 patients (ventricular group, group 1). The synchronous group (group 2) consisted of 25 patients with pacemaker types AAI (R) ($n = 13$), VDD ($n = 2$), and DDD (R) ($n = 10$). Patients in group 1 were further divided into two subgroups on the basis of cardiac rhythm at the time of the transoesophageal echocardiographic study: subgroup 1A, in sinus rhythm, $n = 42$; subgroup 1B, in atrial fibrillation, $n = 21$. The clinical characteristics of the patients are given in table 1. For comparison, an additional group was enrolled in the study: this control group (group 3) consisted of 25 normal subjects in sinus rhythm and no apparent heart disease.

ECHOCARDIOGRAPHY

All patients were examined by transthoracic and transoesophageal echocardiography on the same day. M mode echocardiography was performed in all patients and left atrial size was determined. Transoesophageal echocardiographic examination was performed using a 5 MHz biplane ($n = 87$; 35 subgroup 1A, 17 subgroup 1B, 16 group 2, 19 group 3) or multiplane transducer ($n = 26$). Patients had

fasted for six hours before transoesophageal echocardiography. Local pharyngeal anaesthesia with 8% lignocaine spray was the only pre-medication. During echocardiography, a one lead electrocardiogram was recorded. Informed consent was obtained from all patients and normal control subjects. The study was approved by the institutional research ethics committee.

Left atrial appendage area and flow

Images of the left atrial appendage were obtained in the transverse and longitudinal planes and recorded on Super VHS videotapes for off-line analysis. The boundary of the base of the appendage was defined by a line drawn from the limbus of the left upper pulmonary vein to the anteriormost portion of the mitral annulus. Maximum and minimum left atrial appendage areas were determined by computed planimetry along the endocardial border of the appendage (average of three consecutive values) (fig 1). The ejection fraction of the left atrial appendage was calculated as (maximum area – minimum area)/maximum area. Left atrial appendage velocity profiles were obtained by pulsed wave Doppler interrogation at the orifice of the appendage. The maximum forward positive flow velocity of Doppler left atrial appendage represented the peak emptying velocity (fig 2, *a* wave) and the maximum backward negative flow velocity repre-

Figure 2 Velocity profiles in VVI patients with preserved sinus activities (subgroup 1A), (A) with ventriculoatrial conduction and (B) without ventriculoatrial conduction. Both had an underlying disease of sick sinus syndrome. Both show left atrial appendage flow patterns. Regular and well defined emptying (*a*) waves regularly follow atrial electric activity (seen in A, but not seen in B). Filling (*b*) velocity waves follow the emptying wave. A respiratory trace is shown at the bottom.

Figure 3 Velocity profiles in a VVI patient with atrial fibrillation (subgroup 1B), an underlying disease of sick sinus syndrome. Left atrial appendage flow patterns show irregular low emptying and filling waves. A respiratory trace is shown at the bottom.

Figure 4 Velocity profiles in a patient with DDD pacemaker (group 2), an underlying disease of complete atrioventricular block. Left atrial appendage flow patterns show well defined emptying (a wave) after electrocardiographically antegrade P wave and filling (b wave) velocities following the a wave. A respiratory trace is shown at the bottom.

sented the peak filling velocity (fig 2, b wave). There were 25 patients in subgroup 1A with quadriphasic flows (c and d waves). Because of their smaller flow velocities, c and d waves did not affect the measurement of peak filling and emptying flow velocities. The ejection fraction of the left atrial appendage and the peak emptying and filling velocities were averaged with each RR interval over a minimum of five cardiac cycles for patients with atrial fibrillation, and over three cardiac cycles for those in sinus rhythm. Interobserver differences were resolved by consensus.

Spontaneous echo contrast and thrombus
Left atrial spontaneous echo contrast was diagnosed by the presence of dynamic smoke-like echoes within the atrial cavity, with a characteristic swirling motion distinct from white noise artefact.¹² The gain was continuously adjusted to ensure the best possible visualisation and to avoid noise artefact. Spontaneous echo contrast was graded as previously described¹³: grade 0 = no contrast; grade 1 = slight contrast localised in parts of the left atrium with a low density; grade 2 = heavy contrast with intense, swirling, smoke-like echoes in the whole atrium. Thrombi were defined as masses adherent to the wall of the left atrial appendage with different echogenic density. Particular attention was paid to differentiating these from pectinate muscles. The presence or absence of spontaneous echo contrast and thrombi was defined by the consensus of two experienced echocardiographers.

STATISTICS
Values are reported as mean (SD). A χ^2 test was used to compare categorical variables. The continuous variables between groups were compared with Student's *t* test for two tailed unpaired observation. For comparison of multiple groups, analysis of variance was applied. For the identification of independent predictors of spontaneous echo contrast a multivariate regression analysis was used. Two tailed Fisher's exact test was used for patient numbers less than five. A P value < 0.05 was considered statistically significant.

Results
Figures 2, 3, and 4 show typical findings of left atrial appendage flow in subgroups 1A and 1B and group 2.

BASELINE VARIABLES
The selection of pacing mode was not randomised but was dependent on patients' age, physical activity, prognosis, and cost. The echo studies were performed at a mean time from implantation of 48 (52) months for subgroup 1A, 54 (47) months for subgroup 1B, and 12 (10) months for group 2. The average follow up time in subgroups 1A and 1B was much longer than in group 2. The patient age

Table 1 Characteristics of patients and left atrial appendage in different groups

	Group 1		Group 2 (n = 25)	Control (n = 25)
	1A (n = 42)	1B (n = 21)		
Age (years)	66.4 (10.9)*	72.4 (7.1)‡	62.1 (12.9)	64.6 (7.9)
Male/female	25/17	11/10	14/11	15/10
SSS/CAVB	32/10	19/2‡	16/9	—
Set heart rate (beats/min)	66.4 (4.8)†	66.7 (4.8)‡	72.9 (5.2)	62.3 (6.8)¶
Follow up period	48 (52)†	54 (47)‡	12 (10)	—
Left atrial size (mm)	36.5 (3.5)*	46.3 (4.6)‡	34.2 (3.2)	34.8 (4.2)
Left atrial appendage parameters				
Maximum area (cm ²)	5.4 (1.4)*	6.1 (1.4)‡	4.7 (1.3)	4.9 (2.1)
Minimum area (cm ²)	3.2 (1.0)*†	5.6 (1.2)‡	2.6 (1.2)	2.8 (1.6)
Ejection fraction (%)	40.6 (12.0)*†	7.6 (5.0)‡	48.2 (16.2)	45.7 (13.6)
Peak filling velocity (cm/s)	51.2 (19.6)*	10.8 (8.6)‡	59.8 (17.6)	52.3 (16.5)
Peak emptying velocity (cm/s)	49.3 (17.8)*	11.4 (6.5)‡	51.8 (18.3)	53.8 (23.1)
Spontaneous echo contrast	8 (19%)*	19 (90%)‡	0	0
Thrombus	0	2 (9.5%)	0	0

Values are number (%) of patients or mean (SD).
*P < 0.05 v subgroup 1B; †P < 0.05 v group 2; ‡P < 0.05 v group 2; ¶P < 0.05 v group 2.
CAVB, complete atrioventricular block; SSS, sick sinus syndrome.

in subgroup 1B was significantly higher than in subgroup 1A and group 2, at 72.4 (7.1) *v* 66.4 (10.9) and 62.1 (12.9) (both $P < 0.05$). The set heart rate of the pacemakers was significantly higher in group 2 than in subgroups 1A and 1B. The size of the left atrium was significantly greater in subgroup 1B than in subgroups 1A and groups 2 and 3, at 46.3 (4.6) *v* 36.5 (3.5), 34.2 (3.2), and 34.8 (4.2) mm (all $P < 0.05$) (table 1).

LEFT ATRIAL APPENDAGE

In all patients in subgroup 1A, the left atrial appendage flow revealed an organised biphasic flow pattern with a peak filling and emptying velocity (fig 2). In contrast, all patients from subgroup 1B showed an irregular, very low peak filling and emptying velocity (fig 3). Table 1 compares the left atrial appendage transoesophageal echocardiography parameters in the three groups. The maximum appendage areas were significantly larger in subgroup 1B than in subgroup 1A and group 2. More striking is the difference in the left atrial appendage ejection fraction, which was significantly reduced in subgroups 1A and 1B. There were almost no visible appendage contractions in subgroup 1B. There were no significant differences of the ejection fraction of the left atrial appendage detected by biplane transoesophageal echocardiography compared with multiplane transoesophageal echocardiography in subgroup 1A (41.2 (11.8) *v* 39.4 (12.4) %, NS) (table 1).

Two patterns of left atrial appendage flow were identified in group 1. In subgroup 1A and groups 2 and 3, Doppler image revealed a biphasic flow pattern with a peak filling and emptying velocity. The *a* wave occurred after the P wave and the *b* wave followed the *a* wave. In contrast, patients from subgroup 1B showed irregular, very low peak filling veloci-

ties (range 3 to 25 cm/s, mean 10.8 (8.6) cm/s) and peak emptying velocities (3 to 30 cm/s, mean 11.4 (6.5) cm/s). The left atrial appendage peak filling and emptying velocities were significantly reduced in patients from subgroup 1B compared with patients from subgroups 1A, 2, and 3. There were no significant differences in variable echocardiographic parameters between groups 2 and 3.

SPONTANEOUS ECHO CONTRAST AND THROMBUS FORMATION

The spontaneous echo contrast phenomenon was not observed in groups 2 and 3; however, it was present in 90% of subgroup 1B patients. The echo contrast was seen in eight patients (19%) of subgroup 1A. In subgroup 1B, left atrial spontaneous echo contrast was graded as slight in 12 patients (63%) and heavy in seven (36%). In subgroup 1A, all left atrial spontaneous echo contrast was graded as slight. There was a trend of an increased prevalence of spontaneous echo contrast in subgroup 1A compared with in group 2, although this was not statistically significant ($P = 0.053$). However, the statistic power of the study was 0.45 for $\alpha = 0.05$. Two patients from subgroup 1B showed a left atrial appendage thrombus; one experienced ischaemic stroke. The frequency of atrial fibrillation was higher ($P < 0.0001$) in patients with echo contrast than in those without echo contrast (table 2), as were the minimum and maximum atrial appendage areas ($P < 0.001$ and $P = 0.02$, respectively). The ejection fraction of the atrial appendage was lower in patients with echo contrast than in patients without (42.6 (14.6) *v* 15.5 (15.3) %, $P < 0.0001$). The peak filling and emptying velocities for all patients with echo contrast were lower than for patients without echo contrast (both $P < 0.0001$).

Although univariate analysis showed that older age, ventricular pacing, atrial fibrillation, left atrial size, and maximum/minimum areas and peak filling/emptying velocities of the left atrial appendage were significantly related to occurrence of spontaneous echo contrast (table 2), multivariate analysis identified atrial fibrillation as the variable associated with a significantly increased risk of left atrial spontaneous echo contrast in patients with ventricular pacing (table 3). The value of -2 log L (likelihood ratio test) compared with the full model was 3.00, $df = 1$ ($P = NS$). All patients with spontaneous echo contrast had ventricular pacing. Ventricular pacing was an independent determinant in the univariate analysis of spontaneous echo contrast.

Table 2 Univariate analysis: clinical and echocardiographic factors related to left atrial spontaneous echo contrast (SEC) in pacemaker patients

	SEC (–) (n = 61)	SEC (+) (n = 27)	P value
Age	64.6 (11.5)	72.1 (7.7)	0.003
Male/female	36/25	14/13	NS
SSS/CAVB	43/18	24/3	NS
Ventricular/synchronous	36/25	27/0	<0.0001
Atrial fibrillation	2 (3%)	19 (70%)	<0.0001
Set heart rate	69.0 (6.0)	66.3 (4.9)	0.049
Left atrial appendage	35.4 (4.2)	44.5 (3.7)	<0.0001
Left atrial appendage parameters			
Maximum area (cm ²)	5.2 (1.5)	5.9 (1.3)	0.02
Minimum area (cm ²)	3.0 (1.2)	5.0 (1.5)	<0.0001
Ejection fraction (%)	42.6 (14.6)	15.5 (15.3)	<0.0001
Peak filling velocity (cm/s)	53.2 (19.2)	20.6 (22.4)	<0.0001
Peak emptying velocity (cm/s)	48.2 (18.2)	22.9 (23.4)	<0.0001
Thrombus	0	2 (7%)	NS

Values are number (%) of patients or mean (SD).

SSS, sick sinus syndrome; CAVB, complete atrioventricular block.

Table 3 Logistic regression model analysis of risk factors for development of spontaneous echo contrast in patients with ventricular pacing

Predictors	Odds ratio	95% confidence interval	P value
Atrial fibrillation	21.7	2.52 to 187.21	0.005
Sick sinus syndrome	1.8	0.47 to 25.42	0.447
LA size	1.6	0.55 to 12.73	0.364
LAA peak emptying velocity	0.8	0.83 to 1.05	0.142
LAA ejection fraction	15.8	0.02 to 9874.31	0.372

LA, left atrium; LAA, left atrial appendage.

Discussion

Our study is the first study to show that patients with ventricular pacing have a high prevalence of impaired left atrial appendage function and the spontaneous echo contrast phenomenon in the left atrium, especially in the subgroup with concomitant atrial fibrillation. For patients with ventricular pacing and sinus rhythm, there is a tendency toward an increased prevalence of spontaneous echo

contrast, but this was not statistically significant ($P = 0.053$) because of the small number of patients in this subgroup (power = 0.45).

LEFT ATRIAL APPENDAGE

Left atrial appendage velocities and flow patterns vary according to rhythm.¹⁴ In sinus rhythm, biphasic or quadriphasic clearly defined emptying and filling flows associated with appendage contraction and relaxation were observed. The *a* wave, representing forward flow towards the left atrial cavity, was associated with left atrial appendage contraction and began from the initial deflection of the P wave on the electrocardiogram. The *b* wave, representing retrograde flow toward the left atrial appendage, was associated with re-expansion of the left atrial appendage and began immediately after the *a* wave. Some patients with sinus rhythm had quadriphasic flow patterns in the left atrial appendage. The *c* and *d* waves are passive flows. The *c* wave, coinciding in timing with early mitral inflow, was associated with a pressure gradient between the left atrial appendage and the left ventricle on opening of the mitral valve. The *d* wave, occurring just after the *c* wave but before appendage contraction, indicated left atrial appendage filling during the diastolic period of mitral inflow. Because the *c* and *d* waves were significantly lower in velocity, they did not affect measurement of peak filling and emptying velocities.

In atrial fibrillation, a disorganised flow pattern and reduced velocities of emptying and filling flows are seen. Mugge *et al*⁷ meticulously divided patients with non-rheumatic atrial fibrillation into two subgroups on the basis of left atrial appendage function. One group had a high flow profile with high peak filling and emptying velocities of the left atrial appendage; the other had a low flow profile with very low peak filling and emptying velocities. Their results show that the incidence of spontaneous echo contrast was significantly increased in patients with low left atrial appendage flow than in those with high flow. Porte *et al*¹⁵ found that left atrial appendage contractile function was a predictor of spontaneous echo contrast. Because the appendage patterns were of the low flow profile type with impaired contractile function in our subgroup 1B, it was expected that there would be a high incidence of spontaneous echo contrast. Our results are consistent with previous studies^{8,16,17} that patients with low flow profiles of the left atrial appendage have a high incidence of spontaneous echo contrast.

SPONTANEOUS ECHO CONTRAST AND THROMBUS FORMATION

The left atrial spontaneous echo contrast phenomenon was significantly more frequent in the ventricular group with atrial fibrillation. One study showed that patients with left atrial spontaneous echo contrast are 27 times more likely to have had a previous stroke or peripheral embolism than those without echo contrast.¹⁸ Daniel *et al*⁹ found that in patients with mitral valve disease, left atrial echo contrast

was an independent predictor of left atrial thrombus and cardiogenic embolism. However, only one patient developed cerebral embolism in our study, although there was a higher prevalence of spontaneous echo contrast in the ventricular group. Because only surviving patients were examined and the loss of some patients with severe stroke cannot be ruled out, the real incidence of thromboembolism is probably higher than shown by our study.

MULTIVARIATE PREDICTORS OF SPONTANEOUS ECHO CONTRAST

In this study population, atrial fibrillation was significantly associated with left atrial spontaneous echo contrast in patients with ventricular pacing. The left atrial size, underlying sick sinus syndrome, and left atrial appendage parameters were not predictors of spontaneous echo contrast in patients with ventricular pacing. These results agree with previous observations^{7,11}, that atrial fibrillation is associated with an increased risk of developing spontaneous echo contrast. However, the relation between ventricular pacing and spontaneous echo contrast has not previously been reported. Our results showed that there was a prevalence of spontaneous echo contrast in patients with ventricular pacing: 19% in patients with preserved atrial activity and 90% in patients with atrial fibrillation. One explanation for the high prevalence of spontaneous echo contrast is that ventricular pacing results in a dilated left atrium¹⁹ and relatively reduced cardiac output.²⁰ Conditions favouring stasis of left atrial blood, including left atrial enlargement and low cardiac output, were associated with left atrial spontaneous echo contrast. Therefore ventricular pacing, even with sinus rhythm, was associated with a trend towards the occurrence of spontaneous echo contrast.

CLINICAL IMPLICATIONS

The effect of ventricular pacing and atrial fibrillation on left atrial appendage function and left atrial spontaneous echo contrast suggests a hidden benefit of synchronous pacing. Left atrial spontaneous echo contrast is a phenomenon that usually appears in regions of blood stasis including an enlarged left atrium, as in this study; however, the mechanism of this phenomenon remains unclear. The pathogenesis of spontaneous echo contrast is complex and includes not only the velocity or shear rate of local blood flow but also factors such as abnormalities of blood components. Siegel *et al*²¹ suggested that rouleaux formation of erythrocytes and increased level of serum fibrinogen may be responsible. However, Erbel *et al*²² found increased platelet aggregation in all their patients. Mahony *et al*²³ reported a patient with left ventricular spontaneous echo contrast despite heparin treatment, in whom platelet aggregates were detected in the peripheral blood. Complete spontaneous echo contrast resolution was noted after five days of antiplatelet treatment. Sasaki *et al*²⁴ significantly reduced the incidence of stroke by using anticoagulants in patients with ventricular

pacing. Thus further studies are warranted comparing the therapeutic value of anticoagulant and antiplatelet drugs in paced patients with spontaneous echo contrast.

STUDY LIMITATIONS

This study has limitations. First, it was not randomised. The selection of pacemaker modes depended on patient age, the preference of primary cardiologists, and the cost. Second, the number of patients with ventricular pacing and sinus rhythm was small, which precluded any definite conclusion about the relation between ventricular pacing and spontaneous echo contrast. Third, estimation of the ejection fraction of the left atrial appendage can be influenced by movement of the transoesophageal probe, which will affect the echo plane in which the atrial appendage is evaluated and may cause overestimation or underestimation of the appendage ejection fraction. However, this did not seem to be a problem because this variable was averaged over a minimum of five cardiac cycles in patients with atrial fibrillation and over three cardiac cycles in those in sinus rhythm. Fourth, the rarity of thromboembolic events in this series of patients precluded a definite correlation between systemic embolism and the proposed mechanism involved in thromboembolism in patients with ventricular pacing. Fifth, the presence or absence of spontaneous echo contrast is subjective. However, interpretations were by two experienced cardiologists and any differences were resolved by consensus, so we feel there was little likelihood of misinterpretation. Finally, although some basic clinical variables were not considered in the multivariate analysis, this will not affect the conclusions because all variables with a P value < 0.05 in univariate analysis were considered. Values of χ^2 for covariates compared with the full model were used to check the differences of $-2 \log L$ of various models and to select the optimum model for interpretation.

CONCLUSIONS

Patients with ventricular pacing had a decreased ejection fraction of the left atrial appendage compared with patients with synchronous pacing. With ventricular pacing, there was a trend towards increased prevalence of left atrial spontaneous echo contrast in patients in sinus rhythm, and a significantly high prevalence in patients with atrial fibrillation.

We thank Yuh-Chen Huang for statistical assistance and advice.

- 1 Langenfeld H, Grimm W, Maisch B, Kochsiek K. Atrial fibrillation and embolic complications in paced patients. *PACE* 1988;11:1667-72.
- 2 Bathen J, Sparr S, Rosketh E. Embolism in sinoatrial dis-

- ease. *Acta Med Scand* 1978;203:7-11.
- 3 Santini M, Alexidou G, Ansalone G, Cacciato G, Cini R, Turitto G. Relation of prognosis in sick sinus syndrome to age, conduction defects and modes of permanent cardiac pacing. *Am J Cardiol* 1990;65:729-35.
- 4 Sutton R, Kenny R-A. The natural history of sick sinus syndrome. *PACE* 1986;9:1110-13.
- 5 Reimold SC, Lamas GA, Cantillon CO, Antman EM. Risk factors for development of recurrent atrial fibrillation: role of pacing and clinical variables. *Am Heart J* 1995;129:1127-32.
- 6 Santiago D, Warshofsky M, Mandri GL, Di Tullio M, Coromilas J, Reiffel J, et al. Left atrial appendage function and thrombus formation in atrial fibrillation-flutter: a transesophageal echocardiographic study. *J Am Coll Cardiol* 1994;24:159-64.
- 7 Mugge A, Kuhn H, Nikutta P, Grote J, Lopez AG, Daniel WG. Assessment of left atrial appendage function by biplane transesophageal echocardiography in patients with nonrheumatic atrial fibrillation: identification of a subgroup of patients at increased embolic risk. *J Am Coll Cardiol* 1994;23:599-607.
- 8 Verhorst PMJ, Kamp O, Visser CA, Verheugt FWA. Left atrial appendage flow velocity assessment using transesophageal echocardiography in nonrheumatic atrial fibrillation and systemic embolism. *Am J Cardiol* 1993;71:192-6.
- 9 Daniel WG, Nellessen U, Schroder E, Nonnast-Daniel B, Bednarski P, Nikutta P, et al. Left atrial spontaneous contrast in mitral valve disease: an indicator for an increased thromboembolic risk. *J Am Coll Cardiol* 1988;11:1204-11.
- 10 Black IW, Hopkins AP, Lee LCL, Walsh WF, Jacobson BM. Left atrial spontaneous echo contrast: a clinical and echocardiographic analysis. *J Am Coll Cardiol* 1991;18:398-404.
- 11 De Belder MA, Lovat LB, Tourikis L, Leech G, Camm A. Left atrial spontaneous contrast echoes: markers of thromboembolic risk in patients with atrial fibrillation. *Eur Heart J* 1993;14:326-35.
- 12 Castello R, Pearson AC, Labovitz AJ. Prevalence and clinical implications of atrial spontaneous contrast in patients undergoing transesophageal echocardiography. *Am J Cardiol* 1990;65:1149-53.
- 13 Cormier B, Vahanian A, Iung B, Porte JM, Dadez E, Lazarus A, et al. Influence of percutaneous mitral commissurotomy on left atrial spontaneous contrast of mitral stenosis. *Am J Cardiol* 1993;71:842-7.
- 14 Jue J, Winslow T, Fazio G, Redberg RF, Foster E, Schiller NB. Pulsed Doppler characterization of left atrial appendage flow. *J Am Soc Echocardiogr* 1993;6:237-44.
- 15 Porte JM, Cormier B, Iung B, Dadez E, Starkman C, Nallet O, et al. Early assessment by transesophageal echocardiography of left atrial appendage function after percutaneous mitral commissurotomy. *Am J Cardiol* 1996;77:72-6.
- 16 Asanuma T, Tanabe K, Yoshitomi H, Shimizu H, Murakami Y, Sano Y, et al. Left atrial appendage in patients with single-chamber ventricular pacing. *Am J Cardiol* 1995;76:840-2.
- 17 Fatkin D, Kelly RP, Feneley MP. Relations between left atrial blood velocity, spontaneous echocardiographic contrast and thromboembolic risk in vivo. *J Am Coll Cardiol* 1994;23:961-9.
- 18 Chimowitz MI, DeGeorgia MA, Poole RM, Hepner A, Armstrong WM. Left atrial spontaneous echo contrast is highly associated with previous stroke in patients with atrial fibrillation or mitral stenosis. *Stroke* 1993;24:1015-9.
- 19 Kubica J, Stolarczyk L, Krzyminska E, Krasowski R, Raczak G, Lubinski A, et al. Left atrial size and wall motion in patients with permanent ventricular and atrial pacing. *PACE* 1990;13:1737-41.
- 20 Leclercq C, Gras D, Helloc AL, Nicol L, Mabo P, Daubert C. Hemodynamic importance of preserving the normal sequence of ventricular activation in permanent cardiac pacing. *Am Heart J* 1995;129:1133-41.
- 21 Sigel B, Coelho JC, Spigos DG, Flanagan DP, Schuler JJ, Kasprisin DO, et al. Ultrasonography of blood during stasis and coagulation. *Invest Radiol* 1981;16:71-6.
- 22 Erbel R, Stern H, Ehrenthal W, Schreiner G, Treese N, Kramer G, et al. Detection of spontaneous echocardiographic contrast within the left atrium by transesophageal echocardiography: spontaneous echocardiographic contrast. *Clin Cardiol* 1986;9:245-52.
- 23 Mahony C, Sublett KL, Harrison MR. Resolution of spontaneous contrast with platelet disaggregatory therapy (trifluoperazine). *Am J Cardiol* 1989;63:1009-10.
- 24 Sasaki S, Takeuchi A, Ohzeki M, et al. Long-term follow-up of paced patients with sick sinus syndrome. In: Steinbach K et al, eds. *Cardiac Pacing. Proceedings of the Seventh World Symposium on cardiac Pacing*. Darmsadt: Steinklppft Verlag, 1983:85-90.