

Subject index

- 22q11 microdeletion**, Common arterial trunk in the fetus: characteristics, associations, and outcome in a multicentre series of 23 cases, **1437**
- ablation**, Ablate and pace revisited: long term survival and predictors of permanent atrial fibrillation, **1035**
- aborted myocardial infarction**, Abortion of acute ST segment elevation myocardial infarction after reperfusion: incidence, patients' characteristics, and prognosis, **496**
- ACE genotype**, Prospective evaluation of the effect of an angiotensin I converting enzyme gene polymorphism on the long term risk of major adverse cardiac events after percutaneous coronary intervention, **321**
- ACE inhibitors**, A rational basis for selection among drugs of the same class, **687**
- acquired fistula**, Acquired right coronary artery fistula draining to the right ventricle: angiographic documentation of first appearance following reperfusion after acute myocardial infarction, with subsequent spontaneous closure, **e22**
- acquired immunodeficiency virus**, HIV and cardiovascular medicine, **793**
- active-fixation permanent pacing**, Prolonged temporary cardiac pacing using subcutaneous tunnelled active-fixation permanent pacing leads, **209**
- acute aortic syndrome**, Acute aortic syndrome: proposal for a novel classification, **928**
- acute coronary syndromes**, A single serum glucose measurement predicts adverse outcomes across the whole range of acute coronary syndromes, **512**
- Are troponins confusing?, **1285**
- Associations between ST depression, four year mortality, and in-hospital revascularisation in unselected patients with non-ST elevation acute coronary syndromes, **490**
- C reactive protein for risk stratification in acute coronary syndromes? Verdict: unproven, **1288**
- Creatinine clearance and adverse hospital outcomes in patients with acute coronary syndromes: findings from the global registry of acute coronary events (GRACE), **1003**
- Effect of diabetes on serum potassium concentrations in acute coronary syndromes, **31**
- Multicentre evaluation of the diagnostic value of cardiac troponin T, CK-MB mass, and myoglobin for assessing patients with suspected acute coronary syndromes in routine clinical practice, **280**
- Plaque stabilisation in acute coronary syndromes: clinical considerations, **1268**
- Predictors of mortality in patients with acute coronary syndrome undergoing percutaneous coronary intervention, **1245**
- Revascularisation for acute coronary syndromes: PCI or CABG?, **967**
- The primacy of clinical effectiveness for cost effectiveness analysis, **249**
- acute myocardial infarction**, Acquired right coronary artery fistula draining to the right ventricle: angiographic documentation of first appearance following reperfusion after acute myocardial infarction, with subsequent spontaneous closure, **e22**
- Are we underestimating the full potential of early thrombolytic treatment in patients with acute myocardial infarction?, **483**
- Assessing perfusion and function in acute myocardial infarction: how and when?, **701**
- Assessment of myocardial viability: using coronary pressure and flow after acute myocardial infarction, **9**
- Bangladeshi patients present with non-classic features of acute myocardial infarction and are treated less aggressively in east London, UK, **276**
- Carbohydrates, dietary glycaemic load and glycaemic index, and risk of acute myocardial infarction, **722**
- Emergency coronary stenting of unprotected critical left main coronary artery stenosis in acute myocardial infarction and cardiogenic shock, **e24**
- GREAT: 10 year survival of patients with suspected acute myocardial infarction in a randomised comparison of prehospital and hospital thrombolysis, **563**
- Impact of microvascular integrity and local viability on left ventricular remodelling after reperfused acute myocardial infarction, **393**
- Myocardial contrast echocardiography is superior to other known modalities for assessing myocardial reperfusion after acute myocardial infarction, **139**
- Prospective evaluation of the role of routine cardiac troponin T measurement to identify left ventricular ejection fraction < 40% after first myocardial infarction, **559**
- Relation of coronary flow pattern to myocardial blush grade in patients with first acute myocardial infarction, **1147**
- Role of myocardial contrast echocardiography in the clinical evaluation of acute myocardial infarction, **1398**
- The variation of plasma concentrations of a novel, adipocyte derived protein, adiponectin, in patients with acute myocardial infarction, **667**
- Trends in incidence and case fatality rates of acute myocardial infarction in Denmark and Sweden, **507**
- Usefulness of myocardial perfusion SPECT in patients with left bundle branch block and previous myocardial infarction, **1039**
- Variation during the week in the incidence of acute myocardial infarction: increased risk for Japanese women on Saturdays, **398**
- acute ventricular septal rupture**, Optimal management of acute ventricular septal rupture, **1462**
- adiponectin**, Adipocyte derived plasma protein, adiponectin, is associated with smoking status in patients with coronary artery disease, **663**
- The variation of plasma concentrations of a novel, adipocyte derived protein, adiponectin, in patients with acute myocardial infarction, **667**
- adolescents**, Asymptomatic ventricular pre-excitation in children and adolescents: a 15 year follow up study, **215**
- β adrenoceptor blocker**, Different responses to dobutamine in the presence of carvedilol or metoprolol in patients with chronic heart failure, **621**
- adult**, Cor triatriatum sinister presenting in the adult as mitral stenosis, **e26**
- adult congenital heart disease**, Challenges in the care of adult patients with congenital heart defects, **465**
- The adult patient with native coarctation of the aorta: balloon angioplasty or primary stenting?, **77**
- adverse events**, Adverse events in coronary artery bypass graft (CABG) trials: a systematic review and analysis, **767**
- Flecainide induced ventricular fibrillation in a neonate, **1261**
- adverse outcomes**, A single serum glucose measurement predicts adverse outcomes across the whole range of acute coronary syndromes, **512**
- Air pollution**, Why cardiologists should be interested in air pollution, **1383**
- ALLHAT**, Drug treatment of hypertension: implications of ALLHAT, **589**
- alteplase**, Thrombolysis: newer thrombolytic agents and their role in clinical medicine, **1358**
- ambulatory blood pressure monitoring**, Ambulatory blood pressure monitoring in the management of hypertension, **571**
- ambulatory electrocardiography**, Novel use of an "insertable" loop recorder, **e18**
- amino-terminal pro brain natriuretic peptide**, The influence of age, sex and other variables on the plasma level of N-terminal pro brain natriuretic peptide in a large sample of the general population, **745**
- Amplatzer septal occluder**, Experience with transcatheter closure of secundum atrial septal defects using the Amplatzer septal occluder: a single centre study in 236 consecutive patients, **199**
- Amplatzer VSD occluder**, Transcatheter closure of perimembranous ventricular septal defects with the Amplatzer asymmetric ventricular septal defect occluder: preliminary experience in children, **918**
- aneurysm**, Low pressure giant pulmonary artery aneurysms in the adult: natural history and management strategies, **1067**
- Open surgery for thoracic aortic disease, **825**
- Unruptured congenital aneurysm of the left sinus of Valsalva in an adult with complex left heart malformations, **e3**

- angina**, Same day discharge following elective percutaneous coronary intervention in patients with stable angina, **665**
Successful brachytherapy of coronary vasospasm, **e25**
- angina pectoris**, Enhanced external counterpulsation for refractory angina pectoris, **830**
The current cost of angina pectoris to the National Health Service in the UK, **848**
- angiogenesis**, Angiogenesis, thrombogenesis, endothelial dysfunction and angiographic severity of coronary artery disease, **1411**
Decreased serum vascular endothelial growth factor concentrations in patients with congestive heart failure, **207**
The human coronary collateral circulation, **1352**
- angiography**, How many cath labs do we need?, **827**
Invasive assessment of myocardial bridges, **699**
Long term effects of nisoldipine on the progression of coronary atherosclerosis and the occurrence of clinical events: the NICOLE study, **887**
Prognostic and clinical correlates of angiographically diffuse non-obstructive coronary lesions, **1009**
- angioplasty**, In-stent stenosis: pathology and implications for the development of drug eluting stents, **218**
Mechanism of lumen enlargement with direct stenting versus predilatation stenting: influence of remodelling and plaque characteristics assessed by volumetric intracoronary ultrasound, **84**
Prospective evaluation of the effect of an angiotensin I converting enzyme gene polymorphism on the long term risk of major adverse cardiac events after percutaneous coronary intervention, **321**
Same day discharge following elective percutaneous coronary intervention in patients with stable angina, **665**
The CARDia trial protocol, **1125**
- angiosarcoma**, Unexplained recurrent pericardial effusion: a lethal warning?, **e11**
- angiotensin blockers**, A rational basis for selection among drugs of the same class, **687**
- angiotensin converting enzyme**, DD ACE gene polymorphism is associated with increased coronary artery endothelial dysfunction: the PREFACE trial, **557**
Lack of association between angiotensin converting enzyme (ACE) genotype, serum ACE activity, and haemodynamics in patients with primary pulmonary hypertension, **445**
- angiotensin converting enzyme inhibitors**, A rational basis for selection among drugs of the same class, **687**
Plaque stabilisation in acute coronary syndromes: clinical considerations, **1268**
- angiotensin I converting enzyme gene**, Deletion polymorphism of the angiotensin I converting enzyme gene is a potent risk factor for coronary artery ectasia, **213**
- angiotensin II receptor blocker**, Angiotensin II receptor blockade prevents microangiopathy and preserves diastolic function in the diabetic rat heart, **1236**
- antiarrhythmics**, Drug induced QT prolongation and torsades de pointes, **1363**
- antibiotic treatment**, Prosthetic valve endocarditis: early and late outcome following medical or surgical treatment, **269**
- antidepressants**, Drug induced QT prolongation and torsades de pointes, **1363**
- antiepileptic drug**, Differential diagnosis of cardiogenic syncope and seizure disorders, **353**
- antihistamines**, Drug induced QT prolongation and torsades de pointes, **1363**
- antimicrobials**, Drug induced QT prolongation and torsades de pointes, **1363**
- antiplatelet treatment**, Antiplatelet treatment in stable coronary artery disease, **1273**
- aorta**, Dissection of the aorta: a new approach, **6**
- aortic arch pathology**, Aortic valve and aortic arch pathology after coarctation repair, **1074**
- aortic atresia**, Concordance for hypoplastic left heart syndrome in a monozygotic twin pregnancy, **e13**
- aortic coarctation**, Challenges in the care of adult patients with congenital heart defects, **465**
Coarctation of the aorta in adults: do we need surgeons?, **3**
The adult patient with native coarctation of the aorta: balloon angioplasty or primary stenting?, **77**
The efficacy and safety of stent redilatation in congenital heart disease, **905**
- aortic dissection**, Endovascular treatment of thoracic aortic disease, **823**
- aortic regurgitation**, Excessive pulmonary autograft dilatation causes important aortic regurgitation, **933**
- aortic stenosis**, Cardiovascular risk factors in patients with aortic stenosis predict prevalence of coronary artery disease but not of aortic stenosis: an angiographic pair matched case-control study, **1019**
- aortic valve**, Three dimensional volume quantification of aortic valve calcification using multislice computed tomography, **1191**
- aortic valve disease**, Calcific aortic stenosis: from bench to the bedside—emerging clinical and cellular concepts, **801**
- aortic valve replacement**, Risk factors for pacemaker implantation following aortic valve replacement: a single centre experience, **901**
- aphrodisiac**, Toad venom poisoning: resemblance to digoxin toxicity and therapeutic implications, **e14**
- apical ballooning**, Apical ballooning of the left ventricle: first series in white patients, **1027**
- apoptosis**, Relation between circulating soluble Fas ligand and subsequent ventricular remodelling following myocardial infarction, **339**
- arrhythmia**, Management of fetal tachyarrhythmia based on superior vena cava/aorta Doppler flow recordings, **1211**
QRS duration: a simple marker for predicting cardiac mortality in ICD patients with heart failure, **1157**
- arrhythmias**, Asymptomatic ventricular pre-excitation in children and adolescents: a 15 year follow up study, **215**
- arterial switch operation**, Transoesophageal atrial pacing combined with transthoracic two dimensional echocardiography: experience in patients operated on with arterial switch operation for transposition of the great arteries, **91**
- arterial trunks**, Development of the heart: (1) Formation of the cardiac chambers and arterial trunks, **806**
- arterial valves, intrapericardial arterial trunks**, Development of the heart: (3) Formation of the ventricular outflow tracts, arterial valves, and intrapericardial arterial trunks, **1110**
- arteriogenesis**, The human coronary collateral circulation, **1352**
- ascending aorta**, Aortic valve and aortic arch pathology after coarctation repair, **1074**
- aspirin**, The central role of platelet activation in determining the severity of acute coronary syndromes, **1253**
- asynchrony**, High prevalence of left ventricular systolic and diastolic asynchrony in patients with congestive heart failure and normal QRS duration, **54**
Interventricular and intra-left ventricular electromechanical delays in right ventricular paced patients with heart failure: implications for upgrading to biventricular stimulation, **1401**
- atherosclerosis**, Arterial dysfunction in syndrome X: results of arterial reactivity and pulse wave propagation tests, **422**
Calcific aortic stenosis: from bench to the bedside—emerging clinical and cellular concepts, **801**
Impact of preprocedural white blood cell count on long term mortality after percutaneous coronary intervention: insights from the EPIC, EPILOG, and EPISTENT trials, **1200**
Improving the positive predictive value of exercise testing in women, **1416**
Long term effects of nisoldipine on the progression of coronary atherosclerosis and the occurrence of clinical events: the NICOLE study, **887**
On-line detection of cholesterol and calcification by catheter based Raman spectroscopy in human atherosclerotic plaque ex vivo, **1078**
Prognostic and clinical correlates of angiographically diffuse non-obstructive coronary lesions, **1009**
- athlete's heart**, Athlete's heart, **1455**
- atrial fibrillation**, Ablate and pace revisited: long term survival and predictors of permanent atrial fibrillation, **1035**
Atrial fibrillation: classification, pathophysiology, mechanisms and drug treatment, **939**
Atrial fibrillation: relation between clinical risk factors and transoesophageal echocardiographic risk factors for thromboembolism, **165**
Comparison of monophasic and biphasic shocks for transthoracic cardioversion of atrial fibrillation, **1032**
Frequent atrial premature contractions as a surrogate marker for paroxysmal atrial fibrillation in patients with acute ischaemic stroke, **1247**
Off-pump coronary artery bypass graft surgery: the incidence of postoperative atrial fibrillation, **1134**

- Pacing for atrial fibrillation, **106**
- Platelet activation, haemorrhology and thrombogenesis in acute atrial fibrillation: a comparison with permanent atrial fibrillation, **1093**
- Role of the autonomic nervous system in vagal atrial fibrillation, **333**
- The role of echocardiography in atrial fibrillation and cardioversion, **1447**
- Warfarin for non-valvar atrial fibrillation: still underused in the 21st century?, **553**
- atrial flutter**, Review of diagnosis, treatment, and outcome of fetal atrial flutter compared with supraventricular tachycardia, **913**
- atrial premature contractions**, Frequent atrial premature contractions as a surrogate marker for paroxysmal atrial fibrillation in patients with acute ischaemic stroke, **1247**
- atrial septal aneurysm**, Learning on the web: Case 2: Patent foramen ovale (PFO) and paradoxical embolism, **1362**
- atrial septal defect**, Challenges in the care of adult patients with congenital heart defects, **465**
- atriopulmonary connection**, Fontan conversion to total cavopulmonary connection and arrhythmia ablation: clinical and functional results, **193**
- atrioventricular node**, The relative contribution of the sinus and AV node to heart rate variability, **337**
- atria**, Development of the heart: (2) Septation of the atria and ventricles, **949**
- atypical Fabry's disease**, Sisters with atypical Fabry's disease with complete atrioventricular block, **e2**
- automated implantable cardioverter defibrillator**, Comparative follow up of patients with implanted cardioverter-defibrillators after induction of sustained monomorphic ventricular tachycardias or ventricular fibrillation by programmed stimulation, **629**
- autonomic agents**, Acetylcholinesterase inhibition with pyridostigmine improves heart rate recovery after maximal exercise in patients with chronic heart failure, **854**
- autonomic nervous system**, Acetylcholinesterase inhibition with pyridostigmine improves heart rate recovery after maximal exercise in patients with chronic heart failure, **854**
- Role of the autonomic nervous system in vagal atrial fibrillation, **333**
- Why cardiologists should be interested in air pollution, **1383**
- autosomal recessive disorder**, Incidence of pericardial effusion during attacks of familial Mediterranean fever, **1257**
- B type natriuretic peptide**, B Type natriuretic peptide: a good omen in myocardial ischaemia?, **707**
- Normal values of B type natriuretic peptide in infants, children, and adolescents, **875**
- balloon dilatation**, The adult patient with native coarctation of the aorta: balloon angioplasty or primary stenting?, **77**
- bed rest**, Position and Mobilisation Post-Angiography Study (PAMPAS): a comparison of 4.5 hours and 2.5 hours bed rest, **447**
- bicuspid valve**, Aortic valve and aortic arch pathology after coarctation repair, **1074**
- Bicuspid aortic valve and coarctation: two villains part of a diffuse problem, **965**
- biodegradation**, Biocorrosion of magnesium alloys: a new principle in cardiovascular implant technology?, **651**
- bioprosthesis**, Which heart valve prosthesis for patients aged between 60 and 70 years?, **481**
- bioprosthetic valves**, The Edinburgh heart valve study, **697**
- bisoprolol**, Obstacles to the initiation of β blockers for heart failure in a specialised clinic within a district general hospital, **442**
- biventricular pacing**, Colour tissue velocity imaging can show resynchronisation of longitudinal left ventricular contraction pattern by biventricular pacing in patients with severe heart failure, **859**
- Interventricular and intra-left ventricular electromechanical delays in right ventricular paced patients with heart failure: implications for upgrading to biventricular stimulation, **1401**
- Restrictive transmitral filling patterns predict improvements in left ventricular function after biventricular pacing, **1087**
- β blockers**, A rational basis for selection among drugs of the same class, **687**
- Importance of β blockade in the treatment of advanced heart failure, **1442**
- blood culture negative endocarditis**, Blood culture negative endocarditis: analysis of 63 cases presenting over 25 years, **258**
- blood flow**, Assessment of myocardial viability using coronary zero flow pressure after successful angioplasty in patients with acute anterior myocardial infarction, **71**
- blood oxygen level dependent magnetic resonance imaging**, Detection of scarred and viable myocardium using a new magnetic resonance imaging technique: blood oxygen level dependent (BOLD) MRI, **738**
- blood pressure**, Exercise systolic blood pressures are of questionable value in the assessment of the adult with a previous coarctation repair, **189**
- blunt chest trauma**, Diagnosing cardiac contusion: old wisdom and new insights, **485**
- BNP**, B Type natriuretic peptide: a good omen in myocardial ischaemia?, **707**
- brachytherapy**, Histopathology of coronary in-stent restenosis following γ brachytherapy, **11**
- Successful brachytherapy of coronary vasospasm, **e25**
- brain**, Cardiac surgery and the brain: differences between adult and paediatric studies, **365**
- brain natriuretic peptide**, Brain natriuretic peptide facilitates severity classification of stable chronic heart failure with left ventricular dysfunction, **661**
- Circulating concentrations of cardiac proteins indicate the severity of congestive heart failure, **1303**
- Screening for left ventricular dysfunction in the community: role of hand held echocardiography and brain natriuretic peptides, **iii24**
- breathlessness**, Pattern of ventilation during exercise in chronic heart failure, **610**
- Bruce**, Persisting fever in a patient with brucella endocarditis: occult splenic abscess, **e20**
- Brugada syndrome**, Abnormal response to sodium channel blockers in patients with Brugada syndrome: augmented localised wall motion abnormalities in the right ventricular outflow tract region detected by electron beam computed tomography, **169**
- Obstacles to the initiation of β blockers for heart failure in a specialised clinic within a district general hospital, **442**
- Relation between heart rate, heart rhythm, and reverse left ventricular remodelling in response to carvedilol in patients with chronic heart failure: a single centre, observational study, **293**
- Treatment of chronic heart failure with β adrenergic blockade beyond controlled clinical trials: the BRING-UP experience, **299**
- C reactive protein**, C reactive protein for risk stratification in acute coronary syndromes? Verdict: unproven, **1288**
- calcific aortic stenosis**, Calcific aortic stenosis: from bench to the bedside—emerging clinical and cellular concepts, **801**
- calcium**, Three dimensional volume quantification of aortic valve calcification using multislice computed tomography, **1191**
- calcium channel**, The therapeutic potential of new insights into myocardial excitation-contraction coupling, **371**
- calcium channel blocker**, Catecholaminergic polymorphic ventricular tachycardia: electrocardiographic characteristics and optimal therapeutic strategies to prevent sudden death, **66**
- Campbell**, Maurice Campbell: first editor of *Heart*, **1379**
- capillaries**, Rarefaction of skin capillaries in normotensive offspring of individuals with essential hypertension, **175**
- carbohydrate intake**, Carbohydrates, dietary glycaemic load and glycaemic index, and risk of acute myocardial infarction, **722**
- cardiac arrest**, Factors associated with survival to hospital discharge among patients hospitalised alive after out of hospital cardiac arrest: change in outcome over 20 years in the community of Göteborg, Sweden, **25**
- Presentation, management, and outcome of out of hospital cardiopulmonary arrest: comparison by underlying aetiology, **839**
- cardiac arrhythmia**, Ablate and pace revisited: long term survival and predictors of permanent atrial fibrillation, **1035**
- The therapeutic potential of new insights into myocardial excitation-contraction coupling, **371**
- cardiac catheterisation**, Position and Mobilisation Post-Angiography Study (PAMPAS): a comparison of 4.5 hours and 2.5 hours bed rest, **447**
- cardiac chambers**, Development of the heart: (1) Formation of the cardiac chambers and arterial trunks, **806**
- cardiac contusion**, Diagnosing cardiac contusion: old wisdom and new insights, **485**
- cardiac death**, Long QTc predicts future cardiac death in stroke survivors, **377**
- cardiac function**, Athlete's heart, **1455**
- cardiac imaging**, Contrast echocardiography: from left ventricular opacification to myocardial perfusion. Are the promises to be realised?, **1389**

- cardiac related events**, Accuracy of reporting cardiac related events during long term follow up, **331**
- cardiac remodelling**, B Type natriuretic peptide: a good omen in myocardial ischaemia?, **707**
- cardiac sarcoidosis**, A 28 year old woman with ventricular tachycardia and an abnormal chest radiograph, **e19**
- cardiac troponins T and I**, Are troponins confusing?, **1285**
Multicentre evaluation of the diagnostic value of cardiac troponin T, CK-MB mass, and myoglobin for assessing patients with suspected acute coronary syndromes in routine clinical practice, **280**
Prospective evaluation of the role of routine cardiac troponin T measurement to identify left ventricular ejection fraction - 40% after first myocardial infarction, **559**
- cardiac tumours**, Primary cardiac valve tumours, **1259**
- cardiogenic shock**, Emergency coronary stenting of unprotected critical left main coronary artery stenosis in acute myocardial infarction and cardiogenic shock, **e24**
- cardiomyopathy**, Mutations of the γ myosin heavy chain gene in hypertrophic cardiomyopathy: critical functional sites determine prognosis, **1179**
Risk of competitive sport in young athletes with heart disease, **710**
Risk progression to chronic Chagas cardiomyopathy: influence of male sex and of parasitaemia detected by polymerase chain reaction, **1186**
- cardioplegia**, Protecting the damaged heart during coronary surgery, **367**
- cardiopulmonary resuscitation**, Presentation, management, and outcome of out of hospital cardiopulmonary arrest: comparison by underlying aetiology, **839**
- cardiovascular disease**, Diabetes: can we stop the time bomb?, **ii28**
Ethnic differences in cardiovascular disease, **681**
Microarray analysis: a novel research tool for cardiovascular scientists and physicians, **597**
The obesity epidemic: can we turn the tide?, **ii22**
What level of physical activity protects against premature cardiovascular death? The Caerphilly study, **502**
Why cardiologists should be interested in air pollution, **1383**
- cardiovascular evaluation**, Risk of competitive sport in young athletes with heart disease, **710**
- cardiovascular risk**, Role of exercise treadmill testing in the management of erectile dysfunction: a joint cardiovascular/erectile dysfunction clinic, **671**
- cardioversion**, Comparison of monophasic and biphasic shocks for transthoracic cardioversion of atrial fibrillation, **1032**
The role of echocardiography in atrial fibrillation and cardioversion, **1447**
- carotid stenosis**, Carotid artery stenting, **944**
- carvedilol**, Relation between heart rate, heart rhythm, and reverse left ventricular remodelling in response to carvedilol in patients with chronic heart failure: a single centre, observational study, **293**
- cased based learning**, Cased based learning on the web, **979**
- catecholamines**, Haemodynamic and catecholamine response to simulated ventricular tachycardia in man: effect of baseline left ventricular function, **306**
- catheter ablation**, Catheter ablation of sinoatrial re-entry tachycardia in a 2 month old infant, **e1**
Successful radiofrequency catheter ablation of "clockwise" and "counterclockwise" bundle branch re-entrant ventricular tachycardia in the absence of myocardial or valvar dysfunction without detecting bundle branch potentials, **e12**
- catheter laboratories**, How many cath labs do we need?, **827**
- catheterisation**, Percutaneous technique for the reduction of knotted coronary catheters, **1132**
- CD14 polymorphism**, CD14 promoter polymorphism is associated with acute myocardial infarction resulting from insignificant coronary artery stenosis, **931**
- cell proliferation**, In-stent stenosis: pathology and implications for the development of drug eluting stents, **218**
- cerebral blood flow**, Syncope: what is the trigger?, **123**
- cerebral dysfunction**, Neuroprotection during cardiac surgery: a randomised trial of a platelet activating factor antagonist, **897**
- Chagas' disease**, Risk progression to chronic Chagas cardiomyopathy: influence of male sex and of parasitaemia detected by polymerase chain reaction, **1186**
- chemosensitivity**, Attenuation of hypercapnic carbon dioxide chemosensitivity after postinfarction exercise training: possible contribution to the improvement in exercise hyperventilation, **404**
- children**, Asymptomatic ventricular pre-excitation in children and adolescents: a 15 year follow up study, **215**
Heart rate variability in children with hypertrophic cardiomyopathy, **205**
Left atrial and left ventricular function in healthy children and young adults assessed by three dimensional echocardiography, **544**
Normal values of B type natriuretic peptide in infants, children, and adolescents, **875**
- cholesterol**, High preprocedural non-HDL cholesterol is associated with enhanced oxidative stress and monocyte activation after coronary angioplasty: possible implications in restenosis, **773**
Lipid lowering to delay the progression of coronary artery disease, **451**
- cholesterol crystal embolisation**, Trash feet after coronary angiography, **e17**
- chromosomal anomalies**, Tricuspid regurgitation in the diagnosis of chromosomal anomalies in the fetus at 11–14 weeks of gestation, **1071**
- chronic cor pulmonale**, Chronic cor pulmonale, **225**
- chronic heart failure**, Pattern of ventilation during exercise in chronic heart failure, **610**
The effects of α and β blockade on ventilatory responses to exercise in chronic heart failure, **1169**
- chronic obstructive pulmonary disease**, Chronic cor pulmonale, **225**
- chronic thromboembolic pulmonary hypertension**, Pulmonary thromboendarterectomy almost 50 years after the first surgical attempts, **369**
- classification**, Acute aortic syndrome: proposal for a novel classification, **928**
- clinical targets**, Current targets: where are we going?, **ii6**
Current targets: where are we today?, **ii10**
- clinical usefulness**, Clinical utility and cost effectiveness of a personal ultrasound imager for cardiac evaluation during consultation rounds in patients with suspected cardiac disease, **727**
- clopidogrel**, Antiplatelet treatment in stable coronary artery disease, **1273**
- coaptation point**, Increased distance between mitral valve coaptation point and mitral annular plane: significance and correlations in patients with heart failure, **1174**
- coarctation**, Bicuspid aortic valve and coarctation: two villains part of a diffuse problem, **965**
Exercise systolic blood pressures are of questionable value in the assessment of the adult with a previous coarctation repair, **189**
- coarctation repair**, Aortic valve and aortic arch pathology after coarctation repair, **1074**
Exercise blood pressure in congenital heart disease and in patients after coarctation repair, **125**
- cognitive impairment**, Neuroprotection during cardiac surgery: a randomised trial of a platelet activating factor antagonist, **897**
- cohort study**, Is maternal transmission of coronary heart disease risk stronger than paternal transmission?, **834**
What level of physical activity protects against premature cardiovascular death? The Caerphilly study, **502**
- cold pressor test**, Regional myocardial perfusion defects during exercise, as assessed by three dimensional integration of morphology and function, in relation to abnormal endothelium dependent vasoreactivity of the coronary microcirculation, **517**
- collagen**, Interstitial fibrosis in the dilated non-ischaemic myocardium, **1255**
- collateral circulation**, The human coronary collateral circulation, **1352**
- collateral flow**, Tumour necrosis factor α concentration and collateral flow in patients with coronary artery disease and normal systolic left ventricular function, **96**
- colour tissue velocity imaging**, Colour tissue velocity imaging can show resynchronisation of longitudinal left ventricular contraction pattern by biventricular pacing in patients with severe heart failure, **859**
- common arterial trunk**, Common arterial trunk in the fetus: characteristics, associations, and outcome in a multicentre series of 23 cases, **1437**
- competing risks analysis**, Mode of death in heart failure: findings from the ATLAS trial, **42**
- competitive sports**, Risk of competitive sport in young athletes with heart disease, **710**

- complete atrioventricular block**, Sisters with atypical Fabry's disease with complete atrioventricular block, **e2**
- complete atrioventricular septal defect**, Williams syndrome associated with complete atrioventricular septal defect, **e15**
- complications**, Coronary artery aneurysm with a fistulous connection to the right atrium mimicking a sinus of Valsalva aneurysm, **e4**
Pregnancy related complications in women with hypertrophic cardiomyopathy, **752**
- computed tomography**, Abnormal response to sodium channel blockers in patients with Brugada syndrome: augmented localised wall motion abnormalities in the right ventricular outflow tract region detected by electron beam computed tomography, **169**
Non-invasive coronary imaging: computed tomography or magnetic resonance imaging?, **591**
Three dimensional volume quantification of aortic valve calcification using multislice computed tomography, **1191**
- conduction**, Usefulness of myocardial perfusion SPECT in patients with left bundle branch block and previous myocardial infarction, **1039**
- conduit arterial stiffness**, Conduit arterial stiffness is associated with impaired left ventricular subendocardial function, **449**
- congenital cardiovascular malformation**, Cardiovascular malformations in infants of diabetic mothers, **1217**
- congenital coronary artery dilatation**, Congenital coronary artery dilatation, **595**
- congenital cyanotic heart disease**, Long term somatic growth after repair of tetralogy of Fallot: evidence for restoration of genetic growth potential, **1340**
- congenital heart disease**, Acute endothelin A receptor antagonism improves pulmonary and systemic haemodynamics in patients with pulmonary arterial hypertension that is primary or autoimmune and related to congenital heart disease, **1221**
Cardiac surgery and the brain: differences between adult and paediatric studies, **365**
Coarctation of the aorta in adults: do we need surgeons?, **3**
Cor triatriatum sinister presenting in the adult as mitral stenosis, **e26**
Exercise blood pressure in congenital heart disease and in patients after coarctation repair, **125**
Neurodevelopment at 1 year of age in infants with congenital heart disease, **436**
Pericardial effusion after open heart surgery for congenital heart disease, **780**
Single coronary artery: a familial clustering, **e27**
- congenitally corrected transposition**, Impaired myocardial blood flow and coronary flow reserve of the anatomical right systemic ventricle in patients with congenitally corrected transposition of the great arteries, **1231**
- congestive heart failure**, Decreased serum vascular endothelial growth factor concentrations in patients with congestive heart failure, **207**
Enhanced endothelin-1 degradation by intravenous morphine in patients with congestive heart failure: role of neutral endopeptidase 24.11, **211**
Mode of death in heart failure: findings from the ATLAS trial, **42**
Serum magnesium aberrations in furosemide (frusemide) treated patients with congestive heart failure: pathophysiological correlates and prognostic evaluation, **411**
Trends and ethnic differences in hospital admissions and mortality for congestive heart failure in the elderly in Singapore, 1991 to 1998, **865**
- contrast echocardiography**, Contrast echocardiography, **342**
Contrast echocardiography for left ventricular opacification, **1394**
Contrast echocardiography: from left ventricular opacification to myocardial perfusion. Are the promises to be realised?, **1389**
Isolated left ventricular non-compaction: cardiomyopathy with homogeneous transmural and heterogeneous segmental perfusion, **e21**
Myocardial contrast echocardiography is superior to other known modalities for assessing myocardial reperfusion after acute myocardial infarction, **139**
- contrast media**, Temporal evolution and functional outcome of no reflow: sustained and spontaneously reversible patterns following successful coronary recanalisation, **731**
- coronary angiography**, A prospective study of long term prognosis in young myocardial infarction survivors: the prognostic value of angiography and exercise testing, **843**
Apical ballooning of the left ventricle: first series in white patients, **1027**
Mitral annular calcification: a marker of severe coronary artery disease in patients under 65 years old, **161**
Percutaneous technique for the reduction of knotted coronary catheters, **1132**
Trash feet after coronary angiography, **e17**
- coronary angioplasty**, Coronary angioplasty enhances platelet reactivity through von Willebrand factor release, **329**
- coronary anomalies**, Anomalous origin of the circumflex artery and patent foramen ovale: a rare cause of myocardial ischaemia after percutaneous closure of the defect, **e23**
- coronary anomaly**, Single coronary artery: a familial clustering, **e27**
- coronary arteries**, Invasive assessment of myocardial bridges, **699**
- coronary artery aneurysm**, Coronary artery aneurysm with a fistulous connection to the right atrium mimicking a sinus of Valsalva aneurysm, **e4**
- coronary artery bypass graft surgery**, Adverse events in coronary artery bypass graft (CABG) trials: a systematic review and analysis, **767**
Coronary stenting versus coronary bypass surgery in patients with multiple vessel disease and significant proximal LAD stenosis: results from the ERACI II study, **184**
Indications for percutaneous and surgical revascularisation: how far does the evidence base guide us?, **565**
Neuroprotection during cardiac surgery: a randomised trial of a platelet activating factor antagonist, **897**
Simvastatin attenuates leucocyte-endothelial interactions after coronary revascularisation with cardiopulmonary bypass, **538**
Validation of four different risk stratification systems in patients undergoing off-pump coronary artery bypass surgery: a UK multicentre analysis of 2223 patients, **432**
Revascularisation for acute coronary syndromes: PCI or CABG?, **967**
Socioeconomic deprivation is a predictor of poor postoperative cardiovascular outcomes in patients undergoing coronary artery bypass grafting, **1062**
- coronary artery disease**, Adipocyte derived plasma protein, adiponectin, is associated with smoking status in patients with coronary artery disease, **663**
Angiogenesis, thrombogenesis, endothelial dysfunction and angiographic severity of coronary artery disease, **1411**
Antiplatelet treatment in stable coronary artery disease, **1273**
Cardiovascular effects of testosterone: implications of the "male menopause", **121**
Cardiovascular risk factors in patients with aortic stenosis predict prevalence of coronary artery disease but not of aortic stenosis: an angiographic pair matched case-control study, **1019**
Coronary flow velocity analysis during short term follow up after coronary reperfusion: use of transthoracic Doppler echocardiography to predict regional wall motion recovery in patients with acute myocardial infarction, **1163**
Cost effectiveness of extended treatment with low molecular weight heparin (dalteparin) in unstable coronary artery disease: results from the FRISC II trial, **287**
Drug eluting stents: are human and animal studies comparable?, **133**
Effects of L-arginine on flow mediated dilatation induced by atrial pacing in diseased epicardial coronary arteries, **531**
Enhanced external counterpulsation for refractory angina pectoris, **830**
Epistatic interaction between variations in the angiotensin I converting enzyme and angiotensin II type 1 receptor genes in relation to extent of coronary atherosclerosis, **1195**
Lipid lowering to delay the progression of coronary artery disease, **451**
Non-invasive intravenous coronary angiography using electron beam tomography and multislice computed tomography, **633**
Prognostic and clinical correlates of angiographically diffuse non-obstructive coronary lesions, **1009**
Prospective evaluation of the effect of an angiotensin I converting enzyme gene polymorphism on the long term risk of major adverse cardiac events after percutaneous coronary intervention, **321**
Quality of life assessment using the short form 12 questionnaire is as reliable and sensitive as the short form 36 in distinguishing symptom severity in myocardial infarction survivors, **1445**
Role of myocardial perfusion imaging for risk stratification in suspected or known coronary artery disease, **1291**
Same day discharge following elective percutaneous coronary intervention in patients with stable angina, **665**
Stress echocardiography, **113**

- Tumour necrosis factor α concentration and collateral flow in patients with coronary artery disease and normal systolic left ventricular function, **96**
- coronary artery ectasia**, Deletion polymorphism of the angiotensin I converting enzyme gene is a potent risk factor for coronary artery ectasia, **213**
- coronary artery fistula**, Acquired right coronary artery fistula draining to the right ventricle: angiographic documentation of first appearance following reperfusion after acute myocardial infarction, with subsequent spontaneous closure, **e22**
- coronary artery spasm**, Increased platelet sodium–hydrogen exchanger activity in patients with variant angina, **935**
- coronary artery stenosis**, CD14 promoter polymorphism is associated with acute myocardial infarction resulting from insignificant coronary artery stenosis, **931**
- coronary blood flow**, Coronary flow: clinical considerations, **361**
Persistence of systolic coronary flow reversal predicts irreversible dysfunction after reperfused anterior myocardial infarction, **382**
- coronary calcification**, Coronary calcifications in young patients with first, unheralded myocardial infarction: a risk factor matched analysis by electron beam tomography, **625**
- coronary circulation**, The human coronary collateral circulation, **1352**
- coronary flow**, Relation of coronary flow pattern to myocardial blush grade in patients with first acute myocardial infarction, **1147**
- coronary flow reserve**, Myocardial viability, coronary flow reserve, and in-hospital predictors of late recovery of contractility following successful primary stenting for acute myocardial infarction, **179**
- coronary heart disease**, Are there sex differences in risk factors for coronary heart disease? Maternal versus paternal transmission, **817**
Coronary risk factors predicting early and late coronary deaths, **19**
Current targets: where are we going?, **ii6**
Current targets: where are we today?, **ii10**
Global risk of cardiovascular disease, **ii2**
Improving the delivery of coronary care for ethnic minorities, **479**
Improving the positive predictive value of exercise testing in women, **1416**
Is maternal transmission of coronary heart disease risk stronger than paternal transmission?, **834**
The applicability of the Framingham coronary heart disease prediction function to black and minority ethnic groups in the UK, **785**
The nurse as expert practitioner in global cardiovascular risk management, **ii33**
- coronary microembolisation**, Coronary microembolisation: introduction, **980**
Pathology of coronary microembolisation and no reflow, **983**
Pathophysiology of coronary microembolisation, **981**
Spontaneous and interventional coronary microembolisation, **986**
- coronary ostial lesions**, Clinical and angiographic outcome of directional atherectomy followed by stent implantation in de novo lesions located at the ostium of the left anterior descending coronary artery, **1050**
- coronary risk factors**, Coronary risk factors predicting early and late coronary deaths, **19**
- coronary spasm**, Successful brachytherapy of coronary vasospasm, **e25**
- coronary stenosis**, The elusive link between stenosis severity and prognosis in stable ischaemic heart disease, **961**
- coronary surgery**, Coronary surgery with non-cardioplegic methods in patients with advanced left ventricular dysfunction: immediate and long term results, **427**
Protecting the damaged heart during coronary surgery, **367**
The risk model of choice for coronary surgery in the UK, **98**
- cost effectiveness**, Clinical utility and cost effectiveness of a personal ultrasound imager for cardiac evaluation during consultation rounds in patients with suspected cardiac disease, **727**
Continued improvement of clinical outcome and cost effectiveness following intravascular ultrasound guided PCI: insights from a prospective, randomised study, **1043**
Cost effectiveness of extended treatment with low molecular weight heparin (dalteparin) in unstable coronary artery disease: results from the FRISC II trial, **287**
- cost effectiveness analysis**, The primacy of clinical effectiveness for cost effectiveness analysis, **249**
- cost evaluation**, The current cost of angina pectoris to the National Health Service in the UK, **848**
- creatinine kinase isoenzyme MB**, Circulating concentrations of cardiac proteins indicate the severity of congestive heart failure, **1303**
- creatinine clearance**, Creatinine clearance and adverse hospital outcomes in patients with acute coronary syndromes: findings from the global registry of acute coronary events (GRACE), **1003**
- cryotherapy catheter ablation**, Comparison of radiofrequency versus cryotherapy catheter ablation of septal accessory pathways, **1091**
- culture negative endocarditis**, Diagnosis of culture negative endocarditis: novel strategies to prove the suspect guilty, **241**
- definition**, Sources of diagnostic inaccuracy of conventional versus new diagnostic criteria for myocardial infarction in an unselected UK population with suspected cardiac chest pain, and investigation of independent prognostic variables, **1406**
- deletion polymorphism**, Deletion polymorphism of the angiotensin I converting enzyme gene is a potent risk factor for coronary artery ectasia, **213**
- Denmark**, Trends in incidence and case fatality rates of acute myocardial infarction in Denmark and Sweden, **507**
- depression**, The psychological perspective: a professional view, **ii16**
- diabetes**, Angiotensin II receptor blockade prevents microangiopathy and preserves diastolic function in the diabetic rat heart, **1236**
Cardiovascular malformations in infants of diabetic mothers, **1217**
Diabetes: can we stop the time bomb?, **ii28**
Effect of diabetes on serum potassium concentrations in acute coronary syndromes, **31**
The CARDia trial protocol, **1125**
The obesity epidemic: can we turn the tide?, **ii22**
- diagnosis**, Sources of diagnostic inaccuracy of conventional versus new diagnostic criteria for myocardial infarction in an unselected UK population with suspected cardiac chest pain, and investigation of independent prognostic variables, **1406**
- diagnostic criteria**, Blood culture negative endocarditis: analysis of 63 cases presenting over 25 years, **258**
- diagnostic protocol**, Diagnosing cardiac contusion: old wisdom and new insights, **485**
- diamorphine**, Safety and potential benefits of low dose diamorphine during exercise in patients with chronic heart failure, **1085**
- diastole**, Cardiac function and obesity, **1127**
Clinical assessment of left ventricular diastolic function, **231**
Diastolic heart failure: fact or fiction?, **1281**
- diastolic function**, Effects of isolated obesity on systolic and diastolic left ventricular function, **1152**
- digoxin**, Effects of digoxin on left atrial function in heart failure, **1308**
Management of fetal tachyarrhythmia based on superior vena cava/aorta Doppler flow recordings, **1211**
- digoxin specific Fab fragments**, Toad venom poisoning: resemblance to digoxin toxicity and therapeutic implications, **e14**
- dilated cardiomyopathy**, HIV and cardiovascular medicine, **793**
Progression of familial and non-familial dilated cardiomyopathy: long term follow up, **757**
- dilated myocardium**, Interstitial fibrosis in the dilated non-ischaemic myocardium, **1255**
- directional atherectomy**, Clinical and angiographic outcome of directional atherectomy followed by stent implantation in de novo lesions located at the ostium of the left anterior descending coronary artery, **1050**
- dissection**, Dissection of the aorta: a new approach, **6**
Open surgery for thoracic aortic disease, **825**
- diuretics**, Drug treatment of hypertension: implications of ALLHAT, **589**
- DNA**, Microarray analysis: a novel research tool for cardiovascular scientists and physicians, **597**
- dobutamine**, Diagnostic accuracy of a new shorter dobutamine infusion protocol in stress echocardiography, **1089**
- dobutamine echocardiography**, Effects of glucose–insulin–potassium infusion on chronic ischaemic left ventricular dysfunction, **61**
- dobutamine stress echocardiography**, Early detection of left ventricular dysfunction in patients with β thalassaemia major by dobutamine stress echocardiography, **669**
- Doppler**, Portable spectral Doppler echocardiographic device: overcoming limitations, **1014**
- Doppler echocardiography**, A clinical approach to the assessment of left ventricular diastolic function by Doppler echocardiography: update 2003, **iii18**
Left ventricular remodelling, and systolic and diastolic function in young adults with β thalassaemia major: a Doppler echocardiographic assessment and correlation with haematological data, **762**

- dose reduction**, Radiation exposure to cardiologists: how it could be reduced, **1123**
- dosimetry**, Radiation exposure benefit of a lead cap in invasive cardiology, **1205**
- drug eluting stents**, Drug eluting stents: are human and animal studies comparable?, **133**
Drug eluting stents: maximising benefit and minimising cost, **127**
- drug induced QT prolongation**, Drug induced QT prolongation and torsades de pointes, **1363**
- drugs**, Drugs and professional interactions: the modern day pharmacist, **ii31**
- ductus arteriosus**, Stenting of the ductus arteriosus and banding of the pulmonary arteries: basis for various surgical strategies in newborns with multiple left heart obstructive lesions, **645**
- Duke criteria**, Diagnosis of culture negative endocarditis: novel strategies to prove the suspect guilty, **241**
- echocardiography**, A 28 year old woman with ventricular tachycardia and an abnormal chest radiograph, **e19**
Cardiac function and obesity, **1127**
Clinical criteria and the appropriate use of transthoracic echocardiography for the exclusion of infective endocarditis, **273**
Coronary flow velocity analysis during short term follow up after coronary reperfusion: use of transthoracic Doppler echocardiography to predict regional wall motion recovery in patients with acute myocardial infarction, **1163**
Effects of isolated obesity on systolic and diastolic left ventricular function, **1152**
Fabry disease, **819**
High prevalence of left ventricular systolic and diastolic asynchrony in patients with congestive heart failure and normal QRS duration, **54**
Increased distance between mitral valve coaptation point and mitral annular plane: significance and correlations in patients with heart failure, **1174**
Management of ischaemic mitral regurgitation, **459**
Mitral annular calcification: a marker of severe coronary artery disease in patients under 65 years old, **161**
Persistence of systolic coronary flow reversal predicts irreversible dysfunction after reperfused anterior myocardial infarction, **382**
Portable spectral Doppler echocardiographic device: overcoming limitations, **1014**
The role of echocardiography in atrial fibrillation and cardioversion, **1447**
Ultrasound stethoscopy: a renaissance of the physical examination?, **971**
- elderly**, Which heart valve prosthesis for patients aged between 60 and 70 years?, **481**
- electrocardiography**, A meta-analysis comparing the prognostic accuracy of six diagnostic tests for predicting perioperative cardiac risk in patients undergoing major vascular surgery, **1327**
Abnormal response to sodium channel blockers in patients with Brugada syndrome: augmented localised wall motion abnormalities in the right ventricular outflow tract region detected by electron beam computed tomography, **169**
Athlete's heart, **1455**
High prevalence of left ventricular systolic and diastolic asynchrony in patients with congestive heart failure and normal QRS duration, **54**
- electromagnetic interference**, Artefact mimicking tachycardia during magnetic resonance imaging in a patient with an implantable loop recorder, **e10**
- electron beam tomography**, Coronary calcifications in young patients with first, unheralded myocardial infarction: a risk factor matched analysis by electron beam tomography, **625**
Non-invasive intravenous coronary angiography using electron beam tomography and multislice computed tomography, **633**
- embolic protection devices**, Embolic protection devices, **990**
- embolism**, Incidence of embolism and paravalvar leak after St Jude Silzone valve implantation: experience from the Cardiff Embolic Risk Factor Study, **1055**
- embolotherapy**, Prevalence of haemoptysis in adults with pulmonary atresia and ventricular septal defect, and the role of mammary artery collateral vessels, **937**
- endocarditis**, Clinical criteria and the appropriate use of transthoracic echocardiography for the exclusion of infective endocarditis, **273**
Diagnosis of culture negative endocarditis: novel strategies to prove the suspect guilty, **241**
HIV and cardiovascular medicine, **793**
- Innocent victim of a localised outbreak: legionella endocarditis, **e16**
- endoluminal repair**, Endovascular treatment of thoracic aortic disease, **823**
- endothelial dysfunction**, DD ACE gene polymorphism is associated with increased coronary artery endothelial dysfunction: the PREFACE trial, **557**
Erectile dysfunction and the cardiovascular patient: endothelial dysfunction is the common denominator, **251**
- endothelial function**, Arterial dysfunction in syndrome X: results of arterial reactivity and pulse wave propagation tests, **422**
- endothelin antagonism**, Acute endothelin A receptor antagonism improves pulmonary and systemic haemodynamics in patients with pulmonary arterial hypertension that is primary or autoimmune and related to congenital heart disease, **1221**
- endothelin-1**, Enhanced endothelin-1 degradation by intravenous morphine in patients with congestive heart failure: role of neutral endopeptidase 24.11, **211**
- endothelium**, Angiogenesis, thrombogenesis, endothelial dysfunction and angiographic severity of coronary artery disease, **1411**
Effects of L-arginine on flow mediated dilatation induced by atrial pacing in diseased epicardial coronary arteries, **531**
Regional myocardial perfusion defects during exercise, as assessed by three dimensional integration of morphology and function, in relation to abnormal endothelium dependent vasoreactivity of the coronary microcirculation, **517**
- endovascular stents**, Coarctation of the aorta in adults: do we need surgeons?, **3**
- enhanced external counterpulsation**, Enhanced external counterpulsation for refractory angina pectoris, **830**
- enoxaparin**, Experience with enoxaparin in patients with mechanical heart valves who must withhold acenocumarol, **527**
- entrainment**, Successful radiofrequency catheter ablation of "clockwise" and "counterclockwise" bundle branch re-entrant ventricular tachycardia in the absence of myocardial or valvar dysfunction without detecting bundle branch potentials, **e12**
- epidemiology**, A "natural experiment" in cardiovascular epidemiology in the early 21st century, **255**
Heart failure and the aging population: an increasing burden in the 21st century?, **49**
Prevalence of impaired left ventricular systolic function and heart failure in a middle aged and elderly urban population segment of Copenhagen, **1422**
Prognosis for patients newly admitted to hospital with heart failure: survival trends in 12 220 index admissions in Leicestershire 1993–2001, **615**
- epilepsy**, Differential diagnosis of cardiogenic syncope and seizure disorders, **353**
- epoprostenol**, Intravascular ultrasound of the elastic pulmonary arteries: a new approach for the evaluation of primary pulmonary hypertension, **311**
- erectile dysfunction**, Erectile dysfunction and the cardiovascular patient: endothelial dysfunction is the common denominator, **251**
Role of exercise treadmill testing in the management of erectile dysfunction: a joint cardiovascular/erectile dysfunction clinic, **671**
- ergometric test**, Fontan conversion to total cavopulmonary connection and arrhythmia ablation: clinical and functional results, **193**
- essential hypertension**, Rarefaction of skin capillaries in normotensive offspring of individuals with essential hypertension, **175**
- ethnic differences**, Trends and ethnic differences in hospital admissions and mortality for congestive heart failure in the elderly in Singapore, 1991 to 1998, **865**
- ethnic groups**, The applicability of the Framingham coronary heart disease prediction function to black and minority ethnic groups in the UK, **785**
- ethnic minorities**, Improving the delivery of coronary care for ethnic minorities, **479**
- ethnicity**, Ethnic differences in cardiovascular disease, **681**
- exercise**, Comprehensive cardiac rehabilitation programme for implantable cardioverter defibrillator patients: a randomised controlled trial, **155**
Exercise systolic blood pressures are of questionable value in the assessment of the adult with a previous coarctation repair, **189**
- exercise blood pressure**, Exercise blood pressure in congenital heart disease and in patients after coarctation repair, **125**
- exercise capacity**, Safety and potential benefits of low dose diamorphine during exercise in patients with chronic heart failure, **1085**

- exercise physiology**, Acetylcholinesterase inhibition with pyridostigmine improves heart rate recovery after maximal exercise in patients with chronic heart failure, **854**
- exercise testing**, Improving the positive predictive value of exercise testing in women, **1416**
The effects of α and β blockade on ventilatory responses to exercise in chronic heart failure, **1169**
- exercise training**, Attenuation of hypercapnic carbon dioxide chemosensitivity after postinfarction exercise training: possible contribution to the improvement in exercise hyperventilation, **404**
- exercise treadmill testing**, Role of exercise treadmill testing in the management of erectile dysfunction: a joint cardiovascular/erectile dysfunction clinic, **671**
- Fabry disease**, Fabry disease, **819**
Fabry disease: a mimic for obstructive hypertrophic cardiomyopathy?, **929**
- familial dilated cardiomyopathy**, Progression of familial and non-familial dilated cardiomyopathy: long term follow up, **757**
- familial hypercholesterolaemia**, Long term statin treatment reduces lipoprotein(a) concentrations in heterozygous familial hypercholesterolaemia, **893**
- familial Mediterranean fever**, Incidence of pericardial effusion during attacks of familial Mediterranean fever, **1257**
- family history**, Family history of premature death from ischaemic heart disease is associated with an increased risk of delivering a low birth weight baby, **1249**
- Fas**, Relation between circulating soluble Fas ligand and subsequent ventricular remodelling following myocardial infarction, **339**
- fetal tachycardia**, Management of fetal tachyarrhythmia based on superior vena cava/aorta Doppler flow recordings, **1211**
- fetal/neonatal outcome**, Common arterial trunk in the fetus: characteristics, associations, and outcome in a multicentre series of 23 cases, **1437**
- fetus**, Review of diagnosis, treatment, and outcome of fetal atrial flutter compared with supraventricular tachycardia, **913**
Tricuspid regurgitation in the diagnosis of chromosomal anomalies in the fetus at 11–14 weeks of gestation, **1071**
- fibrillatory arrest**, Coronary surgery with non-cardioplegic methods in patients with advanced left ventricular dysfunction: immediate and long term results, **427**
- flash pulmonary oedema**, Flash pulmonary oedema: accurate evaluation of the renal arteries with multislice computed tomography, **1251**
- flecainide**, Flecainide induced ventricular fibrillation in a neonate, **1261**
- Fontan operation**, Acquired combined immunodeficiency associated with protein losing enteropathy complicating Fontan operation, **1130**
- Fontan procedure**, Modified technique of stent fenestration of the atrial septum, **1227**
- football**, Lower myocardial infarction mortality in French men the day France won the 1998 World Cup of football, **555**
- force**, Dissection of the aorta: a new approach, **6**
- frequency**, Pattern of ventilation during exercise in chronic heart failure, **610**
- gangrene**, Trash feet after coronary angiography, **e17**
- gated SPECT**, Role of myocardial perfusion imaging for risk stratification in suspected or known coronary artery disease, **1291**
- gene polymorphism**, DD ACE gene polymorphism is associated with increased coronary artery endothelial dysfunction: the PREFACE trial, **557**
- gene-gene interaction**, Epistatic interaction between variations in the angiotensin I converting enzyme and angiotensin II type 1 receptor genes in relation to extent of coronary atherosclerosis, **1195**
- general practice**, Survival differences between heart failure in general practices and in hospitals, **1298**
- genetic polymorphism**, Epistatic interaction between variations in the angiotensin I converting enzyme and angiotensin II type 1 receptor genes in relation to extent of coronary atherosclerosis, **1195**
- genetics**, Fabry disease, **819**
Mutations of the β myosin heavy chain gene in hypertrophic cardiomyopathy: critical functional sites determine prognosis, **1179**
- genome**, Microarray analysis: a novel research tool for cardiovascular scientists and physicians, **597**
- globalisation**, Heart failure: a global disease requiring a global response, **585**
- glucose-insulin-potassium infusion**, Effects of glucose-insulin-potassium infusion on chronic ischaemic left ventricular dysfunction, **61**
- glycaemic index**, Carbohydrates, dietary glycaemic load and glycaemic index, and risk of acute myocardial infarction, **722**
- glycaemic load**, Carbohydrates, dietary glycaemic load and glycaemic index, and risk of acute myocardial infarction, **722**
- glycoprotein IIb/IIIa inhibitors**, Antiplatelet treatment in stable coronary artery disease, **1273**
Who would I not give IIb/IIIa inhibitors to during percutaneous coronary intervention?, **477**
- granuloma**, A 28 year old woman with ventricular tachycardia and an abnormal chest radiograph, **e19**
- guanylyl cyclase**, B Type natriuretic peptide: a good omen in myocardial ischaemia?, **707**
- haemochromatosis**, The prevalence of haemochromatosis gene mutations in the West of Scotland and their relation to ischaemic heart disease, **1023**
- haemodynamics**, Clinical and haemodynamic profiles of young, middle aged, and elderly patients with mitral stenosis undergoing mitral balloon valvotomy, **1430**
Improved postoperative outcomes with stentless aortic valve: a community hospital experience, **551**
- haemolysis**, Prevalence and clinical significance of incidental paraprostatic valvar regurgitation: a prospective study using transoesophageal echocardiography, **1316**
- haemoptysis**, Prevalence of haemoptysis in adults with pulmonary atresia and ventricular septal defect, and the role of mammary artery collateral vessels, **937**
- haemorheology**, Platelet activation, haemorheology and thrombogenesis in acute atrial fibrillation: a comparison with permanent atrial fibrillation, **1093**
- haemorrhage**, Deaths associated with platelet glycoprotein IIb/IIIa inhibitor treatment, **535**
- haemorrhagic events**, Experience with enoxaparin in patients with mechanical heart valves who must withhold acenocumarol, **527**
- hand held echocardiography**, Screening for left ventricular dysfunction in the community: role of hand held echocardiography and brain natriuretic peptides, **iii24**
- hand held ultrasound device**, Clinical utility and cost effectiveness of a personal ultrasound imager for cardiac evaluation during consultation rounds in patients with suspected cardiac disease, **727**
- harmonic power Doppler**, Assessment of myocardial perfusion abnormalities by intravenous myocardial contrast echocardiography with harmonic power Doppler imaging: comparison with positron emission tomography, **145**
- heart**, Angiotensin II receptor blockade prevents microangiopathy and preserves diastolic function in the diabetic rat heart, **1236**
- heart development**, Development of the heart: (1) Formation of the cardiac chambers and arterial trunks, **806**
Development of the heart: (2) Septation of the atriums and ventricles, **949**
Development of the heart: (3) Formation of the ventricular outflow tracts, arterial valves, and intrapericardial arterial trunks, **1110**
- heart failure**, Brain natriuretic peptide facilitates severity classification of stable chronic heart failure with left ventricular dysfunction, **661**
Colour tissue velocity imaging can show resynchronisation of longitudinal left ventricular contraction pattern by biventricular pacing in patients with severe heart failure, **859**
Detection of scarred and viable myocardium using a new magnetic resonance imaging technique: blood oxygen level dependent (BOLD) MRI, **738**
Diastolic heart failure: fact or fiction?, **1281**
Different responses to dobutamine in the presence of carvedilol or metoprolol in patients with chronic heart failure, **621**
Estimating prognosis in heart failure: time for a better approach, **587**
Heart failure: a global disease requiring a global response, **585**
Heart failure and the aging population: an increasing burden in the 21st century?, **49**
High prevalence of left ventricular systolic and diastolic asynchrony in patients with congestive heart failure and normal QRS duration, **54**
Importance of β blockade in the treatment of advanced heart failure, **1442**
Increased distance between mitral valve coaptation point and mitral annular plane: significance and correlations in patients with heart failure, **1174**

- Increased urinary 15-F_{2t}-isoprostane concentrations in patients with non-ischaemic congestive heart failure: a marker of oxidative stress, **871**
- Interventricular and intra-left ventricular electromechanical delays in right ventricular paced patients with heart failure: implications for upgrading to biventricular stimulation, **1401**
- Lack of association between angiotensin converting enzyme (ACE) genotype, serum ACE activity, and haemodynamics in patients with primary pulmonary hypertension, **445**
- N-terminal pro B type natriuretic peptide, but not the new putative cardiac hormone relaxin, predicts prognosis in patients with chronic heart failure, **879**
- Obstacles to the initiation of β blockers for heart failure in a specialised clinic within a district general hospital, **442**
- Pathophysiology of coronary microembolisation, **981**
- Predicting mortality in patients with heart failure: a pragmatic approach, **605**
- Prevalence of impaired left ventricular systolic function and heart failure in a middle aged and elderly urban population segment of Copenhagen, **1422**
- Prognosis for patients newly admitted to hospital with heart failure: survival trends in 12 220 index admissions in Leicestershire 1993–2001, **615**
- QRS duration: a simple marker for predicting cardiac mortality in ICD patients with heart failure, **1157**
- Relation between heart rate, heart rhythm, and reverse left ventricular remodelling in response to carvedilol in patients with chronic heart failure: a single centre, observational study, **293**
- Restrictive transmitral filling patterns predict improvements in left ventricular function after biventricular pacing, **1087**
- Safety and potential benefits of low dose diamorphine during exercise in patients with chronic heart failure, **1085**
- Survival differences between heart failure in general practices and in hospitals, **1298**
- The influence of age, sex and other variables on the plasma level of N-terminal pro brain natriuretic peptide in a large sample of the general population, **745**
- The putative role of isoprostanes in human cardiovascular physiology and disease: following the fingerprints, **821**
- The therapeutic potential of new insights into myocardial excitation–contraction coupling, **371**
- Therapeutic inhibition of tumour necrosis factor α in patients with heart failure: cooling an inflamed heart, **14**
- Treatment of chronic heart failure with β adrenergic blockade beyond controlled clinical trials: the BRING-UP experience, **299**
- heart fatty acid binding protein**, Circulating concentrations of cardiac proteins indicate the severity of congestive heart failure, **1303**
- heart rate variability**, Heart rate variability in children with hypertrophic cardiomyopathy, **205**
- The relative contribution of the sinus and AV node to heart rate variability, **337**
- heart valve replacement**, Which heart valve prosthesis for patients aged between 60 and 70 years?, **481**
- heart valves**, The Edinburgh heart valve study, **697**
- Twenty year comparison of a Bjork-Shiley mechanical heart valve with porcine bioprostheses, **715**
- heterozygote**, Sisters with atypical Fabry's disease with complete atrioventricular block, **e2**
- HFE gene**, The prevalence of haemochromatosis gene mutations in the West of Scotland and their relation to ischaemic heart disease, **1023**
- high density lipoprotein**, Lipid lowering to delay the progression of coronary artery disease, **451**
- high risk patients**, Risk assessment in coronary artery surgery, **1**
- histology**, Histopathology of coronary in-stent restenosis following γ brachytherapy, **11**
- history**, Cor triatriatum sinister presenting in the adult as mitral stenosis, **e26**
- HIV**, HIV and cardiovascular medicine, **793**
- hospital admission**, N-terminal pro B type natriuretic peptide, but not the new putative cardiac hormone relaxin, predicts prognosis in patients with chronic heart failure, **879**
- The current cost of angina pectoris to the National Health Service in the UK, **848**
- hospital survival**, Factors associated with survival to hospital discharge among patients hospitalised alive after out of hospital cardiac arrest: change in outcome over 20 years in the community of Göteborg, Sweden, **25**
- human immunodeficiency virus**, HIV and cardiovascular medicine, **793**
- hypertension**, Ambulatory blood pressure monitoring in the management of hypertension, **571**
- Aortic valve and aortic arch pathology after coarctation repair, **1074**
- Cardiac and vascular pathophysiology in hypertension, **1104**
- Drug treatment of hypertension: implications of ALLHAT, **589**
- Long QTc predicts future cardiac death in stroke survivors, **377**
- hypertrophic cardiomyopathy**, Cardiac abnormalities in patients with Leber's hereditary optic neuropathy, **791**
- Contemporary considerations for risk stratification, sudden death and prevention in hypertrophic cardiomyopathy, **977**
- Fabry disease: a mimic for obstructive hypertrophic cardiomyopathy?, **929**
- Heart rate variability in children with hypertrophic cardiomyopathy, **205**
- Intramyocardial coronary flow characteristics in patients with hypertrophic cardiomyopathy: non-invasive assessment by transthoracic Doppler echocardiography, **657**
- Measurements of cardiac troponin T in patients with hypertrophic cardiomyopathy, **659**
- Pregnancy related complications in women with hypertrophic cardiomyopathy, **752**
- QT dispersion and left ventricular morphology in patients with hypertrophic cardiomyopathy, **882**
- hypertrophy**, Fabry disease, **819**
- Mutations of the β myosin heavy chain gene in hypertrophic cardiomyopathy: critical functional sites determine prognosis, **1179**
- hypoplastic left heart**, Stenting of the ductus arteriosus and banding of the pulmonary arteries: basis for various surgical strategies in newborns with multiple left heart obstructive lesions, **645**
- hypoplastic left heart syndrome**, Concordance for hypoplastic left heart syndrome in a monozygotic twin pregnancy, **e13**
- Images in cardiology**, A case of bepridil induced interstitial pneumonitis, **1415**
- A case of rheumatic fever, **1405**
- A diagnostic pitfall in a patient with an implantable cardioverter-defibrillator, **470**
- A giant thrombus aspirated from a coronary artery, **131**
- A novel form of spontaneous coronary vasospasm: bead-like vasospasm in an elderly man, **18**
- A rare cause of coronary obstruction and angina pectoris, **1120**
- Acquired coronary artery fistula leading to acute myocardial infarction after endomyocardial biopsy, **495**
- Acquired supraventricular type of left ventricular to right atrial communication following non-penetrating cardiac trauma caused by traffic accident, **341**
- Acute anterior myocardial infarction obscured by diaphragmatic eventration, **556**
- Acute left ventricular rupture after myocardial infarction, **138**
- Acute myocardial infarction caused by gunshot wound, **326**
- An incidental case of thrombus formation in a patient with a portacath inserted for regular blood transfusions, **244**
- An unusual complication of transvenous temporary pacing, **448**
- An unusual presentation of delayed cardiac perforation caused by atrial screw-in lead, **364**
- An unusual "right atrial mass" on echocardiography, **268**
- Anomalous origin of the pulmonary artery from the left coronary artery, **1436**
- Aortic abscess, **316**
- Aortic dissection extending to both carotid arteries, **5**
- Apical thrombus associated with left ventricular apical ballooning, **927**
- Area ablation of ventricular tachycardia in a patient with arrhythmogenic right ventricular cardiomyopathy, **703**
- Artefactual occlusion of the left subclavian artery during graft angiography, **1335**
- Brugada-like acute myocardial infarction, **1199**
- Cardiogenic shock in a young woman revealing juvenile haemochromatosis, **870**
- Catheter induced spasm: a trap for the unwary, **511**
- Complete occlusion of the abdominal aorta with a solitary kidney, **76**
- Computed tomography imaging of cardiac tamponade secondary to a posterior pericardial abscess, **364**
- Congenitally corrected transposition of the great arteries (L transposition), **976**

- Coronary arterial sling operation, **744**
- Cystic tumour of the atrioventricular node, **122**
- Complications of pacemaker leads, **1421**
- Coronary rupture as a rare cause of acute myocardial ischaemia two months after angioplasty and stenting, **1315**
- Different presentation of hypertrophic cardiomyopathy in monozygotic twins, **751**
- Effect of temperature on the Brugada ECG, **272**
- Electrocardiographic gated multislice computed tomography of the thoracic aorta, **838**
- Epidural haematoma presenting as polymorphic ventricular tachycardia, **316**
- Echocardiographic detection of systemic air embolism during positive pressure ventilation, **1321**
- Femoral vein delivery of contrast improves transoesophageal detection of intracardiac shunting through a patent foramen ovale, **784**
- First human use of a new PFM "Babystent", **83**
- Fragile left atrial thrombus successfully treated with anticoagulants, **60**
- Giant P waves, **1151**
- Giant right atrial thrombosis associated with Hickman catheter, **1252**
- Himalayan P waves in a patient with combined tricuspid and pulmonic stenosis, **1216**
- Huge mass bordering on left atrium raises coronary artery, **138**
- Iatrogenic left coronary artery dissection, **886**
- ICD lead implantation via persistent left superior vena cava, **1307**
- Imaging of a huge atrial myxoma, **99**
- Imitating ventricular tachycardia, **1382**
- Improved detection of protruding apical thrombus with contrast echocardiography, **1297**
- Improved myocardial viability following intramyocardial autologous bone marrow injection after acute myocardial infarction, **930**
- Incidence of embolism and paravalvar leak after St Jude Silzone valve implantation: experience from the Cardiff Embolic Risk Factor Study, **1055**
- Infective endocarditis with progressive periaortic abscess formation in a previously healthy girl, **730**
- In-stent restenosis and atherosclerosis in a human saphenous vein graft, **132**
- Intermittent opening of a mitral valve tilting disc prosthesis 23 years after implantation caused by circumferential pannus, **1049**
- Large B cell lymphoma with cardiac infiltration, **1282**
- Left atrial standstill in a patient with mitral stenosis and sinus rhythm: a risk of thrombus hidden by left and right atrial electrical dissociation, **1173**
- Left atrial thrombus caused by the primary antiphospholipid syndrome causing critical functional mitral stenosis, **262**
- Left circumflex coronary artery arising as a terminal extension of the right coronary artery, **286**
- Left main coronary artery fistula exiting into the right atrium, **656**
- Left ventricular microaneurysms as a cause of apparently idiopathic ventricular fibrillation, **644**
- Levoatriocardinal vein without cardiac malformation and normal pulmonary venous return, **1444**
- Life threatening total occlusion of the left coronary artery after acetylcholine infusion, **1137**
- Long term warfarin associated with bilateral blindness in a patient with atrial fibrillation and macular degeneration, **985**
- Low output, low gradient aortic stenosis, **474**
- Malposition of ventricular pacemaker lead as an incidental finding, **332**
- Management of complex, two chamber false aneurysm by thrombin injections, **564**
- Misplaced pacemaker wire as a cause of mitral regurgitation, **246**
- Multiple coronary artery fistulas visualised by three dimensional computed tomography, **1381**
- Multiple coronary rupture after blunt chest trauma, **594**
- Multiple cystic aneurysms in aortitis demonstrated by three dimensional volume rendering images of multislice computed tomography, **257**
- Multi-vessel dissections in Marfan syndrome demonstrated by multislice computed tomography, **1146**
- Myocardial bridge-like arterial graft compression by draining tube, **822**
- Myocardial infarction with myocardial rupture, **1077**
- Non-haemorrhagic cardiac tamponade caused by a horse kick, **330**
- Pacemaker lead fracture, **783**
- Pacing the left ventricle through the coronary sinus in a patient with a prosthetic tricuspid valve replacement, **1444**
- Partial anomalous pulmonary venous connection or scimitar syndrome, **761**
- Patency of coronary artery lumen surrounded by metallic stent evaluated by three dimensional volume rendering images using ECG gated multislice computed tomography, **388**
- Patency of multiple coronary artery bypass grafts demonstrated by multislice CT, **818**
- Percutaneous balloon fenestration of the intimal flap for management of limb threatening ischaemia in acute aortic dissection, **973**
- Percutaneous closure of a false aneurysm of the superficial femoral artery under fluoroscopic guidance, **326**
- Pituitary apoplexy following anticoagulation for acute coronary syndrome, **10**
- Prevention of embolisation during percutaneous vein graft intervention using a Filter Wire distal protection device, **376**
- Prolonged asymptomatic sinus pause indicated by implantable loop recording, **244**
- Prolonged cardiac arrest caused by glossopharyngeal neuralgia, **381**
- Prolonged sinus arrest after coronary artery spasm, **558**
- Pulmonary arteriovenous fistula, **41**
- Pulmonary artery sarcoma, **1388**
- Pulmonary valve endocarditis, **480**
- Rheumatic carditis in an 18 year old man, **1066**
- Right atrial thrombus following closure of an atrial septal defect, **726**
- Right atrial tumour mass of unusual origin, **1335**
- Right atrium and testis tumour: is there a link?, **1129**
- Ruptured thoracic aortic aneurysm resulting in cutaneous haematoma, **1382**
- Severe pulmonary hypertension in a patient with hypertrophic cardiomyopathy: response to alcohol septal ablation, **790**
- Should you remove an implantable loop recorder after the diagnosis is made?, **1013**
- Significant inducible perfusion abnormality in an asymptomatic patient with hypertrophic cardiomyopathy demonstrated by radionuclide myocardial perfusion imaging, **989**
- Spontaneous coronary artery dissection: stent it or wait for healing?, **164**
- Spontaneous coronary artery dissection in a pre-menopausal woman presenting with transient ST segment elevation, **1326**
- Spontaneous dissection of left atrium presenting as pulmonary oedema, **1178**
- Spontaneous pneumopericardium, **1250**
- Subclavian artery stenosis as a cause for recurrent angina after LIMA graft stenting, **1429**
- Submitral aneurysm: unusual echocardiographic features, **552**
- Sudden cardiac death caused by hypertrophic cardiomyopathy associated with midventricular obstruction and apical aneurysm, **178**
- Surgical resection of multiple right atrial masses in a child with β thalassaemia and receiving desferioxamine, **501**
- Suspected bilateral renal artery stenosis, **1070**
- Synergistic anticoagulation and transoesophageal echocardiographic surveillance for persistent left atrial appendage thrombus, **1022**
- β Thalassaemia major and severe left ventricular dysfunction, **1162**
- The dynamics of an ascending aorta dissection by 16 row multislice computed tomography, **970**
- The "pop phenomenon" detected by phased array intracardiac echocardiography, **706**
- Thinned myocardial fibrosis with thrombus in the dilated form of hypertrophic cardiomyopathy demonstrated by multislice computed tomography, **858**
- Thirty years after percutaneous closure of patent ductus arteriosus, **650**
- Three dimensional images with ECG gated multislice computed tomography revealed stenosis of the descending aorta in 2 month old baby, **444**
- Transcatheter closure of direct communication between right pulmonary artery and left atrium using Amplatzer device, **1210**
- Transcatheter closure of large fistula between left main coronary artery and right atrium using Amplatzer duct occluder, **772**
- Transcatheter embolisation of a large unilateral pulmonary arteriovenous malformation, **737**
- Transoesophageal echocardiography of a large tricuspid valve vegetation: a perfect image of reality, **696**
- Unusual morphologic changes in apical hypertrophic cardiomyopathy, **1290**

- Vasospastic angina, **198**
- Ventricular tachycardia following head injury, **829**
- imaging**, Evaluation of intracoronary stenting by intravascular optical coherence tomography, **317**
- immunodeficiency**, Acquired combined immunodeficiency associated with protein losing enteropathy complicating Fontan operation, **1130**
- impaired left ventricular function**, Prospective evaluation of the role of routine cardiac troponin T measurement to identify left ventricular ejection fraction 40% after first myocardial infarction, **559**
- implantable cardioverter-defibrillators**, An audit of the implications of implementing NICE guidance on the use of implantable cardioverter-defibrillators, **787**
- Comprehensive cardiac rehabilitation programme for implantable cardioverter defibrillator patients: a randomised controlled trial, **155**
- QRS duration: a simple marker for predicting cardiac mortality in ICD patients with heart failure, **1157**
- implantable cardioverter-defibrillator leads**, Symptomatic occlusion of the access vein after pacemaker or ICD lead extraction, **1348**
- infant**, Catheter ablation of sinoatrial re-entry tachycardia in a 2 month old infant, **e1**
- infarction**, B Type natriuretic peptide: a good omen in myocardial ischaemia?, **707**
- infective endocarditis**, Judicious use of transthoracic echocardiography in the diagnosis of infective endocarditis, **1283**
- Molecular diagnosis of culture negative infective endocarditis: clinical validation in a group of surgically treated patients, **263**
- inflammation**, C reactive protein for risk stratification in acute coronary syndromes? Verdict: unproven, **1288**
- CD14 promoter polymorphism is associated with acute myocardial infarction resulting from insignificant coronary artery stenosis, **931**
- Impact of preprocedural white blood cell count on long term mortality after percutaneous coronary intervention: insights from the EPIC, EPILOG, and EPISTENT trials, **1200**
- Inflammatory and thrombotic mechanisms in coronary atherosclerosis, **993**
- Pathophysiology of coronary microembolisation, **981**
- The elusive link between stenosis severity and prognosis in stable ischaemic heart disease, **961**
- Why cardiologists should be interested in air pollution, **1383**
- inotropic agents**, Different responses to dobutamine in the presence of carvedilol or metoprolol in patients with chronic heart failure, **621**
- insertable loop recorder**, Novel use of an "insertable" loop recorder, **e18**
- interactive**, Cased based learning on the web, **979**
- interleukin 1 β** , High preprocedural non-HDL cholesterol is associated with enhanced oxidative stress and monocyte activation after coronary angioplasty: possible implications in restenosis, **773**
- international mortality**, International comparison of mortality rates in patients with non-ST elevation acute coronary events, **1083**
- interstitial fibrosis**, Interstitial fibrosis in the dilated non-ischaemic myocardium, **1255**
- interventional cardiology**, Coarctation of the aorta in adults: do we need surgeons?, **3**
- interventional catheterisation**, Use of 3 French catheters for diagnostic and interventional procedures in newborns and small infants, **1350**
- intima-media thickness**, Long term statin treatment reduces lipoprotein(a) concentrations in heterozygous familial hypercholesterolaemia, **893**
- intra-aortic balloon pump**, Emergency coronary stenting of unprotected critical left main coronary artery stenosis in acute myocardial infarction and cardiogenic shock, **e24**
- intracoronary Doppler**, Invasive assessment of myocardial bridges, **699**
- intramyocardial coronary flow**, Intramyocardial coronary flow characteristics in patients with hypertrophic cardiomyopathy: non-invasive assessment by transthoracic Doppler echocardiography, **657**
- intravascular ultrasonography**, Continued improvement of clinical outcome and cost effectiveness following intravascular ultrasound guided PCI: insights from a prospective, randomised study, **1043**
- intravascular ultrasound**, Intravascular ultrasound of the elastic pulmonary arteries: a new approach for the evaluation of primary pulmonary hypertension, **311**
- Invasive assessment of myocardial bridges, **699**
- Mechanism of lumen enlargement with direct stenting versus predilatation stenting: influence of remodelling and plaque characteristics assessed by volumetric intracoronary ultrasound, **84**
- intravenous drug use**, Injection drug use and right sided endocarditis, **577**
- iron toxicity**, Left ventricular remodelling, and systolic and diastolic function in young adults with β thalassaemia major: a Doppler echocardiographic assessment and correlation with haematological data, **762**
- ischaemia**, B Type natriuretic peptide: a good omen in myocardial ischaemia?, **707**
- Left ventricular remodelling after myocardial infarction: importance of residual myocardial viability and ischaemia, **1121**
- Stress echocardiography, **113**
- ischaemic heart disease**, Detection of scarred and viable myocardium using a new magnetic resonance imaging technique: blood oxygen level dependent (BOLD) MRI, **738**
- Effects of glucose-insulin-potassium infusion on chronic ischaemic left ventricular dysfunction, **61**
- Enhanced external counterpulsation for refractory angina pectoris, **830**
- Ethnic differences in cardiovascular disease, **681**
- Family history of premature death from ischaemic heart disease is associated with an increased risk of delivering a low birth weight baby, **1249**
- Long term outcome of patients with postinfarction left ventricular pseudoaneurysm, **1144**
- Management of ischaemic mitral regurgitation, **459**
- Smoking: can we really make a difference?, **ii25**
- The prevalence of haemochromatosis gene mutations in the West of Scotland and their relation to ischaemic heart disease, **1023**
- isoprostane**, The putative role of isoprostanes in human cardiovascular physiology and disease: following the fingerprints, **821**
- isovolumic relaxation**, Clinical assessment of left ventricular diastolic function, **231**
- Japanese women**, Variation during the week in the incidence of acute myocardial infarction: increased risk for Japanese women on Saturdays, **398**
- lanotepase**, Thrombolysis: newer thrombolytic agents and their role in clinical medicine, **1358**
- L-arginine**, Effects of L-arginine on flow mediated dilatation induced by atrial pacing in diseased epicardial coronary arteries, **531**
- lead extraction**, Symptomatic occlusion of the access vein after pacemaker or ICD lead extraction, **1348**
- Leber's hereditary optic atrophy**, Cardiac abnormalities in patients with Leber's hereditary optic neuropathy, **791**
- left atrial failure**, Effects of digoxin on left atrial function in heart failure, **1308**
- left atrial volume**, Left atrial and left ventricular function in healthy children and young adults assessed by three dimensional echocardiography, **544**
- left bundle branch block**, Body surface mapping improves early diagnosis of acute myocardial infarction in patients with chest pain and left bundle branch block, **998**
- left main coronary artery**, Long term follow up after elective percutaneous coronary intervention for unprotected non-bifurcational left main stenosis: is it time to change the guidelines?, **1336**
- left main stem stenosis**, Takayasu's arteritis: management of left main stem stenosis, **e9**
- left sinus of Valsalva**, Unruptured congenital aneurysm of the left sinus of Valsalva in an adult with complex left heart malformations, **e3**
- left ventricle**, Long term outcome of patients with postinfarction left ventricular pseudoaneurysm, **1144**
- left ventricular assessment**, Left ventricular assessment using real time three dimensional echocardiography, **iii29**
- left ventricular diastolic function**, A clinical approach to the assessment of left ventricular diastolic function by Doppler echocardiography: update 2003, **iii18**
- Clinical applications of tissue Doppler imaging: a promise fulfilled, **1377**
- Clinical assessment of left ventricular diastolic function, **231**
- left ventricular dysfunction**, Brain natriuretic peptide facilitates severity classification of stable chronic heart failure with left ventricular dysfunction, **661**
- Coronary surgery with non-cardioplegic methods in patients with advanced left ventricular dysfunction: immediate and long term results, **427**

- Early detection of left ventricular dysfunction in patients with β thalassaemia major by dobutamine stress echocardiography, **669**
- Left ventricular remodelling, and systolic and diastolic function in young adults with β thalassaemia major: a Doppler echocardiographic assessment and correlation with haematological data, **762**
- Protecting the damaged heart during coronary surgery, **367**
- Screening for left ventricular dysfunction in the community: role of hand held echocardiography and brain natriuretic peptides, **iii24**
- left ventricular function**, Diastolic heart failure: fact or fiction?, **1281**
- Haemodynamic and catecholamine response to simulated ventricular tachycardia in man: effect of baseline left ventricular function, **306**
- left ventricular hypertrophy**, Athlete's heart, **1455**
- left ventricular morphology**, QT dispersion and left ventricular morphology in patients with hypertrophic cardiomyopathy, **882**
- left ventricular opacification**, Contrast echocardiography, **342**
- Contrast echocardiography for left ventricular opacification, **1394**
- left ventricular remodelling**, Left ventricular remodelling after myocardial infarction: importance of residual myocardial viability and ischaemia, **1121**
- left ventricular subendocardial function**, Conduit arterial stiffness is associated with impaired left ventricular subendocardial function, **449**
- left ventricular systolic contraction**, Clinical applications of tissue Doppler imaging: a promise fulfilled, **1377**
- left ventricular systolic function**, Techniques for comprehensive two dimensional echocardiographic assessment of left ventricular systolic function, **iii2**
- left ventricular volume**, Left atrial and left ventricular function in healthy children and young adults assessed by three dimensional echocardiography, **544**
- Legionella pneumophila**, Innocent victim of a localised outbreak: legionella endocarditis, **e16**
- leisure time physical activity**, What level of physical activity protects against premature cardiovascular death? The Caerphilly study, **502**
- lesion size**, Real time quantification of low temperature radiofrequency ablation lesion size using phased array intracardiac echocardiography in the canine model: comparison of two dimensional images with pathological lesion characteristics, **923**
- lipid lowering**, Lipid lowering to delay the progression of coronary artery disease, **451**
- lipid lowering drugs**, Evolution of statin prescribing 1994–2001: a case of agism but not of sexism?, **417**
- lipid peroxidation**, The putative role of isoprostanes in human cardiovascular physiology and disease: following the fingerprints, **821**
- lipoprotein(a)**, Long term statin treatment reduces lipoprotein(a) concentrations in heterozygous familial hypercholesterolaemia, **893**
- lisinopril**, Mode of death in heart failure: findings from the ATLAS trial, **42**
- long term follow up**, Coronary risk factors predicting early and late coronary deaths, **19**
- long term outcome**, Prospective evaluation of the effect of an angiotensin I converting enzyme gene polymorphism on the long term risk of major adverse cardiac events after percutaneous coronary intervention, **321**
- long term survival**, A prospective study of long term prognosis in young myocardial infarction survivors: the prognostic value of angiography and exercise testing, **843**
- loop recorder**, Artefact mimicking tachycardia during magnetic resonance imaging in a patient with an implantable loop recorder, **e10**
- low birth weight**, Family history of premature death from ischaemic heart disease is associated with an increased risk of delivering a low birth weight baby, **1249**
- low density lipoprotein**, Lipid lowering to delay the progression of coronary artery disease, **451**
- low molecular weight heparin**, Cost effectiveness of extended treatment with low molecular weight heparin (dalteparin) in unstable coronary artery disease: results from the FRISC II trial, **287**
- The primacy of clinical effectiveness for cost effectiveness analysis, **249**
- low pressure giant pulmonary artery aneurysm**, Low pressure giant pulmonary artery aneurysms in the adult: natural history and management strategies, **1067**
- low temperature radiofrequency ablation**, Real time quantification of low temperature radiofrequency ablation lesion size using phased array intracardiac echocardiography in the canine model: comparison of two dimensional images with pathological lesion characteristics, **923**
- Lp(a)**, Long term statin treatment reduces lipoprotein(a) concentrations in heterozygous familial hypercholesterolaemia, **893**
- magnesium**, Serum magnesium aberrations in furosemide (frusemide) treated patients with congestive heart failure: pathophysiological correlates and prognostic evaluation, **411**
- magnetic resonance imaging**, Anomalous origin of the circumflex artery and patent foramen ovale: a rare cause of myocardial ischaemia after percutaneous closure of the defect, **e23**
- Artefact mimicking tachycardia during magnetic resonance imaging in a patient with an implantable loop recorder, **e10**
- Non-invasive coronary imaging: computed tomography or magnetic resonance imaging?, **591**
- major vascular surgery**, A meta-analysis comparing the prognostic accuracy of six diagnostic tests for predicting perioperative cardiac risk in patients undergoing major vascular surgery, **1327**
- “male menopause”**, Cardiovascular effects of testosterone: implications of the “male menopause”?, **121**
- mapping**, Body surface mapping improves early diagnosis of acute myocardial infarction in patients with chest pain and left bundle branch block, **998**
- maternal diabetes**, Cardiovascular malformations in infants of diabetic mothers, **1217**
- mean platelet component**, The central role of platelet activation in determining the severity of acute coronary syndromes, **1253**
- mechanical heart valve**, Experience with enoxaparin in patients with mechanical heart valves who must withhold acenocumarol, **527**
- mechanical prosthesis**, Which heart valve prosthesis for patients aged between 60 and 70 years?, **481**
- mechanical valves**, The Edinburgh heart valve study, **697**
- meta-analysis**, Adverse events in coronary artery bypass graft (CABG) trials: a systematic review and analysis, **767**
- metabolism**, Fabry disease, **819**
- microarray**, Microarray analysis: a novel research tool for cardiovascular scientists and physicians, **597**
- microcirculation**, Rarefaction of skin capillaries in normotensive offspring of individuals with essential hypertension, **175**
- Relation of coronary flow pattern to myocardial blush grade in patients with first acute myocardial infarction, **1147**
- Temporal evolution and functional outcome of no reflow: sustained and spontaneously reversible patterns following successful coronary recanalisation, **731**
- miniaturisation**, Use of 3 French catheters for diagnostic and interventional procedures in newborns and small infants, **1350**
- mitochondria**, Cardiac abnormalities in patients with Leber's hereditary optic neuropathy, **791**
- mitral annular calcification**, Mitral annular calcification: a marker of severe coronary artery disease in patients under 65 years old, **161**
- mitral atresia**, Concordance for hypoplastic left heart syndrome in a monozygotic twin pregnancy, **e13**
- mitral balloon valvotomy**, Clinical and haemodynamic profiles of young, middle aged, and elderly patients with mitral stenosis undergoing mitral balloon valvotomy, **1430**
- mitral regurgitation**, Management of ischaemic mitral regurgitation, **459**
- Preoperative left ventricular end systolic dimension as a predictor of postoperative ventricular dysfunction in children with mitral regurgitation, **1243**
- Timing of surgery in mitral regurgitation, **100**
- mitral stenosis**, Clinical and haemodynamic profiles of young, middle aged, and elderly patients with mitral stenosis undergoing mitral balloon valvotomy, **1430**
- mitral surgery**, Maurice Campbell: first editor of *Heart*, **1379**
- mitral valve**, Increased distance between mitral valve coaptation point and mitral annular plane: significance and correlations in patients with heart failure, **1174**
- Timing of surgery in mitral regurgitation, **100**
- mobilisation**, Position and Mobilisation Post-Angiography Study (PAMPAS): a comparison of 4.5 hours and 2.5 hours bed rest, **447**
- molecular biology**, Molecular diagnosis of culture negative infective endocarditis: clinical validation in a group of surgically treated patients, **263**
- Mutations of the β myosin heavy chain gene in hypertrophic cardiomyopathy: critical functional sites determine prognosis, **1179**

- monochorionic twin pregnancy**, Concordance for hypoplastic left heart syndrome in a monochorionic twin pregnancy, **e13**
- morbidity**, The current cost of angina pectoris to the National Health Service in the UK, **848**
- morphine**, Enhanced endothelin-1 degradation by intravenous morphine in patients with congestive heart failure: role of neutral endopeptidase 24.11, **211**
- mortality**, Adverse events in coronary artery bypass graft (CABG) trials: a systematic review and analysis, **767**
- Associations between ST depression, four year mortality, and in-hospital revascularisation in unselected patients with non-ST elevation acute coronary syndromes, **490**
- Deaths associated with platelet glycoprotein IIb/IIIa inhibitor treatment, **535**
- Importance of β blockade in the treatment of advanced heart failure, **1442**
- Lower myocardial infarction mortality in French men the day France won the 1998 World Cup of football, **555**
- Predicting mortality in patients with heart failure: a pragmatic approach, **605**
- Prognosis for patients newly admitted to hospital with heart failure: survival trends in 12 220 index admissions in Leicestershire 1993–2001, **615**
- What level of physical activity protects against premature cardiovascular death? The Caerphilly study, **502**
- multiple pulmonary stenosis**, Williams syndrome associated with complete atrioventricular septal defect, **e15**
- multislice computed tomography**, Flash pulmonary oedema: accurate evaluation of the renal arteries with multislice computed tomography, **1251**
- Non-invasive intravenous coronary angiography using electron beam tomography and multislice computed tomography, **633**
- multivessel coronary stenting**, Coronary stenting versus coronary bypass surgery in patients with multiple vessel disease and significant proximal LAD stenosis: results from the ERACI II study, **184**
- multivessel revascularisation**, Coronary stenting versus coronary bypass surgery in patients with multiple vessel disease and significant proximal LAD stenosis: results from the ERACI II study, **184**
- myocardial blood flow**, Coronary flow velocity analysis during short term follow up after coronary reperfusion: use of transthoracic Doppler echocardiography to predict regional wall motion recovery in patients with acute myocardial infarction, **1163**
- Impaired myocardial blood flow and coronary flow reserve of the anatomical right systemic ventricle in patients with congenitally corrected transposition of the great arteries, **1231**
- myocardial bridge**, Invasive assessment of myocardial bridges, **699**
- myocardial contrast echocardiography**, Assessment of myocardial perfusion abnormalities by intravenous myocardial contrast echocardiography with harmonic power Doppler imaging: comparison with positron emission tomography, **145**
- Contrast echocardiography, **342**
- Role of myocardial contrast echocardiography in the clinical evaluation of acute myocardial infarction, **1398**
- myocardial infarction**, A prospective study of long term prognosis in young myocardial infarction survivors: the prognostic value of angiography and exercise testing, **843**
- Abortion of acute ST segment elevation myocardial infarction after reperfusion: incidence, patients' characteristics, and prognosis, **496**
- Apical ballooning of the left ventricle: first series in white patients, **1027**
- Are troponins confusing?, **1285**
- Assessment of myocardial viability using coronary zero flow pressure after successful angioplasty in patients with acute anterior myocardial infarction, **71**
- Attenuation of hypercapnic carbon dioxide chemosensitivity after postinfarction exercise training: possible contribution to the improvement in exercise hyperventilation, **404**
- Body surface mapping improves early diagnosis of acute myocardial infarction in patients with chest pain and left bundle branch block, **998**
- CD14 promoter polymorphism is associated with acute myocardial infarction resulting from insignificant coronary artery stenosis, **911**
- Coronary calcifications in young patients with first, unheralded myocardial infarction: a risk factor matched analysis by electron beam tomography, **625**
- Improving the delivery of coronary care for ethnic minorities, **479**
- Inflammatory and thrombotic mechanisms in coronary atherosclerosis, **993**
- Influence of contractile reserve and inducible ischaemia on left ventricular remodelling after acute myocardial infarction, **1138**
- Left ventricular remodelling after myocardial infarction: importance of residual myocardial viability and ischaemia, **1121**
- Lower myocardial infarction mortality in French men the day France won the 1998 World Cup of football, **555**
- Myocardial viability, coronary flow reserve, and in-hospital predictors of late recovery of contractility following successful primary stenting for acute myocardial infarction, **179**
- Persistence of systolic coronary flow reversal predicts irreversible dysfunction after reperfused anterior myocardial infarction, **382**
- Quality of life assessment using the short form 12 questionnaire is as reliable and sensitive as the short form 36 in distinguishing symptom severity in myocardial infarction survivors, **1445**
- Sources of diagnostic inaccuracy of conventional versus new diagnostic criteria for myocardial infarction in an unselected UK population with suspected cardiac chest pain, and investigation of independent prognostic variables, **1406**
- Stress and myocardial infarction, **475**
- Temporal evolution and functional outcome of no reflow: sustained and spontaneously reversible patterns following successful coronary recanalisation, **731**
- The central role of platelet activation in determining the severity of acute coronary syndromes, **1253**
- The elusive link between stenosis severity and prognosis in stable ischaemic heart disease, **961**
- The psychological perspective: a professional view, **ii16**
- Myocardial Infarction National Audit Project**, Where are we today? Early results from MINAP, the national audit of myocardial infarction project, **ii13**
- myocardial ischaemia**, Anomalous origin of the circumflex artery and patent foramen ovale: a rare cause of myocardial ischaemia after percutaneous closure of the defect, **e23**
- Regional myocardial perfusion defects during exercise, as assessed by three dimensional integration of morphology and function, in relation to abnormal endothelium dependent vasoreactivity of the coronary microcirculation, **517**
- myocardial perfusion imaging**, Role of myocardial perfusion imaging for risk stratification in suspected or known coronary artery disease, **1291**
- myocardial perfusion scintigraphy**, A meta-analysis comparing the prognostic accuracy of six diagnostic tests for predicting perioperative cardiac risk in patients undergoing major vascular surgery, **1327**
- myocardial protection**, Coronary surgery with non-cardioplegic methods in patients with advanced left ventricular dysfunction: immediate and long term results, **427**
- myocardial scintigraphy**, Fontan conversion to total cavopulmonary connection and arrhythmia ablation: clinical and functional results, **193**
- myocardial stunning**, Myocardial viability, coronary flow reserve, and in-hospital predictors of late recovery of contractility following successful primary stenting for acute myocardial infarction, **179**
- myocardial viability**, Assessment of myocardial viability: using coronary pressure and flow after acute myocardial infarction, **9**
- Assessment of myocardial viability using coronary zero flow pressure after successful angioplasty in patients with acute anterior myocardial infarction, **71**
- Left ventricular remodelling after myocardial infarction: importance of residual myocardial viability and ischaemia, **1121**
- myocardium**, Detection of scarred and viable myocardium using a new magnetic resonance imaging technique: blood oxygen level dependent (BOLD) MRI, **738**
- Transcardiac and transpulmonary gradients in the putative new cardiovascular hormone relaxin, **789**
- myoclonic jerks**, Differential diagnosis of cardiogenic syncope and seizure disorders, **353**
- myosin**, Mutations of the β myosin heavy chain gene in hypertrophic cardiomyopathy: critical functional sites determine prognosis, **1179**
- myosin light chain I**, Circulating concentrations of cardiac proteins indicate the severity of congestive heart failure, **1303**
- National Institute for Clinical Excellence**, An audit of the implications of implementing NICE guidance on the use of implantable cardioverter-defibrillators, **787**

- National Service Framework**, Current targets: where are we going?, ii6
Current targets: where are we today?, ii10
Where are we today? Early results from MINAP, the national audit of myocardial infarction project, ii13
- natriuretic peptides**, Importance of β blockade in the treatment of advanced heart failure, 1442
N-terminal pro B type natriuretic peptide, but not the new putative cardiac hormone relaxin, predicts prognosis in patients with chronic heart failure, 879
NT-proBNP: a new diagnostic screening tool to differentiate between patients with normal and reduced left ventricular systolic function, 150
- neonatal catheterisation**, Use of 3 French catheters for diagnostic and interventional procedures in newborns and small infants, 1350
- nesiritide**, B Type natriuretic peptide: a good omen in myocardial ischaemia?, 707
- neurodevelopmental outcome**, Neurodevelopment at 1 year of age in infants with congenital heart disease, 436
- neurogenic stunned myocardium**, Apical ballooning of the left ventricle: a distinct entity?, 974
- neutrophil adhesion**, Simvastatin attenuates leucocyte-endothelial interactions after coronary revascularisation with cardiopulmonary bypass, 538
- newborn**, Flecainide induced ventricular fibrillation in a neonate, 1261
- NICE**, An audit of the implications of implementing NICE guidance on the use of implantable cardioverter-defibrillators, 787
- nisoldipine**, Long term effects of nisoldipine on the progression of coronary atherosclerosis and the occurrence of clinical events: the NICOLE study, 887
- nitric oxide**, Effects of L-arginine on flow mediated dilatation induced by atrial pacing in diseased epicardial coronary arteries, 531
- no reflow**, Assessing perfusion and function in acute myocardial infarction: how and when?, 701
Pathology of coronary microembolisation and no reflow, 983
Temporal evolution and functional outcome of no reflow: sustained and spontaneously reversible patterns following successful coronary recanalisation, 731
- non-compaction cardiomyopathy**, Isolated left ventricular non-compaction: cardiomyopathy with homogeneous transmural and heterogeneous segmental perfusion, e21
- non-ST elevation acute coronary events**, International comparison of mortality rates in patients with non-ST elevation acute coronary events, 1083
- non-ST elevation acute coronary syndromes**, Cumulative risk assessment in unstable angina: clinical, electrocardiographic, autonomic, and biochemical markers, 36
White blood cell count and long term mortality after non-ST elevation acute coronary syndrome treated with very early revascularisation, 389
- Norwood repair**, Stenting of the ductus arteriosus and banding of the pulmonary arteries: basis for various surgical strategies in newborns with multiple left heart obstructive lesions, 645
- N-terminal proBNP**, NT-proBNP: a new diagnostic screening tool to differentiate between patients with normal and reduced left ventricular systolic function, 150
- nurses**, The nurse as expert practitioner in global cardiovascular risk management, ii33
- obesity**, Cardiac function and obesity, 1127
Effects of isolated obesity on systolic and diastolic left ventricular function, 1152
The obesity epidemic: can we turn the tide?, ii22
- observational studies**, Treatment of chronic heart failure with β adrenergic blockade beyond controlled clinical trials: the BRING-UP experience, 299
- occupational exposure**, Radiation exposure to cardiologists: how it could be reduced, 1123
- occupational stress**, Variation during the week in the incidence of acute myocardial infarction: increased risk for Japanese women on Saturdays, 398
- off-pump coronary artery bypass surgery**, Off-pump coronary artery bypass graft surgery: the incidence of postoperative atrial fibrillation, 1134
- out of hospital events**, Factors associated with survival to hospital discharge among patients hospitalised alive after out of hospital cardiac arrest: change in outcome over 20 years in the community of Göteborg, Sweden, 25
- outcome**, Clinical and haemodynamic profiles of young, middle aged, and elderly patients with mitral stenosis undergoing mitral balloon valvotomy, 1430
Long term outcome of patients with postinfarction left ventricular pseudoaneurysm, 1144
Presentation, management, and outcome of out of hospital cardiopulmonary arrest: comparison by underlying aetiology, 839
- oxidant stress**, High preprocedural non-HDL cholesterol is associated with enhanced oxidative stress and monocyte activation after coronary angioplasty: possible implications in restenosis, 773
- oxidative stress**, Increased urinary 15-F_{2t}-isoprostane concentrations in patients with non-ischaemic congestive heart failure: a marker of oxidative stress, 871
- oxygen consumption**, Safety and potential benefits of low dose diamorphine during exercise in patients with chronic heart failure, 1085
- pacemaker**, Ablate and pace revisited: long term survival and predictors of permanent atrial fibrillation, 1035
- pacemaker leads**, Symptomatic occlusion of the access vein after pacemaker or ICD lead extraction, 1348
- pacemakers**, How many cath labs do we need?, 827
- pacing**, Pacing for atrial fibrillation, 106
Prolonged temporary cardiac pacing using subcutaneous tunnelled active-fixation permanent pacing leads, 209
- paclitaxel**, In-stent stenosis: pathology and implications for the development of drug eluting stents, 218
- paradoxical embolism**, Learning on the web: Case 2: Patent foramen ovale (PFO) and paradoxical embolism, 1362
- paraprosthetic regurgitation**, Prevalence and clinical significance of incidental paraprosthetic valvar regurgitation: a prospective study using transoesophageal echocardiography, 1316
- parasympathetic nervous system**, Acetylcholinesterase inhibition with pyridostigmine improves heart rate recovery after maximal exercise in patients with chronic heart failure, 854
- paravalvar leak**, Incidence of embolism and paravalvar leak after St Jude Silzone valve implantation: experience from the Cardiff Embolic Risk Factor Study, 1055
- parental history**, Are there sex differences in risk factors for coronary heart disease? Maternal versus paternal transmission, 817
Is maternal transmission of coronary heart disease risk stronger than paternal transmission?, 834
- patent ductus arteriosus**, Challenges in the care of adult patients with congenital heart defects, 465
- patent foramen ovale**, Anomalous origin of the circumflex artery and patent foramen ovale: a rare cause of myocardial ischaemia after percutaneous closure of the defect, e23
Learning on the web: Case 2: Patent foramen ovale (PFO) and paradoxical embolism, 1362
- pathology**, Primary cardiac valve tumours, 1259
Real time quantification of low temperature radiofrequency ablation lesion size using phased array intracardiac echocardiography in the canine model: comparison of two dimensional images with pathological lesion characteristics, 923
- pathophysiology**, Cardiac and vascular pathophysiology in hypertension, 1104
- patient compliance**, Medicine partnerships, ii19
- patient factors**, Evolution of statin prescribing 1994–2001: a case of agism but not of sexism?, 417
- patient management**, Bangladeshi patients present with non-classic features of acute myocardial infarction and are treated less aggressively in east London, UK, 276
- peptides**, Transcardiac and transpulmonary gradients in the putative new cardiovascular hormone relaxin, 789
- percutaneous coronary intervention**, Continued improvement of clinical outcome and cost effectiveness following intravascular ultrasound guided PCI: insights from a prospective, randomised study, 1043
Drug eluting stents: maximising benefit and minimising cost, 127
Impact of preprocedural white blood cell count on long term mortality after percutaneous coronary intervention: insights from the EPIC, EPILOG, and EPISTENT trials, 1200
Indications for percutaneous and surgical revascularisation: how far does the evidence base guide us?, 565
Long term follow up after elective percutaneous coronary intervention for unprotected non-bifurcational left main stenosis: is it time to change the guidelines?, 1336

- Outcome following percutaneous coronary intervention: not, so far, in our genes, **247**
- Predictors of mortality in patients with acute coronary syndrome undergoing percutaneous coronary intervention, **1245**
- Revascularisation for acute coronary syndromes: PCI or CABG?, **967**
- Same day discharge following elective percutaneous coronary intervention in patients with stable angina, **665**
- Who would I not give IIb/IIIa inhibitors to during percutaneous coronary intervention?, **477**
- perfusion**, Isolated left ventricular non-compaction: cardiomyopathy with homogeneous transmural and heterogeneous segmental perfusion, **e21**
- pericardial effusion**, Pericardial effusion after open heart surgery for congenital heart disease, **780**
- Unexplained recurrent pericardial effusion: a lethal warning?, **e11**
- pericarditis**, Incidence of pericardial effusion during attacks of familial Mediterranean fever, **1257**
- Practical aspects of the management of pericardial disease, **1096**
- perimembranous ventricular septal defects**, Transcatheter closure of perimembranous ventricular septal defects with the Amplatzer asymmetric ventricular septal defect occluder: preliminary experience in children, **918**
- peripheral vascular resistance**, Cardiac and vascular pathophysiology in hypertension, **1104**
- permanent pacemaker**, Risk factors for pacemaker implantation following aortic valve replacement: a single centre experience, **901**
- personal ultrasound imager**, Clinical utility and cost effectiveness of a personal ultrasound imager for cardiac evaluation during consultation rounds in patients with suspected cardiac disease, **727**
- pharmacists**, Drugs and professional interactions: the modern day pharmacist, **ii31**
- pharmacological treatment**, Atrial fibrillation: classification, pathophysiology, mechanisms and drug treatment, **939**
- phased array intracardiac echocardiography**, Real time quantification of low temperature radiofrequency ablation lesion size using phased array intracardiac echocardiography in the canine model: comparison of two dimensional images with pathological lesion characteristics, **923**
- plaque stabilisation**, Plaque stabilisation in acute coronary syndromes: clinical considerations, **1268**
- plaque vulnerability**, Basic aspects of plaque vulnerability, **1262**
- platelet activation**, Platelet activation, haemorrhage and thrombogenesis in acute atrial fibrillation: a comparison with permanent atrial fibrillation, **1093**
- The central role of platelet activation in determining the severity of acute coronary syndromes, **1253**
- platelet aggregation inhibitors**, Deaths associated with platelet glycoprotein IIb/IIIa inhibitor treatment, **535**
- platelet reactivity**, Coronary angioplasty enhances platelet reactivity through von Willebrand factor release, **329**
- platelets**, Increased platelet sodium-hydrogen exchanger activity in patients with variant angina, **935**
- point of care system**, Portable spectral Doppler echocardiographic device: overcoming limitations, **1014**
- polymerase chain reaction**, Molecular diagnosis of culture negative infective endocarditis: clinical validation in a group of surgically treated patients, **263**
- polymorphism**, Prospective evaluation of the effect of an angiotensin I converting enzyme gene polymorphism on the long term risk of major adverse cardiac events after percutaneous coronary intervention, **321**
- population study**, The influence of age, sex and other variables on the plasma level of N-terminal pro brain natriuretic peptide in a large sample of the general population, **745**
- portable ultrasound**, The "ultrasonic stethoscope": is it of clinical value?, **704**
- positron emission computed tomography**, Assessment of myocardial perfusion abnormalities by intravenous myocardial contrast echocardiography with harmonic power Doppler imaging: comparison with positron emission tomography, **145**
- positron emission tomography**, Impaired myocardial blood flow and coronary flow reserve of the anatomical right systemic ventricle in patients with congenitally corrected transposition of the great arteries, **1231**
- postoperative arrhythmias**, Risk factors for pacemaker implantation following aortic valve replacement: a single centre experience, **901**
- practice guidelines**, Do guidelines influence practice?, **349**
- prediction**, Coronary risk factors predicting early and late coronary deaths, **19**
- predictive factors**, Mode of death in heart failure: findings from the ATLAS trial, **42**
- predictive models**, Predicting mortality in patients with heart failure: a pragmatic approach, **605**
- predictors of mortality**, Predictors of mortality in patients with acute coronary syndrome undergoing percutaneous coronary intervention, **1245**
- pregnancy**, Pregnancy related complications in women with hypertrophic cardiomyopathy, **752**
- prehospital**, GREAT: 10 year survival of patients with suspected acute myocardial infarction in a randomised comparison of prehospital and hospital thrombolysis, **563**
- premature death**, Family history of premature death from ischaemic heart disease is associated with an increased risk of delivering a low birth weight baby, **1249**
- prescribing patterns**, Evolution of statin prescribing 1994–2001: a case of agism but not of sexism?, **417**
- pressure**, Assessment of myocardial viability using coronary zero flow pressure after successful angioplasty in patients with acute anterior myocardial infarction, **71**
- prevalence**, Heart failure and the aging population: an increasing burden in the 21st century?, **49**
- Prevalence of impaired left ventricular systolic function and heart failure in a middle aged and elderly urban population segment of Copenhagen, **1422**
- primary pulmonary hypertension**, Lack of association between angiotensin converting enzyme (ACE) genotype, serum ACE activity, and haemodynamics in patients with primary pulmonary hypertension, **445**
- prognosis**, Creatinine clearance and adverse hospital outcomes in patients with acute coronary syndromes: findings from the global registry of acute coronary events (GRACE), **1003**
- N-terminal pro B type natriuretic peptide, but not the new putative cardiac hormone relaxin, predicts prognosis in patients with chronic heart failure, **879**
- Prognosis for patients newly admitted to hospital with heart failure: survival trends in 12 220 index admissions in Leicestershire 1993–2001, **615**
- Prognostic and clinical correlates of angiographically diffuse non-obstructive coronary lesions, **1009**
- Progression of familial and non-familial dilated cardiomyopathy: long term follow up, **757**
- Serum magnesium aberrations in furosemide (frusemide) treated patients with congestive heart failure: pathophysiological correlates and prognostic evaluation, **411**
- programmed stimulation**, Comparative follow up of patients with implanted cardioverter-defibrillators after induction of sustained monomorphic ventricular tachycardias or ventricular fibrillation by programmed stimulation, **629**
- prostaglandin**, Increased urinary 15-F_{2t}-isoprostane concentrations in patients with non-ischaemic congestive heart failure: a marker of oxidative stress, **871**
- prostheses**, Twenty year comparison of a Bjork-Shiley mechanical heart valve with porcine bioprostheses, **715**
- prosthetic valve endocarditis**, Management of prosthetic valve endocarditis: a clinical challenge, **245**
- Prosthetic valve endocarditis: early and late outcome following medical or surgical treatment, **269**
- protection devices**, Carotid artery stenting, **944**
- protection tools**, Radiation exposure to cardiologists: how it could be reduced, **1123**
- protein losing enteropathy**, Acquired combined immunodeficiency associated with protein losing enteropathy complicating Fontan operation, **1130**
- proximal LAD intervention**, Coronary stenting versus coronary bypass surgery in patients with multiple vessel disease and significant proximal LAD stenosis: results from the ERACI II study, **184**
- pseudoaneurysm**, Long term outcome of patients with postinfarction left ventricular pseudoaneurysm, **1144**
- psychological aspects**, The psychological perspective: a professional view, **ii16**
- public place defibrillators**, The debate on public place defibrillators: charged but shockingly ill informed, **1375**

- pulmonary arterial hypertension**, Acute endothelin A receptor antagonism improves pulmonary and systemic haemodynamics in patients with pulmonary arterial hypertension that is primary or autoimmune and related to congenital heart disease, **1221**
Chronic cor pulmonale, **225**
- pulmonary artery**, Low pressure giant pulmonary artery aneurysms in the adult: natural history and management strategies, **1067**
- pulmonary artery pulsatility**, Intravascular ultrasound of the elastic pulmonary arteries: a new approach for the evaluation of primary pulmonary hypertension, **311**
- pulmonary artery stenosis**, The efficacy and safety of stent redilatation in congenital heart disease, **905**
- pulmonary atresia**, Prevalence of haemoptysis in adults with pulmonary atresia and ventricular septal defect, and the role of mammary artery collateral vessels, **937**
- pulmonary autograft**, Excessive pulmonary autograft dilatation causes important aortic regurgitation, **933**
- pulmonary embolism**, Short term clinical outcome of acute saddle pulmonary embolism, **335**
- pulmonary hypertension**, Atrial septostomy for pulmonary arterial hypertension, **1344**
Intravascular ultrasound of the elastic pulmonary arteries: a new approach for the evaluation of primary pulmonary hypertension, **311**
Modified technique of stent fenestration of the atrial septum, **1227**
- pulmonary regurgitation**, Low pressure giant pulmonary artery aneurysms in the adult: natural history and management strategies, **1067**
- pulmonary thromboendarterectomy**, Pulmonary thromboendarterectomy almost 50 years after the first surgical attempts, **369**
- pulse wave velocity**, Arterial dysfunction in syndrome X: results of arterial reactivity and pulse wave propagation tests, **422**
- QRS width**, Interventricular and intra-left ventricular electromechanical delays in right ventricular paced patients with heart failure: implications for upgrading to biventricular stimulation, **1401**
- QT dispersion**, QT dispersion and left ventricular morphology in patients with hypertrophic cardiomyopathy, **882**
- QTc**, Long QTc predicts future cardiac death in stroke survivors, **377**
- quality of life**, Quality of life assessment using the short form 12 questionnaire is as reliable and sensitive as the short form 36 in distinguishing symptom severity in myocardial infarction survivors, **1445**
- racial differences**, Bangladeshi patients present with non-classic features of acute myocardial infarction and are treated less aggressively in east London, UK, **276**
- radiation exposure**, Radiation exposure benefit of a lead cap in invasive cardiology, **1205**
- radiation safety**, Radiation exposure benefit of a lead cap in invasive cardiology, **1205**
- radiofrequency ablation**, Comparison of radiofrequency versus cryotherapy catheter ablation of septal accessory pathways, **1091**
- radionuclide ventriculography**, A meta-analysis comparing the prognostic accuracy of six diagnostic tests for predicting perioperative cardiac risk in patients undergoing major vascular surgery, **1327**
- Raman spectroscopy**, On-line detection of cholesterol and calcification by catheter based Raman spectroscopy in human atherosclerotic plaque ex vivo, **1078**
- randomised controlled trials**, Women, older persons, and ethnic minorities: factors associated with their inclusion in randomised trials of statins 1990 to 2001, **327**
- rapamycin**, In-stent stenosis: pathology and implications for the development of drug eluting stents, **218**
- β receptors**, The effects of α and β blockade on ventilatory responses to exercise in chronic heart failure, **1169**
- redilatation**, The efficacy and safety of stent redilatation in congenital heart disease, **905**
- re-entry**, Successful radiofrequency catheter ablation of "clockwise" and "counterclockwise" bundle branch re-entrant ventricular tachycardia in the absence of myocardial or valvar dysfunction without detecting bundle branch potentials, **e12**
- refractory angina**, Enhanced external counterpulsation for refractory angina pectoris, **830**
- rehabilitation**, Comprehensive cardiac rehabilitation programme for implantable cardioverter defibrillator patients: a randomised controlled trial, **155**
- relaxin**, N-terminal pro B type natriuretic peptide, but not the new putative cardiac hormone relaxin, predicts prognosis in patients with chronic heart failure, **879**
Transcardiac and transpulmonary gradients in the putative new cardiovascular hormone relaxin, **789**
- remodelling**, Angiotensin II receptor blockade prevents microangiopathy and preserves diastolic function in the diabetic rat heart, **1236**
Impact of microvascular integrity and local viability on left ventricular remodelling after reperfused acute myocardial infarction, **393**
Influence of contractile reserve and inducible ischaemia on left ventricular remodelling after acute myocardial infarction, **1138**
Mechanism of lumen enlargement with direct stenting versus predilatation stenting: influence of remodelling and plaque characteristics assessed by volumetric intracoronary ultrasound, **84**
Relation between heart rate, heart rhythm, and reverse left ventricular remodelling in response to carvedilol in patients with chronic heart failure: a single centre, observational study, **293**
- renal artery stenosis**, Flash pulmonary oedema: accurate evaluation of the renal arteries with multislice computed tomography, **1251**
- renal dysfunction**, Creatinine clearance and adverse hospital outcomes in patients with acute coronary syndromes: findings from the global registry of acute coronary events (GRACE), **1003**
Minor renal dysfunction: an emerging independent cardiovascular risk factor, **963**
- renin-angiotensin system**, Epistatic interaction between variations in the angiotensin I converting enzyme and angiotensin II type 1 receptor genes in relation to extent of coronary atherosclerosis, **1195**
- reperfusion**, Assessing perfusion and function in acute myocardial infarction: how and when?, **701**
Myocardial contrast echocardiography is superior to other known modalities for assessing myocardial reperfusion after acute myocardial infarction, **139**
Relation of coronary flow pattern to myocardial blush grade in patients with first acute myocardial infarction, **1147**
- reperfusion injury**, Impact of microvascular integrity and local viability on left ventricular remodelling after reperfused acute myocardial infarction, **393**
- reserve**, Impact of microvascular integrity and local viability on left ventricular remodelling after reperfused acute myocardial infarction, **393**
- restenosis**, Clinical and angiographic acute and follow up results of intracoronary β brachytherapy in saphenous vein bypass grafts: a subgroup analysis of the multicentre European registry of intraluminal coronary β brachytherapy (RENO), **640**
Drug eluting stents: maximising benefit and minimising cost, **127**
High preprocedural non-HDL cholesterol is associated with enhanced oxidative stress and monocyte activation after coronary angioplasty: possible implications in restenosis, **773**
Histopathology of coronary in-stent restenosis following γ brachytherapy, **11**
Successful brachytherapy of coronary vasospasm, **e25**
The CARDia trial protocol, **1125**
- resuscitation**, Resuscitation, **673**
- reteplase**, Thrombolysis: newer thrombolytic agents and their role in clinical medicine, **1358**
- revascularisation**, Associations between ST depression, four year mortality, and in-hospital revascularisation in unselected patients with non-ST elevation acute coronary syndromes, **490**
How many cath labs do we need?, **827**
Indications for percutaneous and surgical revascularisation: how far does the evidence base guide us?, **565**
Long term effects of nisoldipine on the progression of coronary atherosclerosis and the occurrence of clinical events: the NICOLE study, **887**
White blood cell count and long term mortality after non-ST elevation acute coronary syndrome treated with very early revascularisation, **389**
- right bundle branch**, Successful radiofrequency catheter ablation of "clockwise" and "counterclockwise" bundle branch re-entrant ventricular tachycardia in the absence of myocardial or valvar dysfunction without detecting bundle branch potentials, **e12**

- right heart failure**, Sisters with atypical Fabry's disease with complete atrioventricular block, **e2**
- right sided endocarditis**, Injection drug use and right sided endocarditis, **577**
- right ventricular function**, Tissue Doppler, strain, and strain rate echocardiography for the assessment of left and right systolic ventricular function, **iii9**
- risk assessment**, A meta-analysis comparing the prognostic accuracy of six diagnostic tests for predicting perioperative cardiac risk in patients undergoing major vascular surgery, **1327**
Risk assessment in coronary artery surgery, **1**
- risk estimation**, Global risk of cardiovascular disease, **ii2**
The applicability of the Framingham coronary heart disease prediction function to black and minority ethnic groups in the UK, **785**
- risk factors**, Are there sex differences in risk factors for coronary heart disease? Maternal versus paternal transmission, **817**
Cardiovascular risk factors in patients with aortic stenosis predict prevalence of coronary artery disease but not of aortic stenosis: an angiographic pair matched case-control study, **1019**
Global risk of cardiovascular disease, **ii2**
Minor renal dysfunction: an emerging independent cardiovascular risk factor, **963**
QRS duration: a simple marker for predicting cardiac mortality in ICD patients with heart failure, **1157**
Risk factors for pacemaker implantation following aortic valve replacement: a single centre experience, **901**
- risk management**, The nurse as expert practitioner in global cardiovascular risk management, **ii33**
- risk markers**, Cumulative risk assessment in unstable angina: clinical, electrocardiographic, autonomic, and biochemical markers, **36**
- risk model**, The risk model of choice for coronary surgery in the UK, **98**
- risk prediction**, Estimating prognosis in heart failure: time for a better approach, **587**
- risk stratification**, Contemporary considerations for risk stratification, sudden death and prevention in hypertrophic cardiomyopathy, **977**
Cumulative risk assessment in unstable angina: clinical, electrocardiographic, autonomic, and biochemical markers, **36**
The risk model of choice for coronary surgery in the UK, **98**
Validation of four different risk stratification systems in patients undergoing off-pump coronary artery bypass surgery: a UK multicentre analysis of 2223 patients, **432**
- saphenous vein bypass graft**, Clinical and angiographic acute and follow up results of intracoronary β brachytherapy in saphenous vein bypass grafts: a subgroup analysis of the multicentre European registry of intraluminal coronary β brachytherapy (RENO), **640**
- sarcoplasmic reticulum**, The therapeutic potential of new insights into myocardial excitation-contraction coupling, **371**
- Schaumann body**, A 28 year old woman with ventricular tachycardia and an abnormal chest radiograph, **e19**
- scintigraphy**, Usefulness of myocardial perfusion SPECT in patients with left bundle branch block and previous myocardial infarction, **1039**
- secular trend**, Trends in incidence and case fatality rates of acute myocardial infarction in Denmark and Sweden, **507**
- secundum atrial septal defect**, Experience with transcatheter closure of secundum atrial septal defects using the Amplatzer septal occluder: a single centre study in 236 consecutive patients, **199**
- seizure**, Differential diagnosis of cardiogenic syncope and seizure disorders, **353**
- septal accessory pathway**, Comparison of radiofrequency versus cryotherapy catheter ablation of septal accessory pathways, **1091**
- septation**, Development of the heart: (2) Septation of the atriums and ventricles, **949**
- septostomy**, Atrial septostomy for pulmonary arterial hypertension, **1344**
- serum glucose**, A single serum glucose measurement predicts adverse outcomes across the whole range of acute coronary syndromes, **512**
- serum potassium**, Effect of diabetes on serum potassium concentrations in acute coronary syndromes, **31**
- SF-12**, Quality of life assessment using the short form 12 questionnaire is as reliable and sensitive as the short form 36 in distinguishing symptom severity in myocardial infarction survivors, **1445**
- SF-36**, Quality of life assessment using the short form 12 questionnaire is as reliable and sensitive as the short form 36 in distinguishing symptom severity in myocardial infarction survivors, **1445**
- short protocol**, Diagnostic accuracy of a new shorter dobutamine infusion protocol in stress echocardiography, **1089**
- shortening fraction**, Restrictive transmitral filling patterns predict improvements in left ventricular function after biventricular pacing, **1087**
- Silzone valve**, Incidence of embolism and paravalvar leak after St Jude Silzone valve implantation: experience from the Cardiff Embolic Risk Factor Study, **1055**
- simvastatin**, Simvastatin attenuates leucocyte-endothelial interactions after coronary revascularisation with cardiopulmonary bypass, **538**
- sinoatrial re-entry tachycardia**, Catheter ablation of sinoatrial re-entry tachycardia in a 2 month old infant, **e1**
- sinus node**, The relative contribution of the sinus and AV node to heart rate variability, **337**
- sirolimus**, Drug eluting stents: maximising benefit and minimising cost, **127**
- smoking**, Adipocyte derived plasma protein, adiponectin, is associated with smoking status in patients with coronary artery disease, **663**
Smoking: can we really make a difference?, **ii25**
- smooth muscles**, Dissection of the aorta: a new approach, **6**
- socioeconomic deprivation**, Socioeconomic deprivation is a predictor of poor postoperative cardiovascular outcomes in patients undergoing coronary artery bypass grafting, **1062**
- sodium channel blockers**, Abnormal response to sodium channel blockers in patients with Brugada syndrome: augmented localised wall motion abnormalities in the right ventricular outflow tract region detected by electron beam computed tomography, **169**
- sodium-hydrogen exchanger**, Increased platelet sodium-hydrogen exchanger activity in patients with variant angina, **935**
- sotalol**, Management of fetal tachyarrhythmia based on superior vena cava/aorta Doppler flow recordings, **1211**
- SPECT**, Role of myocardial perfusion imaging for risk stratification in suspected or known coronary artery disease, **1291**
- splenic abscess**, Persisting fever in a patient with brucella endocarditis: occult splenic abscess, **e20**
- sporting event**, Lower myocardial infarction mortality in French men the day France won the 1998 World Cup of football, **555**
- ST recovery time**, Improving the positive predictive value of exercise testing in women, **1416**
- stable angina**, The elusive link between stenosis severity and prognosis in stable ischaemic heart disease, **961**
- statins**, A rational basis for selection among drugs of the same class, **687**
Evolution of statin prescribing 1994-2001: a case of agism but not of sexism?, **417**
Long term statin treatment reduces lipoprotein(a) concentrations in heterozygous familial hypercholesterolaemia, **893**
Plaque stabilisation in acute coronary syndromes: clinical considerations, **1268**
Women, older persons, and ethnic minorities: factors associated with their inclusion in randomised trials of statins 1990 to 2001, **327**
- stenosis**, Emergency coronary stenting of unprotected critical left main coronary artery stenosis in acute myocardial infarction and cardiogenic shock, **e24**
Pathophysiology of coronary microembolisation, **981**
- stents**, Biocorrosion of magnesium alloys: a new principle in cardiovascular implant technology?, **651**
Clinical and angiographic acute and follow up results of intracoronary β brachytherapy in saphenous vein bypass grafts: a subgroup analysis of the multicentre European registry of intraluminal coronary β brachytherapy (RENO), **640**
Clinical and angiographic outcome of directional atherectomy followed by stent implantation in de novo lesions located at the ostium of the left anterior descending coronary artery, **1050**
Drug eluting stents: are human and animal studies comparable?, **133**
Evaluation of intracoronary stenting by intravascular optical coherence tomography, **317**
Histopathology of coronary in-stent restenosis following β brachytherapy, **11**
Mechanism of lumen enlargement with direct stenting versus predilatation stenting: influence of remodelling and plaque characteristics assessed by volumetric intracoronary ultrasound, **84**
Myocardial viability, coronary flow reserve, and in-hospital predictors of late recovery of contractility following successful primary stenting for acute myocardial infarction, **179**
Prospective evaluation of the effect of an angiotensin I converting enzyme gene polymorphism on the long term risk of major adverse cardiac events after percutaneous coronary intervention, **321**

- Same day discharge following elective percutaneous coronary intervention in patients with stable angina, **665**
- Stenting of the ductus arteriosus and banding of the pulmonary arteries: basis for various surgical strategies in newborns with multiple left heart obstructive lesions, **645**
- The adult patient with native coarctation of the aorta: balloon angioplasty or primary stenting?, **77**
- The CARDia trial protocol, **1125**
- The efficacy and safety of stent redilatation in congenital heart disease, **905**
- stent fenestration**, Modified technique of stent fenestration of the atrial septum, **1227**
- stent graft**, Endovascular treatment of thoracic aortic disease, **823**
- Open surgery for thoracic aortic disease, **825**
- stenting**, Carotid artery stenting, **944**
- Emergency coronary stenting of unprotected critical left main coronary artery stenosis in acute myocardial infarction and cardiogenic shock, **e24**
- In-stent stenosis: pathology and implications for the development of drug eluting stents, **218**
- strain**, Tissue Doppler, strain, and strain rate echocardiography for the assessment of left and right systolic ventricular function, **iii9**
- strain rate**, Tissue Doppler, strain, and strain rate echocardiography for the assessment of left and right systolic ventricular function, **iii9**
- stress**, Stress and myocardial infarction, **475**
- stress echocardiography**, A meta-analysis comparing the prognostic accuracy of six diagnostic tests for predicting perioperative cardiac risk in patients undergoing major vascular surgery, **1327**
- Diagnostic accuracy of a new shorter dobutamine infusion protocol in stress echocardiography, **1089**
- Influence of contractile reserve and inducible ischaemia on left ventricular remodelling after acute myocardial infarction, **1138**
- Stress echocardiography, **113**
- Transoesophageal atrial pacing combined with transthoracic two dimensional echocardiography: experience in patients operated on with arterial switch operation for transposition of the great arteries, **91**
- stress myocardial contrast echocardiography**, Stress myocardial contrast echocardiography, **1391**
- stroke**, Ethnic differences in cardiovascular disease, **681**
- Frequent atrial premature contractions as a surrogate marker for paroxysmal atrial fibrillation in patients with acute ischaemic stroke, **1247**
- Long QTc predicts future cardiac death in stroke survivors, **377**
- structural heart disease**, Concordance for hypoplastic left heart syndrome in a monozygotic twin pregnancy, **e13**
- stunning**, Apical ballooning of the left ventricle: first series in white patients, **1027**
- sudden death**, Catecholaminergic polymorphic ventricular tachycardia: electrocardiographic characteristics and optimal therapeutic strategies to prevent sudden death, **66**
- Contemporary considerations for risk stratification, sudden death and prevention in hypertrophic cardiomyopathy, **977**
- Heart rate variability in children with hypertrophic cardiomyopathy, **205**
- Risk of competitive sport in young athletes with heart disease, **710**
- single coronary artery**, Single coronary artery: a familial clustering, **e27**
- supravalvar aortic stenosis**, Williams syndrome associated with complete atrioventricular septal defect, **e15**
- supraventricular tachycardia**, Flecainide induced ventricular fibrillation in a neonate, **1261**
- Review of diagnosis, treatment, and outcome of fetal atrial flutter compared with supraventricular tachycardia, **913**
- surgery**, Accuracy of reporting cardiac related events during long term follow up, **331**
- Cardiac surgery and the brain: differences between adult and paediatric studies, **365**
- Improved postoperative outcomes with stentless aortic valve: a community hospital experience, **551**
- Optimal management of acute ventricular septal rupture, **1462**
- Pericardial effusion after open heart surgery for congenital heart disease, **780**
- Primary cardiac valve tumours, **1259**
- Timing of surgery in mitral regurgitation, **100**
- surgical strategy**, Low pressure giant pulmonary artery aneurysms in the adult: natural history and management strategies, **1067**
- surgical treatment**, Prosthetic valve endocarditis: early and late outcome following medical or surgical treatment, **269**
- survival**, Atrial septostomy for pulmonary arterial hypertension, **1344**
- N-terminal pro B type natriuretic peptide, but not the new putative cardiac hormone relaxin, predicts prognosis in patients with chronic heart failure, **879**
- Survival differences between heart failure in general practices and in hospitals, **1298**
- Twenty year comparison of a Bjork-Shiley mechanical heart valve with porcine bioprostheses, **715**
- Sweden**, Trends in incidence and case fatality rates of acute myocardial infarction in Denmark and Sweden, **507**
- symptom severity**, Quality of life assessment using the short form 12 questionnaire is as reliable and sensitive as the short form 36 in distinguishing symptom severity in myocardial infarction survivors, **1445**
- syncope**, Differential diagnosis of cardiogenic syncope and seizure disorders, **353**
- syndrome X**, Arterial dysfunction in syndrome X: results of arterial reactivity and pulse wave propagation tests, **422**
- Regional myocardial perfusion defects during exercise, as assessed by three dimensional integration of morphology and function, in relation to abnormal endothelium dependent vasoreactivity of the coronary microcirculation, **517**
- systemic right ventricle**, Impaired myocardial blood flow and coronary flow reserve of the anatomical right systemic ventricle in patients with congenitally corrected transposition of the great arteries, **1231**
- ystole**, Cardiac function and obesity, **1127**
- Diastolic heart failure: fact or fiction?, **1281**
- systolic dimension**, Preoperative left ventricular end systolic dimension as a predictor of postoperative ventricular dysfunction in children with mitral regurgitation, **1243**
- systolic dysfunction**, NT-proBNP: a new diagnostic screening tool to differentiate between patients with normal and reduced left ventricular systolic function, **150**
- systolic left ventricular function**, Effects of isolated obesity on systolic and diastolic left ventricular function, **1152**
- Prevalence of impaired left ventricular systolic function and heart failure in a middle aged and elderly urban population segment of Copenhagen, **1422**
- Tissue Doppler, strain, and strain rate echocardiography for the assessment of left and right systolic ventricular function, **iii9**
- tachyarrhythmias**, A 28 year old woman with ventricular tachycardia and an abnormal chest radiograph, **e19**
- Review of diagnosis, treatment, and outcome of fetal atrial flutter compared with supraventricular tachycardia, **913**
- Takayasu's arteritis**, Takayasu's arteritis: management of left main stem stenosis, **e9**
- tako-tsubo-like left ventricular dysfunction**, Apical ballooning of the left ventricle: a distinct entity?, **974**
- tecteplase**, Thrombolysis: newer thrombolytic agents and their role in clinical medicine, **1358**
- testosterone**, Cardiovascular effects of testosterone: implications of the "male menopause"?, **121**
- tetralogy of Fallot**, Challenges in the care of adult patients with congenital heart defects, **465**
- Long term somatic growth after repair of tetralogy of Fallot: evidence for restoration of genetic growth potential, **1340**
- thalassaemia major**, Left ventricular remodelling, and systolic and diastolic function in young adults with β thalassaemia major: a Doppler echocardiographic assessment and correlation with haematological data, **762**
- thallasaemia**, Early detection of left ventricular dysfunction in patients with β thalassaemia major by dobutamine stress echocardiography, **669**
- thoracic aneurysm**, Endovascular treatment of thoracic aortic disease, **823**
- thoracic aorta**, Open surgery for thoracic aortic disease, **825**
- three dimensional echocardiography**, Left atrial and left ventricular function in healthy children and young adults assessed by three dimensional echocardiography, **544**
- Left ventricular assessment using real time three dimensional echocardiography, **iii29**
- thrombocytopenia**, Deaths associated with platelet glycoprotein IIb/IIIa inhibitor treatment, **535**

- thromboembolic events**, Experience with enoxaparin in patients with mechanical heart valves who must withhold acenocumarol, **527**
- thromboembolic risk factors**, Experience with enoxaparin in patients with mechanical heart valves who must withhold acenocumarol, **527**
- thromboembolism**, Atrial fibrillation: relation between clinical risk factors and transoesophageal echocardiographic risk factors for thromboembolism, **165**
- thrombogenesis**, Angiogenesis, thrombogenesis, endothelial dysfunction and angiographic severity of coronary artery disease, **1411**
Platelet activation, haemorrhology and thrombogenesis in acute atrial fibrillation: a comparison with permanent atrial fibrillation, **1093**
- thrombogenic milieu**, Atrial fibrillation: relation between clinical risk factors and transoesophageal echocardiographic risk factors for thromboembolism, **165**
- thrombolysis**, Abortion of acute ST segment elevation myocardial infarction after reperfusion: incidence, patients' characteristics, and prognosis, **496**
Are we underestimating the full potential of early thrombolytic treatment in patients with acute myocardial infarction?, **483**
Thrombolysis: newer thrombolytic agents and their role in clinical medicine, **1358**
- thrombolytic agents**, Thrombolysis: newer thrombolytic agents and their role in clinical medicine, **1358**
- thrombolytic therapy**, GREAT: 10 year survival of patients with suspected acute myocardial infarction in a randomised comparison of prehospital and hospital thrombolysis, **563**
Plaque stabilisation in acute coronary syndromes: clinical considerations, **1268**
- thrombophilia**, Learning on the web: Case 2: Patent foramen ovale (PFO) and paradoxical embolism, **1362**
- thrombosis**, Basic aspects of plaque vulnerability, **1262**
Inflammatory and thrombotic mechanisms in coronary atherosclerosis, **993**
- thrombus**, The role of echocardiography in atrial fibrillation and cardioversion, **1447**
- tidal volume**, Safety and potential benefits of low dose diamorphine during exercise in patients with chronic heart failure, **1085**
- time factors**, GREAT: 10 year survival of patients with suspected acute myocardial infarction in a randomised comparison of prehospital and hospital thrombolysis, **563**
- tissue Doppler echocardiography**, Tissue Doppler, strain, and strain rate echocardiography for the assessment of left and right systolic ventricular function, **iii9**
- tissue Doppler imaging**, Clinical applications of tissue Doppler imaging: a promise fulfilled, **1377**
- tissue engineering**, Biocorrosion of magnesium alloys: a new principle in cardiovascular implant technology?, **651**
- toad venom poisoning**, Toad venom poisoning: resemblance to digoxin toxicity and therapeutic implications, **e14**
- tolerability**, Obstacles to the initiation of β blockers for heart failure in a specialised clinic within a district general hospital, **442**
- tomography**, Evaluation of intracoronary stenting by intravascular optical coherence tomography, **317**
- Toronto SPV valve**, Improved postoperative outcomes with stentless aortic valve: a community hospital experience, **551**
- Torsades de pointes**, Drug induced QT prolongation and torsades de pointes, **1363**
- toxicology**, Toad venom poisoning: resemblance to digoxin toxicity and therapeutic implications, **e14**
- training**, Radiation exposure to cardiologists: how it could be reduced, **1123**
- transcatheter closure**, Experience with transcatheter closure of secundum atrial septal defects using the Amplatzer septal occluder: a single centre study in 236 consecutive patients, **199**
Transcatheter closure of perimembranous ventricular septal defects with the Amplatzer asymmetric ventricular septal defect occluder: preliminary experience in children, **918**
- transient left ventricular apical ballooning**, Apical ballooning of the left ventricle: a distinct entity?, **974**
- transoesophageal atrial pacing**, Transoesophageal atrial pacing combined with transthoracic two dimensional echocardiography: experience in patients operated on with arterial switch operation for transposition of the great arteries, **91**
- transoesophageal atrial stimulation**, Fontan conversion to total cavopulmonary connection and arrhythmia ablation: clinical and functional results, **193**
- transoesophageal echocardiography**, Atrial fibrillation: relation between clinical risk factors and transoesophageal echocardiographic risk factors for thromboembolism, **165**
Injection drug use and right sided endocarditis, **577**
Management of prosthetic valve endocarditis: a clinical challenge, **245**
Primary cardiac valve tumours, **1259**
- transposition of the great arteries**, Cardiovascular malformations in infants of diabetic mothers, **1217**
Transoesophageal atrial pacing combined with transthoracic two dimensional echocardiography: experience in patients operated on with arterial switch operation for transposition of the great arteries, **91**
- transradial cardiac procedures**, Implementation of a diagnostic and interventional transradial programme: resource and organisational implications, **561**
- transthoracic Doppler echocardiography**, Intramyocardial coronary flow characteristics in patients with hypertrophic cardiomyopathy: non-invasive assessment by transthoracic Doppler echocardiography, **657**
- transthoracic echocardiography**, Injection drug use and right sided endocarditis, **577**
Judicious use of transthoracic echocardiography in the diagnosis of infective endocarditis, **1283**
- trends**, Heart failure and the aging population: an increasing burden in the 21st century?, **49**
Trends and ethnic differences in hospital admissions and mortality for congestive heart failure in the elderly in Singapore, 1991 to 1998, **865**
- tricuspid regurgitation**, Tricuspid regurgitation in the diagnosis of chromosomal anomalies in the fetus at 11–14 weeks of gestation, **1071**
- triglycerides**, Lipid lowering to delay the progression of coronary artery disease, **451**
- troponin**, C reactive protein for risk stratification in acute coronary syndromes? Verdict: unproven, **1288**
- troponin T**, Circulating concentrations of cardiac proteins indicate the severity of congestive heart failure, **1303**
Measurements of cardiac troponin T in patients with hypertrophic cardiomyopathy, **659**
- tumour necrosis factor α** , Therapeutic inhibition of tumour necrosis factor α in patients with heart failure: cooling an inflamed heart, **14**
Tumour necrosis factor α concentration and collateral flow in patients with coronary artery disease and normal systolic left ventricular function, **96**
- twin pregnancies**, Concordance for hypoplastic left heart syndrome in a monochorionic twin pregnancy, **e13**
- ultrasonic stethoscope**, The "ultrasonic stethoscope": is it of clinical value?, **704**
- ultrasound**, Ultrasound stethoscopy: a renaissance of the physical examination?, **971**
- ultrasound stethoscopy**, Ultrasound stethoscopy: a renaissance of the physical examination?, **971**
- unstable angina**, Cumulative risk assessment in unstable angina: clinical, electrocardiographic, autonomic, and biochemical markers, **36**
International comparison of mortality rates in patients with non-ST elevation acute coronary events, **1083**
The central role of platelet activation in determining the severity of acute coronary syndromes, **1253**
The elusive link between stenosis severity and prognosis in stable ischaemic heart disease, **961**
- VA intervals**, Management of fetal tachyarrhythmia based on superior vena cava/aorta Doppler flow recordings, **1211**
- valve**, Improved postoperative outcomes with stentless aortic valve: a community hospital experience, **551**
- valve disease**, Primary cardiac valve tumours, **1259**
- valve prosthesis**, Incidence of embolism and paravalvar leak after St Jude Silzone valve implantation: experience from the Cardiff Embolic Risk Factor Study, **1055**
- variant angina**, Increased platelet sodium–hydrogen exchanger activity in patients with variant angina, **935**

- vascular brachytherapy**, Clinical and angiographic acute and follow up results of intracoronary β brachytherapy in saphenous vein bypass grafts: a subgroup analysis of the multicentre European registry of intraluminal coronary β brachytherapy (RENO), **640**
- vascular endothelial growth factor**, Decreased serum vascular endothelial growth factor concentrations in patients with congestive heart failure, **207**
- vascular injury**, Use of 3 French catheters for diagnostic and interventional procedures in newborns and small infants, **1350**
- vasodilatation**, Syncope: what is the trigger?, **123**
- vasomotor tone**, Effects of L-arginine on flow mediated dilatation induced by atrial pacing in diseased epicardial coronary arteries, **531**
- vasovagal reaction**, Syncope: what is the trigger?, **123**
- venous thrombosis**, Symptomatic occlusion of the access vein after pacemaker or ICD lead extraction, **1348**
- ventilation**, Attenuation of hypercapnic carbon dioxide chemosensitivity after postinfarction exercise training: possible contribution to the improvement in exercise hyperventilation, **404**
Pattern of ventilation during exercise in chronic heart failure, **610**
- ventilatory response**, The effects of α and β blockade on ventilatory responses to exercise in chronic heart failure, **1169**
- ventricles**, Development of the heart: (2) Septation of the atriums and ventricles, **949**
- ventricular fibrillation**, Catecholaminergic polymorphic ventricular tachycardia: electrocardiographic characteristics and optimal therapeutic strategies to prevent sudden death, **66**
Flecainide induced ventricular fibrillation in a neonate, **1261**
The debate on public place defibrillators: charged but shockingly ill informed, **1375**
- ventricular filling**, Clinical assessment of left ventricular diastolic function, **231**
- ventricular function**, Preoperative left ventricular end systolic dimension as a predictor of postoperative ventricular dysfunction in children with mitral regurgitation, **1243**
- ventricular outflow tracts**, Development of the heart: (3) Formation of the ventricular outflow tracts, arterial valves, and intrapericardial arterial trunks, **1110**
- ventricular pre-excitation**, Asymptomatic ventricular pre-excitation in children and adolescents: a 15 year follow up study, **215**
- ventricular remodelling**, Relation between circulating soluble Fas ligand and subsequent ventricular remodelling following myocardial infarction, **339**
- ventricular septal defect**, Challenges in the care of adult patients with congenital heart defects, **465**
Prevalence of haemoptysis in adults with pulmonary atresia and ventricular septal defect, and the role of mammary artery collateral vessels, **937**
- ventricular tachyarrhythmias**, Comparative follow up of patients with implanted cardioverter-defibrillators after induction of sustained monomorphic ventricular tachycardias or ventricular fibrillation by programmed stimulation, **629**
- ventricular tachycardia**, Catecholaminergic polymorphic ventricular tachycardia: electrocardiographic characteristics and optimal therapeutic strategies to prevent sudden death, **66**
Haemodynamic and catecholamine response to simulated ventricular tachycardia in man: effect of baseline left ventricular function, **306**
- vigorous physical activity**, What level of physical activity protects against premature cardiovascular death? The Caerphilly study, **502**
- von Willebrand factor**, Coronary angioplasty enhances platelet reactivity through von Willebrand factor release, **329**
- vulnerable plaque**, On-line detection of cholesterol and calcification by catheter based Raman spectroscopy in human atherosclerotic plaque ex vivo, **1078**
- warfarin**, Warfarin for non-valvar atrial fibrillation: still underused in the 21st century?, **553**
- weekly variation**, Variation during the week in the incidence of acute myocardial infarction: increased risk for Japanese women on Saturdays, **398**
- white blood cell count**, Impact of preprocedural white blood cell count on long term mortality after percutaneous coronary intervention: insights from the EPIC, EPILOG, and EPISTENT trials, **1200**
White blood cell count and long term mortality after non-ST elevation acute coronary syndrome treated with very early revascularisation, **389**
- white coat hypertension**, Ambulatory blood pressure monitoring in the management of hypertension, **571**
- Williams syndrome**, Williams syndrome associated with complete atrioventricular septal defect, **e15**
- women**, Improving the positive predictive value of exercise testing in women, **1416**
Women, older persons, and ethnic minorities: factors associated with their inclusion in randomised trials of statins 1990 to 2001, **327**
- young**, A prospective study of long term prognosis in young myocardial infarction survivors: the prognostic value of angiography and exercise testing, **843**
- young athlete**, Risk of competitive sport in young athletes with heart disease, **710**