

Systematic review of multidisciplinary interventions in heart failure

R Holland, J Battersby, I Harvey, E Lenaghan, J Smith, L Hay

Heart 2005;91:899–906. doi: 10.1136/hrt.2004.048389

See end of article for authors' affiliations

Correspondence to:
Dr Richard Holland,
School of Medicine, Health
Policy and Practice,
University of East Anglia,
Norwich, Norfolk NR4
7TJ, UK; r.holland@uea.
ac.uk

Accepted
8 November 2004

Objective: To determine the impact of multidisciplinary interventions on hospital admission and mortality in heart failure.

Design: Systematic review. Thirteen databases were searched and reference lists from included trials and related reviews were checked. Trial authors were contacted if further information was required.

Setting: Randomised controlled trials conducted in both hospital and community settings.

Patients: Trials were included if all, or a defined subgroup of patients, had a diagnosis of heart failure.

Interventions: Multidisciplinary interventions were defined as those in which heart failure management was the responsibility of a multidisciplinary team including medical input plus one or more of the following: specialist nurse, pharmacist, dietician, or social worker. Interventions were separated into four mutually exclusive groups: provision of home visits; home physiological monitoring or televideo link; telephone follow up but no home visits; and hospital or clinic interventions alone. Pharmaceutical and exercise based interventions were excluded.

Main outcome measures: All cause hospital admission, all cause mortality, and heart failure hospital admission.

Results: 74 trials were identified, of which 30 contained relevant data for inclusion in meta-analyses. Multidisciplinary interventions reduced all cause admission (relative risk (RR) 0.87, 95% confidence interval (CI) 0.79 to 0.95, $p = 0.002$), although significant heterogeneity was found ($p = 0.002$). All cause mortality was also reduced (RR 0.79, 95% CI 0.69 to 0.92, $p = 0.002$) as was heart failure admission (RR 0.70, 95% CI 0.61 to 0.81, $p < 0.001$). These results varied little with sensitivity analyses.

Conclusion: Multidisciplinary interventions for heart failure reduce both hospital admission and all cause mortality. The most effective interventions were delivered at least partly in the home.

Heart failure is a common and serious public health problem. Recent estimates suggest a prevalence of congestive heart failure in Europe of 2.3%.¹ There has been a steady increase in hospital admissions for heart failure with almost 80 000 recorded in the UK in 2000. The estimated total National Health Service (NHS) cost associated with heart failure in 2000 was £905 million—that is, 1.9% of NHS expenditure.²

Despite progress in heart failure treatments, such as increasing use of angiotensin converting enzyme (ACE) inhibitors and more recently β blockade, many agree that the overall management of heart failure can be improved.^{3–4} Since 1999, several reviews of heart failure disease management programmes have been published, including two meta-analyses.^{5–7} All reviews have concluded that specialised follow up of patients by a multidisciplinary team can reduce hospitalisation but have been unclear about the effect on survival. Since the most recent reviewers completed their search⁶ a further 11 randomised trials have reported results. These have substantially increased the number of randomly assigned patients from 3304 to 8158. It is therefore timely to review this larger body of evidence with its potential to provide more precise effect size estimates. Furthermore, we wished to investigate the relative merits of home, telemonitoring, telephone, and clinic or hospital based interventions. This question of the location of follow up has not been examined to date.

Home visits have the potential to allow educational interventions to be tailored to a patient's social circumstances. Telemonitoring gives people the freedom of their homes while permitting daily (or even continuous) evaluation of physiological state and often face to face contact

through videophone technology. Telephone follow up allows hospital advice to be re-enforced and promotes continuity of care⁸ but, although less time consuming, it is also less personal than a home visit or telemonitoring.

The objective of this systematic review of all available randomised controlled trials was to determine the impact of multidisciplinary interventions on hospital admission and mortality in patients with heart failure. Multidisciplinary interventions were defined as those in which management was the responsibility of a multidisciplinary team that included medical input plus one or more of the following: a specialist nurse, a pharmacist, a health educator, a dietician, or a social worker. Trials of drugs, pacemakers, mechanical appliances, and exercise-alone interventions were excluded.

METHODS

Data sources

The search strategy identified research on patients with congestive heart failure that tested multidisciplinary interventions and reported outcomes of admission, mortality, or quality of life. Multidisciplinary interventions were identified by a broad range of search terms and medical subject headings: home care services, patient or health education, community health nursing, patient care planning,

Abbreviations: ACE, angiotensin converting enzyme; AMED, Allied and Complementary Medicine; CI, confidence interval; CINAHL, Cumulative index to Nursing and Allied Health Literature; COACH, coordinating study evaluating outcomes of advising and counselling in heart failure; DARE, Database of Abstracts of Reviews of Effects; NHS, National Health Service; TEN-HMS, Trans-European Network initiative-homecare management system study

pharmaceutical services, and telemonitoring. This search was developed with reference to the indexing of previously identified studies. The following electronic databases were searched from their inception to 1 June 2004: Medline, Embase, Cumulative index to Nursing and Allied Health Literature (CINAHL), Allied and Complementary Medicine (AMED), Cochrane Controlled Trials Register, the Cochrane Effective Practice and Organization of Care Study Registry, Biomed, the TRIP Database, Meta-Register of Current Controlled Trials, Research Findings Electronic Register, NHS Research Register, the Cochrane Database of Systematic Reviews, and the Database of Abstracts of Reviews of Effects (DARE). Reference lists of all included articles and relevant review articles were searched. There were no language restrictions.

Study selection

One of three investigators (JB, RH, and EL) reviewed all titles retrieved by the literature search. Titles needed to appear potentially relevant to the study area. Two investigators (JB and RH) independently assessed abstracts against three criteria to determine, firstly, that the study was a randomised controlled trial; secondly, that a defined group of patients were identified as having heart failure; and thirdly, that the intervention was non-pharmaceutical and non-exercise based. Interventions needed to be applied to patients and not focus solely on changing physician management. Trials incorporating exercise advice among a range of other heart failure issues were included. Multidisciplinary interventions were nurse led programmes, medication review, medication adherence interventions, patient education, or enhanced monitoring. Studies could be carried out in any setting (hospital, clinic, home, etc). Full papers were retrieved if one or both investigators considered the abstract suitable. The two investigators then independently assessed retrieved studies for their suitability for inclusion in the meta-analysis. Differences were resolved by discussion with reference to a third reviewer (IH) if necessary.

Data extraction

Data were extracted (JB, RH, IH, EL) on to an electronic database based on a previously piloted data extraction form. Extracted data were type of participants, intervention details, outcomes, and trial quality characteristics.

Classification of interventions

Two researchers (JS and LH), who were blinded to all trial results, independently classified trial interventions and then agreed their decisions jointly. Studies were classified into one or more of the following definitions:

- (1) One or more planned intervention visits in the patient's home by a health care professional to educate or improve patient self management (this excludes visits to take blood samples, set up physiological monitoring, or carry out interventions such as wound care)
- (2) Videophone use or any form of home physiological monitoring
- (3) One or more planned telephone calls to the patient at home
- (4) Educational or symptom self management mailings to the patient's home address
- (5) No planned intervention home visits; instead, intervention was provided during hospital admission or during hospital clinic attendance
- (6) No planned intervention home visits; instead, intervention was provided in general practice or a community clinic.

For the meta-analyses, each trial was categorised according to the uppermost category into which it fitted. Thus, a trial classified as having an intervention delivered in hospital (5) with additional planned telephone follow up (3) was classified as category 3. Furthermore, as only one trial was classified in category 4, this was merged with category 3. Similarly, categories 5 and 6 were merged. This resulted in four final intervention subgroups: subgroup A entailed at least some home visiting; subgroup B, home physiological monitoring or televideo contact; subgroup C, telephone or mailing to the patient's home without home visiting; and subgroup D interventions were delivered exclusively in a hospital, clinic, or general practice. These four subgroups were mutually exclusive.

Trial interventions were also classified as high or low intensity. High intensity interventions consisted of planned contact at least monthly with multifaceted interventions; low intensity interventions entailed less frequent contact or narrow interventions (for example, medication review alone).

Outcomes

The primary outcome was the proportion of patients with one or more hospital admission (all cause). Secondary outcomes were all cause mortality, heart failure hospital admission, and mean inpatient days. Data were extracted from studies for their predefined final follow up. Data on quality of life were also extracted but are not presented here owing to the diversity of scales used, which prevented data pooling.

Quality assessment

Many quality scales have been created to judge trials. These scales often give importance to subject blinding, which was impossible within the trials considered by this review. Juni *et al*⁹ recommend estimating trial quality by assessing three key components: concealment of allocation, use of intention to treat analysis, and blinding of outcome assessment. Blinding of outcome assessment was rarely reported for our outcomes (hospital admission or death). We therefore assessed whether studies confirmed outcome data by using at least two sources (for example, hospital data and self report). In addition, trial quality was assessed against criteria recommended by the York Centre for Reviews and Dissemination¹⁰: firstly, explicit statement of inclusion or exclusion criteria; secondly, baseline comparability between groups; and thirdly, a clearly defined primary outcome. The review team also considered the following criteria important in reference to this particular study area: clear diagnostic criteria for heart failure and length of follow up (where six months and over was considered adequate). This gave a total of eight quality criteria against which studies were assessed.

Data synthesis

All trials that reported appropriate data were included in the meta-analyses if interventions were compared with usual care. Authors of trials were contacted to confirm the data extraction and additional data were sought where articles did not report one or more outcomes. For data on proportion of patients admitted (all cause and heart failure) and mortality, the effect of the intervention was reported as a relative risk. For mean inpatient days a weighted mean difference was calculated. Meta-analyses were carried out with the random effects methods in RevMan version 4.2 (Cochrane Collaboration, Oxford, UK). Funnel plots were constructed to assess possible publication bias.

Sensitivity analyses

The robustness of the findings in relation to all cause admission and mortality was investigated in the following sensitivity analyses. Firstly, we investigated reasons for any




Figure 1 Flowchart describing study selection and excluded studies. RCT, randomised controlled trial.

heterogeneity found and repeated analyses with fixed effects methods. Next we explored the effect of excluding poorer quality studies. This was investigated in two ways: by removing those lacking components recommended by Juni *et al*⁹ and by removing those achieving 50% or less of the eight criteria above. Trials where randomisation was not of individual patients (that is, cluster trials) were also excluded from analysis. These report overprecise results if adjustment is not made for the clustering of data. In the context of a meta-analysis such data can potentially bias results. Analyses were also repeated excluding trials yet to formally publish results in a peer reviewed article. Lastly, analyses were repeated excluding trials that recruited broadly, in which patients with heart failure were simply a subgroup.

RESULTS

Search results

We reviewed 9132 titles, which yielded 667 potentially relevant studies. Seventy four studies met the original inclusion criteria (fig 1). Of these, 21 contained sufficient

published data to be included in the meta-analysis and a further nine were included after contact with authors yielded additional data.^{8 11–18} Forty four studies were excluded. Figure 1 lists reasons for exclusion.

Study characteristics

Of the 30 trials included in this review 13 were performed in Europe, 13 in the USA, three in Australia or New Zealand, and one in Argentina (table 1; to view table 1 visit the *Heart* website—<http://www.heartjnl.com/supplemental>). Twenty two trials (73%) were published in the past five years (1999–2004). The earliest included trial was published in 1993.¹⁹ Patient inclusion criteria were reasonably similar across all studies. Four trials recruited patients from a hospital clinic or general practice^{12 20–22} and all others recruited hospitalised (or recently hospitalised) patients admitted with a diagnosis of heart failure. In almost all studies patients were excluded if they were due to be discharged to a long term care facility, had some form of cognitive impairment or psychiatric disorder, or had a

terminal disease or severe co-morbidity. Despite the apparently similar inclusion criteria the mean age of patients varied between 56–86 years and the proportion of men from 27–99%. Fifteen trials recorded baseline ejection fractions, with trial means varying from 22–43%. Eight studies focused their intervention on a category of more severe heart failure (for example, New York Heart Association functional class > II¹⁵).

Trial quality

Only six studies clearly satisfied all three key quality components (concealed allocation, intention to treat, and data cross checked).^{11 23–27} However, when trials were considered against all eight quality criteria the majority (22 of 30) met at least five. Two trials achieved only two criteria,^{12 28} five achieved three criteria,^{13 16 19 22 29 30} and one achieved four criteria.³¹ It should be noted that for one study only an abstract was available, which limited its quality assessment.¹³

Interventions

Twelve trials had a home based component (subgroup A) (table 2; to view table 2 visit the *Heart* website—<http://www.heartjnl.com/supplemental>). Home interventions included between one and eight visits over trial follow up (mean of 0.6 visits for each month of follow up). In two trials patients were visited only if they could not attend a clinic.^{18 25} All but two of these 12 interventions also included telephone

contact.^{12 32} Three trials included remote physiological monitoring (subgroup B). Twelve trials had telephone interventions or letters to patients²⁸ (subgroup C). Patients received between one and 17 telephone calls over follow up (mean 1.4 for each month of follow up). Three trials delivered interventions exclusively in hospital or clinic settings (subgroup D).^{24 29 30}

Effect on all cause admission

Twenty one trials provided data on all cause admission (fig 2). No data were available from home monitoring trials on this outcome. Meta-analysis showed a significant reduction in all cause admission (relative risk 0.87, 95% confidence interval (CI) 0.79 to 0.95, $p = 0.002$). However, significant heterogeneity ($p = 0.002$) was found. Subgroup analysis showed that home visit interventions reduced all cause admission to hospital (relative risk 0.80, 95% CI 0.71 to 0.89, $p < 0.0001$). Telephone-type interventions also reduced admission but this finding was of borderline significance (relative risk 0.86, 95% CI 0.73 to 1.02, $p = 0.09$). Hospital based interventions had no effect (relative risk 0.99, 95% CI 0.90 to 1.10, $p = 0.56$). Heterogeneity was found among telephone-type interventions ($p < 0.001$) but not within the two other subgroups.

Effect on all cause mortality

Twenty seven trials provided data on all cause mortality (fig 3). Meta-analysis showed a significant decrease in all cause mortality (relative risk 0.79, 95% CI 0.69 to 0.92,




Figure 2 Forest plot for all cause admission results. CI, confidence interval; RR, relative risk. Note: data on all cause admission were not available for the three trials in subgroup B (videophone/remote monitoring).




Figure 3 Forest plot of all cause mortality results. The control group from the study of Cleland/Coletta appears twice, as this was a three arm trial (telemonitoring, telephone support, and usual care). As a result, estimates for each subgroup are correctly calculated but the total estimate is slightly inaccurate. This should read 0.79 (95% CI 0.69 to 0.92) and the overall test for heterogeneity should have a χ^2 of 40.07, $p = 0.04$.

$p = 0.002$). Subgroup analysis showed significant reductions in mortality for both telemonitoring (relative risk 0.49, 95% CI 0.33 to 0.73, $p < 0.001$) and telephone follow up (relative risk 0.70, 95% CI 0.53 to 0.94, $p = 0.02$). Reductions in mortality were non-significant in the home (relative risk 0.87, 95% CI 0.72 to 1.06, $p = 0.44$) and clinic subgroups (relative risk 1.00, 95% CI 0.84 to 1.20, $p = 0.98$). All subgroup confidence intervals overlapped, with the exception of subgroups B and D. Heterogeneity was detected for the overall result ($p = 0.04$) and for the telephone subgroup ($p = 0.04$).

Effect on heart failure admission

Sixteen trials provided data on heart failure admission. Meta-analysis showed a significant reduction in heart failure admission with a relative risk of 0.70 (95% CI 0.61 to 0.81, $p < 0.0001$). The benefit was notable and similar for home

and telephone-type interventions, with relative risks of 0.62 (95% CI 0.51 to 0.74, $p < 0.001$) and 0.70 (95% CI 0.57 to 0.85, $p < 0.001$), respectively. No significant effect was apparent in the two studies of patients receiving a hospital or community based intervention (relative risk 0.94, 95% CI 0.78 to 1.13, $p = 0.51$).^{24, 29} No heterogeneity was detected within the subgroups, but it was detected for the overall result ($p = 0.04$).

Effect on total admissions and mean inpatient days

Multidisciplinary interventions appear to reduce not only the proportion of patients admitted but also the total number of admissions. Across 18 of the studies the intervention groups had a mean admission rate of 21.8 admissions versus 29.0/1000 patient weeks for control groups. These data should be interpreted with caution, as formal pooling was not possible due to lack of standard deviation data. The 10 trials that

provided mean inpatient days and standard deviation data (seven home visits, three telephone interventions) were formally pooled. This showed a significant decrease in overall mean inpatient days of 1.9 days in favour of the intervention (95% C.I. 0.71 to 3.1, $p = 0.002$), with no overall heterogeneity ($p = 0.24$).

Sensitivity analyses

Table 3 describes the various sensitivity analyses conducted on the primary outcome (all cause admissions). The overall effect was slightly reduced by considering a fixed effect estimate (relative risk 0.90 ν 0.87). Both results were highly significant ($p = 0.002$). However, in both analyses significant heterogeneity was found. This was principally caused by two outliers. One suggested very strong benefit³¹ and the other potential harm.²⁷ The study showing a very strong benefit provided intensive heart failure support through a day hospital.³¹ Patients could be managed there as day patients with intravenous treatment if necessary, and these events did not then count as admissions. Such care was not available in any other study. The second outlier recruited a wide spectrum of patients (with heart failure and other diagnoses) and delivered an intervention focused particularly on medication before and immediately after discharge.²⁷ Removing these two studies, which provided qualitatively different interventions from all others, reduced heterogeneity notably ($p = 0.39$) but only slightly reduced the overall effectiveness estimate (relative risk 0.89).

Table 3 also shows that sensitivity analyses around quality components had little impact on the effectiveness estimate. The largest impact came from limiting the meta-analysis to studies that clearly used concealed allocation. In this case the relative risk was reduced to 0.92. In contrast to this, when quality was considered across eight trial components higher quality studies (judged as achieving five or more components) showed increased effectiveness (relative risk 0.85 ν

0.94). A similar result was found when cluster trials were excluded, indicating that their inclusion could only have biased results towards no effect. In one of these trials the analysis had accounted for clustering but adjusted data required for the meta-analysis could not be extracted.²⁴ A final quality indicator used was in terms of peer review publication of data. Data from eight studies were considered "unpublished". In these cases, data were either provided directly by the author, extracted from conference abstracts, or from web based material. Relative risk estimates were identical between meta-analysis of these data and published data (0.87 ν 0.86).

Among those providing admission data, 12 trials had high intensity interventions. These studies were from all our intervention categories (that is, home, telemonitoring, telephone, and hospital). These appeared only marginally more effective than low intensity interventions (relative risk 0.86 ν 0.88), with overlapping confidence intervals. A clearer though non-significant difference was evident when mortality was considered (relative risk 0.76 ν 0.84).

Three studies recruited patients with a variety of diagnoses and applied more generic interventions.^{11 27 32} Together these studies did not show a benefit within their subgroup of patients with heart failure (relative risk 1.02). In contrast, studies solely recruiting patients with heart failure found reduced risk of admission, with little difference between those recruiting high risk patients and those that were less selective (relative risk 0.82 ν 0.86).

Identical sensitivity analyses were also performed for mortality results (table 3). These found broadly similar results to those described above for all cause admission.

Funnel plots

Funnel plots (available from the authors) suggested little evidence of publication bias.

Table 3 Sensitivity analyses

Analysis	All cause admission			Mortality		
	No of patients (no of trials)	RR (95% CI)	p Value	No of patients (no of trials)	RR (95% CI)	p Value
Random versus fixed effects						
Random effects	6569 (21)	0.87 (0.79 to 0.95)	0.002	7447 (27)	0.79 (0.69 to 0.92)	0.002
Fixed effects	6569 (21)	0.90 (0.85 to 0.95)	<0.001	7447 (27)	0.85 (0.77 to 0.94)	0.001
Removing outliers						
Base case	6569 (21)	0.87 (0.79 to 0.95)	0.002	7447 (27)	0.79 (0.69 to 0.92)	0.002
Removing extreme trials	5831 (19)	0.89 (0.83 to 0.94)	<0.001	7213 (26)	0.83 (0.73 to 0.95)	0.006
Quality component						
1. Concealed allocation	2238 (10)	0.92 (0.80 to 1.05)	0.22	3161 (11)	0.87 (0.74 to 1.01)	0.07
No clear concealment	4331 (11)	0.82 (0.73 to 0.95)	0.01	4286 (16)	0.74 (0.58 to 0.94)	0.01
2. ITT	5363 (14)	0.88 (0.79 to 0.99)	0.03	5589 (17)	0.76 (0.63 to 0.91)	0.003
Unclear if ITT used	1206 (7)	0.83 (0.73 to 0.94)	0.002	1858 (10)	0.86 (0.67 to 1.12)	0.26
3. Data cross checked	3859 (13)	0.86 (0.75 to 0.99)	0.03	2587 (13)	0.86 (0.72 to 1.03)	0.10
No clear cross check	2710 (8)	0.89 (0.80 to 0.98)	0.02	4860 (14)	0.72 (0.69 to 0.92)	0.007
Overall quality score						
High score (>5 of 8)	4577 (16)	0.85 (0.75 to 0.95)	0.005	4793 (20)	0.82 (0.71 to 0.96)	0.01
Low score (\leq 5 of 8)	1992 (5)	0.94 (0.86 to 1.04)	0.50	2654 (7)	0.69 (0.45 to 1.05)	0.08
Level of randomisation						
Individual patient only	4510 (18)	0.84 (0.75 to 0.94)	0.002	5388 (24)	0.76 (0.64 to 0.90)	0.002
Cluster trial	2059 (3)	0.97 (0.89 to 1.07)	0.56	2059 (3)	0.99 (0.83 to 1.18)	0.92
Origin of data						
Published article	4315 (13)	0.86 (0.75 to 0.98)	0.02	4740 (21)	0.83 (0.81 to 0.97)	0.02
Data from author/abstract	2254 (8)	0.87 (0.79 to 0.95)	0.003	2707 (6)	0.70 (0.48 to 1.02)	0.06
Intensity						
High	3593 (11)	0.85 (0.75 to 0.97)	0.02	4550 (15)	0.76 (0.63 to 0.92)	0.005
Low	2976 (10)	0.88 (0.76 to 1.01)	0.07	2897 (12)	0.84 (0.66 to 1.06)	0.15
Target of intervention						
High risk CHF patient	1354 (7)	0.82 (0.69 to 0.96)	0.02	1586 (7)	0.81 (0.58 to 1.15)	0.25
Any CHF patient	4543 (11)	0.86 (0.78 to 0.96)	0.007	5693 (18)	0.79 (0.67 to 0.94)	0.008
Variety of diagnoses	672 (3)	1.02 (0.72 to 1.44)	0.91	168 (2)	0.73 (0.36 to 1.51)	0.40

CHF, congestive heart failure; CI, confidence interval; ITT, intention to treat; RR, relative risk.

DISCUSSION

This systematic review shows that delivering multidisciplinary interventions to patients with heart failure not only reduces hospital admission but also is an effective method for reducing mortality. Our results show that the risk of all cause admission is reduced by about 13%, mortality by as much as 20%, and heart failure admission by 30%. Results for all cause admission and mortality were little changed by a wide variety of sensitivity analyses that excluded trials on the basis of quality criteria or clustering of data. This is the first meta-analysis of this form of intervention to show a significant reduction in mortality, and it is important to note that the reduction is similar to that achieved by ACE inhibitors in heart failure.⁴⁰

This study also investigated whether effectiveness varied according to where the interventions were carried out. These subgroup analyses generated less clear cut results. Home based interventions successfully reduced all cause admission, heart failure admission, and mean days in hospital. However, they led to a non-significant decrease in deaths. The latter result may simply reflect that insufficient patients have been enrolled in home based trials as yet ($n = 1909$). In contrast, telephone interventions decreased heart failure admission and deaths but led to a non-significant decrease in all cause admission. Remote televideo or physiological monitoring provides a newer form of intervention. A few trials of these interventions have now been conducted. Although this review suggests that these interventions may have an important effect on mortality, data were not available to investigate their effect on admission. We found only three trials where interventions were solely delivered in a hospital, clinic, or general practice.^{24 29 30} None of these studies reported significant benefits.

Studies that we reviewed had reasonably broad inclusion criteria such that the findings are of relevance to most hospitalised patients with heart failure. However, trials tended to restrict participation to patients with no important co-morbidity, terminal disease, confusion, or residence in a long term care facility. Thus, the benefits may not extend to these groups. Equally, it should be noted that trial populations were reasonably heterogeneous, which is likely to have contributed to heterogeneity in the results.

This review has been enhanced by the provision of extra unpublished data from some authors and data from abstracts of trials presented at international conferences. Exclusion of these non-peer reviewed data within our sensitivity analysis did not change the effectiveness estimates but did broaden the confidence limits.

Eleven studies were excluded because they did not report outcome data that could be used by this review. However, eight of these reported other outcomes favouring their intervention groups, while three reported no clear differences. Funnel plots suggested little evidence of publication bias. Nonetheless, given the numbers of patients now randomly assigned (> 8000), it seems unlikely that publication of small negative studies would have an important impact on any of the effectiveness estimates.

A diverse range of interventions could have met this review's inclusion criteria. However, in reality the trials that we found tested relatively similar interventions. Almost all interventions shared two key elements: one to one patient education concerning heart failure, medication, and diet and exercise advice; and symptom monitoring and self management advice. Education tended to be given over a number of encounters. It should be noted that 11 trials incorporated interventions that appeared to be of high intensity. Effectiveness estimates from these appeared to be only marginally greater than those of lower intensity interventions and the confidence intervals overlapped. Equally, targeting the

intervention at high risk patients with heart failure did not seem important. It is also interesting to note that trial results do not appear to have changed over time despite increased use of ACE inhibitors and introduction of low dose β blockade (fig 2). This may suggest that the effectiveness of multidisciplinary interventions is mediated not solely through better compliance with modern drug treatments but also through better symptom self management and lifestyle changes.

No trial included in this review provided any form of placebo intervention. It is difficult to imagine how this would be delivered in practice. It is therefore possible that the effectiveness observed is partly due to increased social contact. A more important limitation of much of the research to date has been the use of a single highly motivated specialist team. Replicating this in other settings, particularly in rural areas where distances to specialist centres are large, is likely to be difficult. It seems important to test whether less specialised nurses or community pharmacists are as effective as these highly specialised teams.

A few studies were excluded from this review because they compared two interventions and did not include a usual care subgroup. These shed some light on the importance of intervention intensity. Benatar *et al*⁴¹ compared telemonitoring with frequent home visiting (nine visits over five weeks, then as needed) of 216 patients over 12 months' follow up. This trial reported heart failure admission data and found a non-significant reduction from telemonitoring (38 admissions) compared with home visiting (63 admissions). Jerant *et al*⁴² randomly assigned 37 patients to telemonitoring, telephone follow up, or usual care (where all groups received two detailed home visits) and followed them up over six months. This study's sample size limited its findings. In total there were nine admissions in the telemonitoring group, five in the telephone group, and 15 in the usual care group ($p = 0.45$). Harrison *et al*⁴³ randomly assigned 192 patients to receive usual home nursing care, which consisted of assessment, monitoring and health teaching, or a transitional care intervention. The transitional care intervention was a comprehensive discharge programme with evidence based counselling and education. Harrison *et al*⁴³ found a non-significant decrease in the proportion of patients admitted after the transitional care intervention. Finally, Coletta *et al* (in the TEN-HMS (Trans-European Network initiative-homecare management system study)),¹³ whose interim results are included in this review, compared usual care with two interventions: a telephone intervention and a televideo intervention. This study has found an improvement in mortality in both intervention groups, but only a small non-significant difference between these groups (13% for telemonitoring *v* 15% for telephone). With the exception of the study of Jerant *et al*,⁴² these studies suggest a small incremental effect of intensive interventions, in line with the finding from our review comparing high intensity interventions versus low intensity interventions. However, this difference, as well as its relative cost effectiveness, needs to be more clearly established. Ongoing trials will add to this analysis over the next three years including the large COACH (coordinating study evaluating outcomes of advising and counselling in heart failure) trial,⁴⁴ which is randomly assigning 1050 patients to either usual care, basic education and support, or intensive education and support, as well as final results from the TEN-HMS study.¹³

In conclusion, it is apparently possible to achieve major reductions in admissions and deaths of patients with heart failure by implementing post-discharge interventions delivering patient education and symptom self management. These interventions appear to be particularly effective when they are at least partly delivered in a patient's own home through visits, telephone calls, or more advanced televideo techniques.

ACKNOWLEDGEMENTS

We thank Jessie McGowan for her kind help searching the Cochrane Effective Practice and Organisation of Care Study Registry. We also thank the following authors for providing further data from their trials or publications: Brian Bixby (for Mary Naylor), Socorro Capomolla, John Cleland, Alison Coletta, Inger Ekman, Wendy Gattis, Larry Goodyer, Margaret Harrison, Tiny Jaarsma, Anthony Jerant, Harlan Krumholz, Michael Rich, Barbara Riegel, and Anna Strömberg. RH was funded by MRC as a research fellow during the course of this review.



To view tables 1 and 2 visit the *Heart* website—<http://www.heartjnl.com/supplemental>.

Authors' affiliations

R Holland, J Battersby, I Harvey, E Lenaghan, J Smith, L Hay, School of Medicine, Health Policy and Practice, University of East Anglia, Norwich, Norfolk, UK

REFERENCES

- Davies M, Hobbs F, Davis R, et al. Prevalence of left-ventricular systolic dysfunction and heart failure in the echocardiographic heart of England screening study: a population based study. *Lancet* 2001;**358**:439–44.
- Stewart S, Jenkins A, Buchan S, et al. The current cost of heart failure to the National Health Service in the UK. *Eur J Heart Fail* 2002;**4**:361–71.
- Jessup M, Brozena S. Heart failure. *N Engl J Med* 2003;**348**:2007–18.
- Horowitz JD. Home-based intervention: the next step in treatment of chronic heart failure? *Eur Heart J* 2000;**21**:1807–9.
- McAlister FA, Lawson FM, Teo KK, et al. A systematic review of randomized trials of disease management programs in heart failure. *Am J Med* 2001;**110**:378–84.
- Phillips CO, Wright SM, Kern DE, et al. Comprehensive discharge planning with postdischarge support for older patients with congestive heart failure: a meta-analysis. *JAMA* 2004;**291**:1358–67.
- Rich MW. Heart failure disease management: a critical review. *J Card Fail* 1999;**5**:64–75.
- Riegel B, Carlson B, Kopp Z, et al. Effect of a standardized nurse case-management telephone intervention on resource use in patients with chronic heart failure. *Arch Intern Med* 2002;**162**:705–12.
- Juni P, Altman DG, Egger M. Assessing the quality of randomised controlled trials. In: *Systematic reviews in health care: meta-analysis in context*. London: BMJ Publishing Group, 2001.
- Centre for Reviews & Dissemination. *Undertaking systematic reviews of research on effectiveness*. York: University of York, 2001.
- Holland R, Lenaghan E, Harvey I, et al. Does home based medication review keep older people out of hospital? The HOMER randomised controlled trial. *BMJ* (in press).
- Goodyer LI, Miskelly F, Milligan P. Does encouraging good compliance improve patients' clinical condition in heart failure? *Br J Clin Pract* 1995;**49**:173–6.
- Coletta AP, Louis AA, Clark AL, et al. Clinical trials update from the European Society of Cardiology: CARMEN, EARTH, OPTIMAAL, ACE, TEN-HMS, MAGIC, SOLVD-X and PATH-CHF II. *Eur J Heart Fail* 2002;**4**:661–6.
- Rainville EC. Impact of pharmacist interventions on hospital readmissions for heart failure. *Am J Health Syst Pharm* 1999;**56**:1339–42.
- Jaarsma T, Halfens R, Huijter Abu-Saad H, et al. Effects of education and support on self-care and resource utilization in patients with heart failure. *Eur Heart J* 1999;**20**:673–82.
- Rich MW, Beckham V, Wittenberg C, et al. A multidisciplinary intervention to prevent the readmission of elderly patients with congestive heart failure. *N Engl J Med* 1995;**333**:1190–5.
- Ekman I, Andersson B, Ehnfors M, et al. Feasibility of a nurse-monitored, outpatient-care programme for elderly patients with moderate-to-severe, chronic heart failure. *Eur Heart J* 1998;**19**:1254–60.
- Krumholz HM, Amatruda J, Smith GL, et al. Randomized trial of an education and support intervention to prevent readmission of patients with heart failure. *J Am Coll Cardiol* 2002;**39**:83–9.
- Rich MW, Vinson JM, Sperry JC, et al. Prevention of readmission in elderly patients with congestive heart failure: results of a prospective, randomized pilot study. *J Gen Intern Med* 1993;**8**:585–90.
- Gattis WA, Hasselblad V, Whellan DJ, et al. Reduction in heart failure events by the addition of a clinical pharmacist to the heart failure management team: results of the pharmacist in heart failure assessment recommendation and monitoring (PHARM) study. *Arch Intern Med* 1999;**159**:1939–45.
- Grancelli H, Varini S, Ferrante D, et al. Randomized trial of telephone intervention in chronic heart failure (DIAL): study design and preliminary observations. *J Card Fail* 2003;**9**:172–9.
- De Lusignan S, Wells S, Johnson P, et al. Compliance and effectiveness of 1 year's home telemonitoring: the report of a pilot study of patients with chronic heart failure. *Eur J Heart Fail* 2001;**3**:723–30.
- Blue L, Lang E, McMurray JJ, et al. Randomised controlled trial of specialist nurse intervention in heart failure. *BMJ* 2001;**323**:715–8.
- Doughty RN, Wright SP, Pearl A, et al. Randomized, controlled trial of integrated heart failure management: the Auckland heart failure management study. *Eur Heart J* 2002;**23**:139–46.
- Kasper EK, Gerstenblith G, Hefter G, et al. A randomized trial of the efficacy of multidisciplinary care in heart failure outpatients at high risk of hospital readmission. *J Am Coll Cardiol* 2002;**39**:471–80.
- Naylor MD, Brooten DA, Campbell RL, et al. Transitional care of older adults hospitalized with heart failure: a randomized, controlled trial. *J Am Geriatr Soc* 2004;**52**:675–84.
- Weinberger M, Oddone EZ, Henderson WG. Does increased access to primary care reduce hospital readmissions? Veterans Affairs Cooperative Study Group on Primary Care and Hospital Readmission. *N Engl J Med* 1996;**334**:1441–7.
- Serxner S, Miyajima M, Jeffords J. Congestive heart failure disease management study: a patient education intervention. *Prev Manage Congest Heart Fail* 1998;**4**:23–8.
- Philbin EF, Rocco TA, Lindenmuth NW, et al. The results of a randomized trial of a quality improvement intervention in the care of patients with heart failure. The MISCHF Study Investigators. *Am J Med* 2000;**109**:443–9.
- Varma S, McElnay JC, Hughes CM, et al. Pharmaceutical care of patients with congestive heart failure: interventions and outcomes. *Pharmacotherapy* 1999;**19**:860–9.
- Capomolla S, Febo O, Ceresa M, et al. Cost/utility ratio in chronic heart failure: comparison between heart failure management program delivered by day-hospital and usual care. *J Am Coll Cardiol* 2002;**40**:1259–66.
- Stewart S, Pearson S, Horowitz JD. Effects of a home-based intervention among patients with congestive heart failure discharged from acute hospital care. *Arch Intern Med* 1998;**158**:1067–72.
- Bouvy ML, Heerdink ER, Urquhart J, et al. Effect of a pharmacist-led intervention on diuretic compliance in heart failure patients: a randomized controlled study. *J Card Fail* 2003;**9**:404–11.
- Cline CM, Israelsson BY, Willenheimer RB, et al. Cost effective management programme for heart failure reduces hospitalisation. *Heart* 1998;**80**:442–6.
- Goldberg LR, Piette JD, Walsh MN, et al. Randomized trial of a daily electronic home monAS, Levinsky SK, Sargent J, et al. Case management in a heterogeneous congestive heart failure population in patients with advanced heart failure: the weight monitoring in heart failure (WHARF) trial. *Am Heart J* 2003;**146**:705–12.
- Laramee AS, Levinsky SK, Sargent J, et al. Case management in a heterogeneous congestive heart failure population: a randomized controlled trial. *Arch Intern Med* 2003;**163**:809–17.
- McDonald K, Ledwidge M, Cahill J, et al. Elimination of early rehospitalization in a randomized, controlled trial of multidisciplinary care in a high-risk, elderly heart failure population: the potential contributions of specialist care, clinical stability and optimal angiotensin-converting enzyme inhibitor dose at discharge. *Eur J Heart Fail* 2001;**3**:209–15.
- Stewart S, Marley JE, Horowitz JD. Effects of a multidisciplinary, home-based intervention on unplanned readmissions and survival among patients with chronic congestive heart failure: a randomised controlled study. *Lancet* 1999;**354**:1077–83.
- Stromberg A, Martensson J, Fridlund B, et al. Nurse-led heart failure clinics improve survival and self-care behaviour in patients with heart failure: results from a prospective, randomised trial. *Eur Heart J* 2003;**24**:1014–23.
- Garg R, Yusuf S. Overview of randomized trials of angiotensin-converting enzyme inhibitors on mortality and morbidity in patients with heart failure. Collaborative Group on ACE Inhibitor Trials. *JAMA* 1995;**273**:1450–6.
- Benatar D, Bondmass M, Ghitelman J, et al. Outcomes of chronic heart failure. *Arch Intern Med* 2003;**163**:347–52.
- Jerant AF, Azari R, Nesbitt TS. Reducing the cost of frequent hospital admissions for congestive heart failure: a randomized trial of a home telecare intervention. *Med Care* 2001;**39**:1234–45.
- Harrison MB, Browne GB, Roberts J, et al. Quality of life of individuals with heart failure: a randomized trial of the effectiveness of two models of hospital-to-home transition. *Med Care* 2002;**40**:271–82.
- Jaarsma T, Van Der Wal MH, Hogenhuis J, et al. Design and methodology of the COACH study: a multicenter randomised coordinating study evaluating outcomes of advising and counselling in heart failure. *Eur J Heart Fail* 2004;**6**:227–33.

Table 1: Included studies – baseline data on trial participants and trial quality

Author & Year	Setting	Total number	Participants			Follow-up (weeks)	Quality criteria achieved ¹ (Max = 8)	Notes regarding design/quality
			Mean age (years)	% Male	Mean ejection fraction			
Blue 2001 ²³	UK	165	75.0	58%	-	52	8	
Bouvy 2003 ³³	Netherlands	152	69.7	66%	-	26	8	
Capomolla 2002 ³¹	Italy	234	56.0	84%	30.0%	52	4	No details of allocation concealment, outcome data from single source only, no clear primary outcome, no clear inclusion/exclusion criteria.
Cline 1998 ³⁴	Sweden	190	75.6	53%	34.0%	52	5	No details of allocation concealment, no clear primary outcome, ITT not specified. NB: Randomised consent procedure used. 16 intervention patients refused consent vs. 0 controls.
Cleland / Coletta 2002 ¹³	Europe	426	-	-	-	57	3	Abstract only available. No details of allocation concealment, ITT not specified, outcome data not clearly cross-checked, baseline comparability unclear, inclusion/exclusion criteria not specified
de Lusignan 2001 ²²	UK	20	-	-	-	52	3	No details of allocation concealment, outcome data from single source only, no clear primary outcome, ITT not specified, no data on baseline comparability.
Doughty 2002 ²⁴	New Zealand	197	73.0	60%	32.2%	52	8	Cluster RCT, small cluster size (mean = 1.5 patients/cluster), authors adjusted for effect of clustering and found no effect on results
Ekman 1998 ¹⁷	Sweden	158	80.3	58%	40.5%	26	6	Outcome data from single source only, no clear primary outcome
Gattis 1999 ²⁰	USA	181	67.2	68%	30%	24	6	No details of allocation concealment, ITT not specified
Goldberg 2003 ³⁵	USA	280	59.1	68%	22%	26	8	
Goodyer 1995 ¹²	UK	100	84.5	27%	-	13	2	Criteria for CHF diagnosis not specified, no details of allocation concealment, outcome data from single source only, no clear primary outcome, ITT not specified, follow-up < 6 months.

Table 1: Included studies – baseline data on trial participants and trial quality (cont)

Author & Year	Setting	Total number	Participants			Follow-up (weeks)	Quality criteria achieved ¹ (Max = 8)	Notes regarding design/quality
			Mean age (years)	% male	Mean ejection fraction			
Grancelli 2003 ²¹	Argentina	1518	65	71%	-	52 (mean)	7	Unclear how outcome data were gathered.
Holland 2004 ¹¹	UK	71	85.9	48%	-	26	6	Criteria for CHF solely determined by discharge diagnosis
Jaarsma 1999 ¹⁵	Netherlands	179	73.0	58%	34.4%	39	7	ITT not specified. NB: 7 subjects excluded post-randomisation.
Kasper 2002 ²⁵	USA	200	61.9	61%	27.3%	26	7	Criteria for diagnosis of CHF not specified
Krumholz 2002 ¹⁸	USA	88	73.8	57%	37.5%	52	6	No details of concealed allocation, groups appeared to differ at baseline
Laramee 2003 ³⁶	USA	287	70.7	54%	-	13	5	No mention of intention to treat, follow-up < 6 months, no details of concealed allocation
McDonald 2002 ³⁷	Ireland	98	69.0	66%	37.0%	12	5	No details of concealed allocation, outcome data from single source only, follow-up < 6 months
Naylor 2004 ²⁶	USA	239	76.0	43%	-	52	7	Validation of diagnosis of heart failure not specified
Philbin 2000 ²⁹	USA	1504	75.7	44%	39.6%	26	3	No concealed allocation, no clear data cross-checking, poor baseline comparability, inclusion/exclusion criteria not specified and diagnostic criteria for heart failure unclear.
Rainville 1999 ¹⁴	USA	34	70.0	50%	-	52	5	Outcome data from single source only, ITT not specified, groups appeared to differ at baseline. NB: 4 subjects excluded post-randomisation.
Rich 1993 ¹⁹	USA	98	79.3	41%	-	13	3	No details of concealed allocation, groups appeared to differ at baseline, outcome data from single source only, follow-up < 6m, ITT not specified. NB: subjects randomised 2:1 intervention vs. control
Rich 1995 ¹⁶	USA	282	78.6	37%	42.5%	13	5	Groups appeared to differ at baseline, outcome data from single source only, follow-up < 6 months

Table 1: Included studies – baseline data on trial participants and trial quality (cont)

Author & Year	Setting	Total number	Participants			Follow-up (weeks)	Quality criteria achieved ¹ (Max = 8)	Notes regarding design/quality
			Mean age (years)	% male	Mean ejection fraction			
Riegel 2002 ⁸	USA	358	73.9	49%	42.7%	26	5	Criteria for diagnosis of CHF not specified, no details of concealed allocation, not analysed by ITT. NB: cluster RCT. No adjustment for clustering but cluster size small (mean = 1.3 patients/cluster). 57 subjects excluded post-randomisation as physician refused intervention (29) or patient withdrew (28)
Serxner 1998 ²⁸	USA	109	71.0	48%	-	26	2	Criteria for diagnosis of CHF not specified, no details of concealed allocation, no data on baseline comparability, no clear primary outcome, outcome data from single source only, ITT not specified.
Stewart 1998 ³²	Australia	97	75.0	48%	38.5%	26	7	Outcome data from single source only. NB: part of larger study of elderly hospital discharges (total n=762).
Stewart 1999 ³⁸	Australia	200	75.7	62%	37.0%	26	7	Outcome data from single source only
Stromberg 2003 ³⁹	Sweden	106	77.5	61%	-	52	6	Unclear if allocation concealed beyond use of sealed envelopes, outcome data from single source only.
Varma 1999 ³⁰	UK	83	75.9	41%	-	52	3	No details of concealed allocation, groups appear to differ at baseline, no clear primary outcome, outcome data from single source only, ITT not specified.
Weinberger 1996 ²⁷	USA	504	62.8	99%	-	26	7	Criteria for diagnosis of CHF not specified. NB: this was sub-group within a large study also recruiting patients with diabetes or chronic obstructive pulmonary disease (total subjects= 1396)

CHF, congestive heart failure; ITT, intention to treat; RCT, randomised controlled trial.

¹Quality criteria used: concealed allocation; use of an intention to treat analysis; outcome data confirmed by using at least two sources; an explicit statement of inclusion or exclusion criteria; baseline comparability between groups; a clearly defined primary outcome; clear diagnostic criteria for heart failure; and length of follow-up (where 6 months and over was considered adequate).

Table 2: Description of interventions provided

Author	Intervention subgroup (category)	Intervention – detail	Intervention provided by	Educational component	Planned contacts/ month of follow-up	Intensity judgement
Blue ²³	A (1)	Home educational and support visits of decreasing frequency, optimising drugs and monitoring electrolytes, plus phone contact as needed and liaison with other professionals	Specialist nurse	CHF knowledge and treatment advice, diet & exercise, promotion of self-management.	1.5	high
Bouvy ³³	C (3,6)	Review in pharmacy and monthly phone follow-up. Intervention focused on drug use and reasons for non-compliance	Community pharmacist	Medication	1	low
Capomolla ³¹	C (3,5)	Heart failure management programme delivered in a day hospital. Tailored therapy, physical training & counselling, correction of risk factors and education. Follow-up determined by need / symptoms.	Cardiologist, nurses, physiotherapist, dietician, psychologist & social assistant.	Monitoring weight, fluid restriction, nutrition & exercise, promotion of self-management	Unclear	high
Cline ³⁴	A (1)	2x30 minute educational visits in hospital plus 1 hour group oral and video presentation for patients and family after discharge plus follow up at easy access nurse clinic. Medication organiser offered. Nurse available by phone. One scheduled visit by nurse at 8 months	Study nurse	CHF knowledge, treatment adherence, promotion of self-management & use of patient symptom diary	0.5	low

Table 2: Description of interventions provided (cont)

Author	Intervention sub-group (category)	Intervention – detail	Intervention provided by	Educational component	Planned contacts/ month of follow-up	Intensity judgement
Cleland / Coletta ¹³	B (2,3)	Home tele-monitoring (weight, BP, ECG recorded twice daily) or monthly phone-calls from a nurse	Cardiologist, specialist nurse	Unclear	unclear for tele-monitored group	high
de Lusignan ²²	B (2)	Remote monitoring and tele-consultation with clinicians via videophone in patients' homes. Weekly consult for 3-months then decreasing frequency	Study nurse & GPs, cardiologist physiologist	Unclear	2	high
Doughty ²⁴	D (5, 6)	4 individual and group outpatient education sessions plus liaison with, and 6 weekly follow up by, GPs & heart failure clinic.	Study nurse, cardiologist, GPs	Medication, exercise, diet, monitoring weight & self-management	1	high
Ekman ¹⁷	C (3,5)	Nurse-monitored outpatient clinic or monthly phone contact by nurse. Written information to patients & regular liaison with primary care providers.	Clinic nurse	Medication, diet & promotion of self-management & weighing	1	high

Table 2: Description of interventions provided (cont)

Author	Intervention sub-group (category)	Intervention – detail	Intervention provided by	Educational component	Planned contacts/ month of follow-up	Intensity judgement
Gattis ²⁰	C (3,5)	Educational session by pharmacist who also optimised therapy & liaised with physicians & provided phone follow-up at 2, 12 & 24 weeks.	Clinical pharmacist	Medication changes discussed and medication education	0.66	low
Goldberg ³⁵	B (2,3)	Home tele-monitoring (weight and symptom checking daily)	Specialist nurse	CHF knowledge, daily weight, dietary & fluid restriction, symptom self-management - to both groups in hospital	daily	high
Goodyer ¹²	A (1)	Three home visits focusing on medication. Provision of medication calendars & information leaflets.	Pharmacist	Counselling on correct use of medication	1	low
Grancelli ²¹	C (3)	Telephone intervention with 5 objectives: diet and drug therapy compliance, symptom monitoring, control of oedema/weight, daily exercise.	Specialist nurse	CHF knowledge, diet & drug compliance, self-control techniques, help seeking if symptoms worsen	>1	high
Holland ¹¹	A (1)	Home visit by pharmacist to perform medication review and educate to improve medication compliance.	Pharmacist	Focused on medication	0.33	low

Table 2: Description of interventions provided (cont)

Author	Intervention sub-group (category)	Intervention – detail	Intervention provided by	Educational component	Planned contacts/ month of follow-up	Intensity judgement
Jaarsma ¹⁵	A (1,3)	Educational programme for patient (& family) during hospital stay followed by phone call from nurse one week after discharge & one home visit to reinforce/continue education.	Study nurse	Education & promotion of self-management, sodium restriction, fluid balance & compliance	0.33	low
Kasper ²⁵	A (1,3)	Treatment plan devised by cardiologist. Regular phone calls (initially weekly then decreasing frequency) & monthly follow-up at clinic or home by nurse to implement plan & titrate medications as required. A pill sorter & written educational materials were provided.	Specialist nurse, cardiologist, primary physician	List of medications, 2g sodium-restricted diet & exercise advice.	3	high
Krumholz ¹⁸	A (1,3)	One hour educational session at home or clinic followed by regular telephone calls (initially weekly then decreasing frequency) from nurse to support/monitor patients.	Specialist nurse	CHF knowledge, medication, health behaviour & symptom awareness.	1.5	high
Laramee ³⁶	C (3,5)	Early discharge planning and co-ordination of care, individualised patient/family education, enhanced telephone surveillance/follow-up, and promotion of optimal CHF medication and dosages.	Specialist nurse	CHF knowledge, diet & fluid advice, risk factor modification, exercise, symptom monitoring, prognosis & counselling	3.33	high

Table 2: Description of interventions provided (cont)

Author	Intervention sub-group (category)	Intervention – detail	Intervention provided by	Educational component	Planned contacts/ month of follow-up	Intensity judgement
McDonald ³⁷	C (3,5)	Education from nurse and dietician during admission plus follow up phone calls from nurse on discharge (weekly for 3 months) plus two extra clinic visits to nurse.	Specialist nurse, dietician	Medication, weight monitoring, salt restriction, dietary review	1.5	high
Naylor ²⁶	A (1)	Daily in-hospital visits to help optimise care, eight or more home visits over 3-months, plus telephone availability. Multiple home visits included assessment, treatment adjustment (in collaboration with physician) and patient/care-giver education.	Specialist nurse	Unclear but appeared wide-ranging	0.66	high
Philbin ²⁹	D (5)	Broad quality improvement intervention with standardised admission orders, health professional education, written information for patients and a home care pathway.	nurse, physicians, administrators	Teaching aids were given to patients – unclear what domains were covered	unclear	low
Rainville ¹⁴	C (3,5)	Medication review pre-discharge plus education and provision of written information. Follow-up by phone at 3 and 7 days, 1, 3 & 12 months post-discharge.	Pharmacist & Specialist nurse	Medication, weight monitoring & risk modification	0.5	low

Table 2: Description of interventions provided (cont)

Author	Intervention sub-group (category)	Intervention – detail	Intervention provided by	Educational component	Planned contacts/ month of follow-up	Intensity judgement
Rich 1993 ¹⁹	A (1,3)	Education, dietician review, medication review and early visit by social worker pre-discharge. Daily hospital visits pre-discharge then 3 home visits by home care nurse in first week then decreasing frequency plus phone contact.	Specialist nurse, dietician, cardiologist, social worker, home care nurse	CHF and its treatment with medication cards given, dietary review, plus weighing advice	>1 (precise no. unclear)	high
Rich 1995 ¹⁶	A (1,3)	As above (Rich 1993)	As above	As above	>1	high
Riegel ⁸	C (3,4)	Regular phone calls from nurse (approx. weekly depending on need) after discharge with management guided by decision support software. Written educational material also mailed monthly.	Specialist nurse	Medication adherence and dietary advice, promotion of symptom monitoring	2	high
Serxner ²⁸	C (4)	4 mailings of information over 12 weeks post-discharge.	Unclear	Medication, lifestyle, and risk factor advice	0.66	low
Stewart 1998 ³²	A (1)	Counselling by nurse and/or pharmacist prior to discharge followed by home visit by nurse and pharmacist and referral for additional community-based support for those deemed to be in need.	Study nurse & pharmacist	Medication review and counselling	0.33	low

Table 2: Description of interventions provided (cont)

Author	Intervention sub-group (category)	Intervention – detail	Intervention provided by	Educational component	Planned contacts/ month of follow-up	Intensity judgement
Stewart 1999 ³⁸	A (1,3)	Home visit by cardiac nurse 7-14 days post-discharge, with further follow up & visits as necessary & two follow up calls	Specialist nurse	Adherence, fluid & sodium intake, exercise, psychosocial support	0.5	low
Stromberg ³⁹	C (3,5)	Visit to nurse-led heart failure clinic 2 weeks after discharge. During visit treatment was optimised (in consultation with cardiologist) and patient given tailored education. Telephone follow-up to provide psycho-social support	Specialist nurse	CHF and its treatment, diet, smoking cessation, exercise, vaccination advice, symptom monitoring	0.1 – 0.2 visits, unclear number of phone calls	low
Varma ³⁰	D (5)	Hospital-based education & provision of information by pharmacist to encourage self-monitoring plus liaison with physicians & community pharmacists. Advice repeated in clinic 3-monthly	Research pharmacist	Medication, diet and self-management, daily weighing	0.4	low
Weinberger ²⁷	C (3,5)	In-patient education by nurse and visit to hospital from primary care physician. Phone call follow-up by nurse post-discharge & review of treatment plan at clinic.	Primary care nurse, primary care physician	Provision of educational materials	0.5	low

Intervention categories:

1 = One or more planned intervention visits in the patient's home by a health care professional to educate or improve their self-management (this excludes visits simply to take blood samples or carry out interventions such as wound care). [A]

2 = Home physiological monitoring or tele-visual contact with patients. [B]

3 = One or more planned telephone call or videophone call to the patient at home. [C]

4 = Educational or symptom self-management mailings to patient's home address. [C]

5 = No planned intervention home visits, instead, intervention occurring during hospital admission or during hospital clinic attendance. [D]

6 = No planned intervention home visits, instead, intervention occurring in general practice or a community clinic. [D]

Note: if intervention was categorised as 3 & 5, this was considered to be in sub-group C.