

Subject index

- a7r5 vascular smooth muscle cells**, Testosterone inhibits extracellular calcium entry via voltage-gated calcium channels in A7r5 vascular smooth muscle cells at physiological concentrations, **A43**
- abciximab**, Evaluation of Novel dose of C7E3Fab (abciximab) On periprocedural release of Troponin I and Thrombin generation marker (PF1+2) in Radial access Coronary Stenting as Elective Day case procedure 'The ENFORCED study', **A38**
- ablation**, Ablation of difficult right sided accessory pathways (AP) aided by simple mapping of the tricuspid VALVE annulus (TVA) using a halo catheter, **A55**
Catheter ablation of atrial fibrillation: primary and long term efficacy for three approaches in a single centre, **A10**
Catheter location trial: a prospective randomised comparison of carto, NavX and fluoroscopy for catheter ablation, **A37**
Experience with 75 consecutive cryoablation procedures in a single UK tertiary centre, **A10**
Percutaneous transthoracic epicardial ablation (PTEA) for failed endocardial ablations: promising results of a new technique, **A54**
Rapid pulmonary vein (PV) electrical isolation using a novel focused ultrasound balloon catheter in patients with atrial fibrillation (AF), **A55**
Real-time integrated X-ray and magnetic resonance imaging (XMR)-A novel strategy to optimise precise catheter placement during pulmonary vein isolation, **A37**
The influence of atrial conduction, refractoriness, and volume on outcomes following pulmonary vein isolation (PVI), **A54**
- accessory pathway**, Ablation of difficult right sided accessory pathways (AP) aided by simple mapping of the tricuspid VALVE annulus (TVA) using a halo catheter, **A55**
- acute chest pain**, Rapid, safe, and effective management of acute chest pain in a dedicated chest pain assessment unit at St James's Hospital, Dublin, **A3**
- acute coronary syndrome**, Anaemia predisposes to heart failure in acute coronary syndromes independently of the severity of myocardial injury, **A7**
Cholesterol content of erythrocyte membranes may contribute to the development of acute coronary syndrome. A new paradigm regarding coronary artery disease instability?, **A45**
Diagnostic accuracy of non-invasive coronary angiography with multislice computed tomography in patients presenting with acute chest pain, **A15**
How long do patients wait with acute coronary syndromes for inter-hospital transfer in the UK? Results from the coronary heart disease collaborative national audit 2004, **A25**
Impact of secondary prevention medication on 30 day mortality for patients admitted to English and Welsh hospitals with acute coronary syndromes, **A25**
Inflammation and acute hyperglycaemia in diabetics with non-ST elevation acute coronary syndromes (ACS). Analyses from OPUS TIMI 16 and TACTICS TIMI 18, **A60**
Is it safe to discharge patients from accident and emergency using a rapid point of care triple cardiac marker test to rule out acute coronary syndrome in patients presenting with chest pain?, **A26**
Patients with non-ST elevation acute coronary syndromes associated with myonecrosis have a similar rate of death or MI as ST elevation MI at 6 months: An audit of all ACS admissions in a UK District General Hospital over 1 year, **A42**
Prognostic value of normal stress echocardiography in patients with suspected Non-ST elevation acute coronary syndrome and negative troponin, **A33**
Rapid reduction of inter hospital Acute Coronary Syndrome (ACS) transfer waiting times; the role of a Regional Transfer Unit (RTU), **A25**
The hospital burden of suspected acute coronary syndromes (ACS): recent trends, **A60**
- acute myocardial infarction**, Declining case fatality rates for acute myocardial infarction in south Asians and whites in the last 15 years, **A41**
Factors influencing left ventricular function following successful rescue percutaneous coronary intervention for failed thrombolysis in acute myocardial infarction, **A24**
Fasting plasma glucose fails to detect new diabetes or abnormal glucose tolerance (GI) in patients admitted with acute myocardial infarction (MI), **A47**
Improved accuracy of low power contrast echocardiography for the assessment of left ventricular remodeling compared with unenhanced harmonic echocardiography following acute myocardial infarction: comparison with cardiac magnetic resonance imaging, **A32**
Prediction of transmuralty of acute myocardial infarction—comparison between myocardial contrast echocardiography and radionuclide perfusion imaging, **A32**
Two years of primary coronary intervention for ST elevation myocardial infarction in a district general hospital, **A59**
- adenosine**, A prospective, randomised, controlled trial to study the effect of intra coronary injection of verapamil and adenosine on coronary blood flow during urgent percutaneous coronary intervention, **A23**
- adult**, Is exercise training beneficial in adults with congenital heart disease?, **A33**
- adult congenital heart disease**, Resynchronization therapy in adults with congenital heart disease, **A35**
- advanced heart failure**, Usage of implantable defibrillators and biventricular pacemakers in cardiac transplant candidates: a survey of UK transplant centres, **A48**
- AED**, Initial results from the national defibrillator programme (NDP), **A63**
- ageing**, Erythropoietin delays endothelial cell senescence by preserving telomere length, **A14**
- air pollution**, Cardiovascular effects of exposure to air pollutants—human exposure studies demonstrating an adverse influence on heart rate variability (HRV), **A67**
Combustion derived nanoparticulate impairs vascular function and endogenous fibrinolysis in man: an explanation for the increased cardiovascular mortality associated with air pollution, **A2**
- alcohol septal myocardial ablation**, Electrocardiography and cardiac magnetic resonance imaging (MRI) before and after alcohol septal myocardial ablation (ASA) in hypertrophic obstructive cardiomyopathy, **A70**
- aldosterone**, Actions of aldosterone blockade in patients with mild-moderately severe heart failure, **A20**
Aldosterone mediates angiotensin II induced interstitial cardiac fibrosis via a Nox2 containing NADPH oxidase, **A3**
Genetic variation at the locus encompassing 11-beta hydroxylase and aldosterone synthase accounts for heritability in cortisol precursor (11-deoxycortisol) urinary metabolite excretion, **A65**
- ambulance**, Direct 24-hour ambulance triage and transfer of patients with acute myocardial infarction (AMI) to a cardiac intervention centre with no accident and emergency department, **A59**
- ambulance transfer**, Logistics and economics of interhospital transfer v direct admission for primary angioplasty, **A22**
Optimising target vessel reperfusion times with primary percutaneous intervention for acute myocardial infarction, **A23**
- AMI**, The REACT (REscue Angioplasty v Conservative treatment or repeat Thrombolysis) Trial: results according to actual treatment received, **A22**
- Amiloride**, Effects of amiloride and spironolactone on QT in stroke survivors—a randomised placebo controlled double-blind cross over study, **A65**
- AMP-activated protein kinase**, Delineating the role of adenosine monophosphate (AMP)-activated protein kinase (AMPK) γ 2 subunit in hypertrophic cardiomyopathy (HCM), **A69**
- anaemia**, Anaemia predisposes to heart failure in acute coronary syndromes independently of the severity of myocardial injury, **A7**
- angina**, Contemporary prognosis of angina and non-cardiac chest pain at the interface between primary and secondary care: multicentre outcome analysis of 8802 patients attending rapid access chest pain clinic, **A12**
Enhanced external counterpulsation (EECP) improves angina control and exercise tolerance in patients with chronic stable refractory angina, **A68**

- Prevalence, incidence, primary care burden, and medical treatment of angina in Scotland: age, sex, and socio-economic disparities, **A40**
- The immediate and 2-year outcomes of enhanced external counterpulsation (EECP) in the treatment of chronic refractory angina—a UK perspective, **A68**
- angiogenesis**, Genetic alteration of marrow derived stromal cells with HIF-1 α /VP16 enhances collateral forming potential in vitro and in vivo, **A1**
- Nitric oxide is not required for vasculogenesis or angiogenesis in zebrafish embryos, **A28**
- The role of angiogenic growth factors in intimal hyperplasia, **A1**
- angioplasty**, Clopidogrel preloading and myocardial necrosis after PCI, **A38**
- Impact of primary angioplasty (PPCI) programme on the management of patients presenting with ST-elevation myocardial infarction (STEMI) in a North West London borough, **A53**
- In-patient, 30-day MACE and MACE at 1–6 months in 73 consecutive bifurcation lesions treated with taxus stents and the ‘CRUSH-T Stent’ Technique, **A38**
- Stent expansion: “A failure to deliver”. A combination of delivery balloon constraint and acute stent recoil reduces predicted stent dimension by 27% irrespective of reference vessel size, **A40**
- Transradial coronary intervention without pre-screening for a dual palmar blood supply, **A39**
- Troponin T (TnT) and creatine kinase MB (CK-MB) release following percutaneous coronary intervention (PCI): are they equivalent?, **A24**
- angiotensin**, Aldosterone mediates angiotensin II induced interstitial cardiac fibrosis via a Nox2 containing NADPH oxidase, **A3**
- angiotensin ii**, Angiotensin II type 1 receptor (AT1R) blockade has no effect on the self perpetuation of AF over 4 weeks of burst pacing in a goat model of atrial fibrillation (AF), **A35**
- anti-apoptosis**, Over-expression of myocyte stress 1 protein in vitro switches on the fetal gene programme and anti-apoptotic pathways, **A44**
- anticoagulation**, Anticoagulation control and outcome in patients with second generation mechanical heart valves, **A19**
- aorta**, An operation for the Marfanoid aorta which spares the aortic valve and the vascular intima: Robiscek’s concept realised by a rapid prototyping engineering solution, **A18**
- aortic stenosis**, Severe left ventricular hypertrophy (LVH) indicates the failure of myocardial adaptation in aortic stenosis (AS), **A71**
- aortic valve disease**, Improved haemodynamic profile and left ventricular function following aortic valve replacement with a stentless rather than stented bioprosthesis: a randomised controlled trial, **A18**
- Non-invasive imaging of the coronary arteries by computed tomography (CT) for excluding significant coronary stenoses in patients requiring aortic valve replacement, **A47**
- aortic valve surgery**, Patient prosthesis mismatch in aortic valve replacement—size does not matter, **A19**
- aortic valves**, A randomised comparison of stentless versus stented valves for aortic stenosis, **A18**
- apolipoprotein B**, Homocysteine predicts cardiovascular risk independently of the metabolic syndrome, **A42**
- arrhythmia**, Cardiovascular effects of exposure to air pollutants—human exposure studies demonstrating an adverse influence on heart rate variability (HRV), **A67**
- Catheter location trial: a prospective randomised comparison of carto, NavX and fluoroscopy for catheter ablation, **A37**
- Latest results of endocardial arrhythmia ablation following the modified Fontan, **A34**
- arrhythmia clinic**, Cardiac physiologist led arrhythmia clinic substantially reduces Consultant clinic workload, **A56**
- arrhythmia surgery**, Left atrial radiofrequency ablation during mitral valve surgery for continuous atrial fibrillation: results of a prospective randomised clinical trial, **A1**
- arterial stiffness**, Increased arterial stiffness in healthy young South Asian men, **A66**
- aspirin**, Thrombogenicity of flowing blood is increased in patients with type 2 diabetes mellitus (T2DM) despite treatment with aspirin, **A61**
- assessment**, Rapid, safe, and effective management of acute chest pain in a dedicated chest pain assessment unit at St James’s Hospital, Dublin, **A3**
- atherosclerosis**, A membrane-bound vaccinia viral protein ‘Mem35K’ blocks CC-chemokine induced cell migration and CC-chemokine activity in vitro, **A30**
- Association between common polymorphisms of the pro-opiomelanocortin (POMC) gene and body fat distribution: a family study, **A41**
- Association between the -174G/C polymorphism of the Interleukin-6 gene and carotid atherosclerosis: family study and meta-analysis, **A15**
- Gene transfer of a broad spectrum CC-chemokine inhibitor reduces macrophage recruitment and smooth muscle content in vein graft atherosclerosis in apolipoprotein E-knockout mice, **A13**
- Inflammatory genotypes predisposing to premature coronary artery disease (CAD) in a large discordant sibship collection and utility as risk predictors, **A67**
- Inhibitory Smads regulate pathological cellular differentiation in atherosclerosis, **A29**
- The role of angiogenic growth factors in intimal hyperplasia, **A1**
- atrial**, A novel rectangular biphasic waveform from a radiofrequency defibrillator compared with a conventional waveform for the transvenous cardioversion of chronic atrial fibrillation in patients, **A9**
- Audit: management of patients with heart failure in a district general hospital, **A12**
- Complications of diagnostic cardiac catheterisation (CC): results from a confidential enquiry into CC complications, **A62**
- How long do patients wait with acute coronary syndromes for inter-hospital transfer in the UK? Results from the coronary heart disease collaborative national audit 2004, **A25**
- Using a coronary heart disease (CHD) register to support evidence-based cardiac care in general practice: lessons from the Scottish national CHD demonstration project, **A61**
- atrial fibrillation**, A myocardial nox2 containing NAD(P)H oxidase contributes to oxidative stress in human atrial fibrillation, **A44**
- Ablation of epicardial autonomic ganglionated plexi during minimally invasive surgical ablation of atrial fibrillation, **A9**
- Angiotensin II type 1 receptor (AT1R) blockade has no effect on the self perpetuation of AF over 4 weeks of burst pacing in a goat model of atrial fibrillation (AF), **A35**
- Association between serum thyrotrophin concentrations and occurrence of atrial fibrillation in a large community-based population of elderly subjects (The Birmingham Elderly Thyroid Study), **A40**
- Atrial fibrillation-induced Connexin 43 redistribution in atria of the Goat, **A43**
- Can atrial fibrillation with a coarse ECG appearance be treated by catheter ablation of the tricuspid valve-inferior vena cava isthmus? Preliminary results of a multicentre randomised controlled trial, **A9**
- Catheter ablation of atrial fibrillation: primary and long term efficacy for three approaches in a single centre, **A10**
- Characterisation of the electrical wavefronts seen in human permanent AF, **A37**
- Determinants of early recurrence of AF in pacemaker AF suppression study patients, **A9**
- Does increased area of pulmonary vein isolation decrease the number of non pulmonary vein initiators of atrial fibrillation?, **A36**
- Does pre-operative atrial cell electrophysiology predict post-operative atrial fibrillation?, **A43**
- Four years experience of a nurse-led elective external cardioversion service within a district general hospital setting, **A56**
- High frequency re-entry circuits driving AF: evidence from global left atrial mapping in humans, **A2**
- Interactions between focal triggers and left atrial substrate in the initiation and maintenance of atrial fibrillation: a study using dominant frequency analysis of non-contact mapping data, **A36**
- Is a pericardial window more effective than conservative treatment in reducing the incidence of post-operative atrial fibrillation (AF) following coronary artery bypass grafting (CABG)?, **A50**
- Left atrial radiofrequency ablation during mitral valve surgery for continuous atrial fibrillation: results of a prospective randomised clinical trial, **A1**
- Rapid pulmonary vein (PV) electrical isolation using a novel focused ultrasound balloon catheter in patients with atrial fibrillation (AF), **A55**
- Real-time integrated X-ray and magnetic resonance imaging (XMR)-A novel strategy to optimise precise catheter placement during pulmonary vein isolation, **A37**
- Surface atrial frequency analysis in patients with atrial fibrillation: assessing the effect of two standard linear ablation lesions sited in the left atrium, **A10**

- The prevalence/incidence paradox of atrial fibrillation in heart failure, **A57**
- The safety of DC cardioversion under sedation: long-term experience, **A36**
- atrial flutter**, Four years experience of a nurse-led elective external cardioversion service within a district general hospital setting, **A56**
- atrial remodeling**, Angiotensin II type 1 receptor (AT1R) blockade has no effect on the self perpetuation of AF over 4 weeks of burst pacing in a goat model of atrial fibrillation (AF), **A35**
- augmentation index**, Increased arterial stiffness in healthy young South Asian men, **A66**
- autonomic**, Ablation of epicardial autonomic ganglionated plexi during minimally invasive surgical ablation of atrial fibrillation, **A9**
- Changes in heart rate variability measures due to cardiac rehabilitation, **A11**
- autonomic dysfunction**, Tetrahydrobiopterin regulates vagal control of heart rate, **A57**
- AV and VV optimisation**, Optimisation of atrioventricular (AV) and interventricular (VV) delay in biventricular pacing using continuous non-invasive measurement of blood pressure (BP), **A52**
- balloon occlusion**, Evidence for the presence of a second window of preconditioning in humans during serial balloon occlusion at angioplasty on consecutive days, **A40**
- bare metal stent**, Low incidence of revascularisation of bare metal stents in the era of drug eluting stents: a single tertiary centre experience, **A5**
- Barlow's disease**, Edge-to-edge repair for Barlow's disease: early and late clinical and echocardiographic results, **A49**
- beta-blockade**, Peri-operative beta-blockade for patients undergoing infra-renal vascular surgery (POBBLE): results of a randomised double blind controlled trial, **A69**
- Beta-blockers inhibit transgene expression under transcriptional regulation of the major immediate-early cytomegalovirus enhancer/promoter in vivo, **A29**
- bifurcation lesions**, In-patient, 30-day MACE and MACE at 1–6 months in 73 consecutive bifurcation lesions treated with taxus stents and the 'CRUSH-T Stent' Technique, **A38**
- Percutaneous coronary intervention with drug eluting stents for bifurcation lesions: 6 month results, **A6**
- biventricular pacing**, Optimisation of atrioventricular (AV) and interventricular (VV) delay in biventricular pacing using continuous non-invasive measurement of blood pressure (BP), **A52**
- blood pressure**, Optimisation of atrioventricular (AV) and interventricular (VV) delay in biventricular pacing using continuous non-invasive measurement of blood pressure (BP), **A52**
- BNP**, BNP detects silent ischaemia on exercise testing in type II diabetic patients, **A45**
- Effects of amiloride and spironolactone on QT in stroke survivors—a randomised placebo controlled double-blind cross over study, **A65**
- bone marrow stromal cell**, Inhibitory Smads regulate pathological cellular differentiation in atherosclerosis, **A29**
- bosentan**, The effect of Bosentan on exercise capacity in patients with Eisenmenger's syndrome, **A35**
- bradykinin**, Effects of bradykinin (BK) on venous capacitance in health and treated chronic heart failure (CHF), **A65**
- brain natriuretic peptide**, Interrelation between NT-pro BNP, QRS width, and severity of heart failure: EuroHeart Failure Survey, **A6**
- Prognostic value of brain natriuretic peptide (BNP) in patients with pulmonary arterial hypertension (PAH), **A58**
- CABG**, Is a pericardial window more effective than conservative treatment in reducing the incidence of post-operative atrial fibrillation (AF) following coronary artery bypass grafting (CABG)?, **A50**
- What is the appropriate ratio between coronary artery bypass graft (CABG) and percutaneous intervention (PCI) in the drug eluting stent era?, **A63**
- calcification**, Calcific crystals directly induce the release of osteopontin from human macrophages via pi3-kinase, ERK1/2 map kinase, and NFκB signalling pathways: implications for atherogenesis and calcific vascular disease, **A14**
- calcium**, Cardiac nNOS regulates myocardial relaxation by stimulating phospholamban phosphorylation, **A3**
- Testosterone inhibits extracellular calcium entry via voltage-gated calcium channels in A7r5 vascular smooth muscle cells at physiological concentrations, **A43**
- calcium flux**, Molecular basis for platelet hyper-reactivity to adenosine diphosphate (ADP) associated with the platelet P2Y1 receptor 1622 G allele, **A60**
- cardiac**, Demographic and temporal trends in out-of-hospital sudden cardiac death in Belfast, **A63**
- cardiac arrest**, False cardiac arrest (FCA) calls—the right time to turn away?, **A63**
- cardiac catheterisation**, Complications of diagnostic cardiac catheterisation (CC): results from a confidential enquiry into CC complications, **A62**
- cardiac index**, The cardiovascular effects of intravenous corticotrophin releasing hormone (CRH) in healthy volunteers, **A66**
- cardiac mortality**, A rapid access cardiology service effectively diagnoses non-cardiac disease and low risk among patients presenting with chest pain, breathlessness, and palpitations, **A13**
- cardiac MRI**, Electrocardiography and cardiac magnetic resonance imaging (MRI) before and after alcohol septal myocardial ablation (ASA) in hypertrophic obstructive cardiomyopathy, **A70**
- cardiac physiologists**, Cardiac physiologist led arrhythmia clinic substantially reduces Consultant clinic workload, **A56**
- cardiac rehabilitation**, Home-based versus supervised centre-based cardiac rehabilitation in a multi-ethnic population: 6 month results of the Birmingham Rehabilitation Uptake Maximisation Study (BRUM), **A53**
- Physical activity behaviour in users of a menu based cardiac rehabilitation programme: a longitudinal survey, **A11**
- Psychological rehabilitation for cardiac patients: systematic review and meta-analysis, **A11**
- The growth of cardiac rehabilitation in the UK since 1998, **A10**
- cardiac resynchronisation**, Clinical benefit from cardiac resynchronisation therapy (CRT) depends on myocardial viability rather than on QRS duration, **A52**
- Increased QT dispersion as a predictor of mortality following biventricular pacing, **A54**
- cardiac risk**, Peri-operative beta-blockade for patients undergoing infra-renal vascular surgery (POBBLE): results of a randomised double blind controlled trial, **A69**
- cardiac transplantation**, Calcineurin inhibitor (CI) withdrawal and replacement with sirolimus (SIR) in CTR with renal dysfunction, **A49**
- Donor hearts with coronary artery disease – an under utilised resource in cardiac transplantation?, **A49**
- cardiology specialist registrars**, Elective coronary angiography by non-medical angiographers. A comparison with cardiology specialist registrars, **A68**
- cardiomyocyte**, FoxO transcription factor expressions and activity are regulated during the development of cardiomyocyte hypertrophy, **A44**
- The tumour suppressor RASSF1A attenuates cardiomyocyte hypertrophy in response to phenylephrine, **A44**
- cardiomyopathy**, Disease severity and abnormal collagen turn over in anderson fabry disease (AFD), **A70**
- Isolated left ventricular noncompaction: two year experience with cardiovascular magnetic resonance, **A17**
- Left ventricular non-compaction (LVNC): insights from cardiovascular magnetic resonance (CMR) imaging, **A71**
- Novel mutations in cardiac MYBPC3 cause early onset malignant hypertrophic cardiomyopathy (HCM), **A69**
- cardiovascular events**, Prognostic value of coronary artery calcium screening in asymptomatic patients with type 2 diabetes, **A16**
- cardioversion**, Four years experience of a nurse-led elective external cardioversion service within a district general hospital setting, **A56**
- The safety of DC cardioversion under sedation: long-term experience, **A36**
- cellular electrophysiology**, Does pre-operative atrial cell electrophysiology predict post-operative atrial fibrillation?, **A43**
- chemokines**, A membrane-bound vaccinia viral protein 'Mem35K' blocks CC-chemokine induced cell migration and CC-chemokine activity in vitro, **A30**
- chest pain**, Is it safe to discharge patients from accident and emergency using a rapid point of care triple cardiac marker test to rule out acute coronary syndrome in patients presenting with chest pain?, **A26**
- Rapid, safe, and effective management of acute chest pain in a dedicated chest pain assessment unit at St James's Hospital, Dublin, **A3**
- circadian**, Circadian clock genes cause activation of the human PAI-1 gene promoter with 4G/5G allelic preference, **A51**

- circulating endothelial cells**, Circulating endothelial cells, von Willebrand factor, and flow mediated dilatation as indices of endothelial damage/dysfunction in patients with acute coronary syndromes, **A14**
Increased numbers of circulating endothelial cells predict adverse cardiovascular events following an acute coronary syndrome, **A27**
- clinical outcome**, Patients with non-ST elevation acute coronary syndromes associated with myonecrosis have a similar rate of death or MI as ST elevation MI at 6 months: an audit of all ACS admissions in a UK district general hospital over 1 year, **A42**
- clopidogrel**, Clopidogrel preloading and myocardial necrosis after PCI, **A38**
Duration of clopidogrel therapy following percutaneous coronary intervention (PCI) with bare metal (BMS) and drug eluting stents (DES). A questionnaire survey in the UK; do we need to be NICE?, **A64**
- CMR**, Left ventricular remodelling in dilated cardiomyopathy (DCM) with and without evidence of late gadolinium enhancement-cardiovascular magnetic resonance (LGE-CMR), **A71**
- collagen**, Disease severity and abnormal collagen turn over in anderson fabry disease (AFD), **A70**
- complication rates**, Introduction of a risk scoring system for percutaneous coronary intervention (PCI) procedures, **A63**
- complications**, Complications of diagnostic cardiac catheterisation (CC): results from a confidential enquiry into CC complications, **A62**
Transradial coronary intervention without pre-screening for a dual palmar blood supply, **A39**
- conduction**, The influence of atrial conduction, refractoriness, and volume on outcomes following pulmonary vein isolation (PVI), **A54**
- congenital**, Is exercise training beneficial in adults with congenital heart disease?, **A33**
- congenital heart defects**, Identification of cardiac abnormalities in the Sonic Hedgehog mouse mutant using optical projection tomography, **A34**
Role of the planar cell polarity pathway in the development of the outflow tract, **A34**
- connexin**, Atrial fibrillation-induced Connexin 43 redistribution in atria of the Goat, **A43**
- consultants**, Impact of a 24-hour primary PCI service for ST elevation myocardial infarction (STEMI) on consultants' work patterns, **A27**
- contractility**, P38-MAPK mediates the early negative inotropic effect of tumour necrosis factor- α (TNF). Evidence of synergy between a direct negative inotropic effect and coronary constriction, **A57**
- contrast echocardiography**, Improved accuracy of low power contrast echocardiography for the assessment of left ventricular remodeling compared with unenhanced harmonic echocardiography following acute myocardial infarction: comparison with cardiac magnetic resonance imaging, **A32**
Prognostic value of myocardial contrast echocardiography in patients presenting to hospital with acute chest pain and negative troponin, **A31**
- control**, Anticoagulation control and outcome in patients with second generation mechanical heart valves, **A19**
Changes in heart rate variability measures due to cardiac rehabilitation, **A11**
- coronary angiography**, Diagnostic accuracy of non-invasive coronary angiography with multislice computed tomography in patients presenting with acute chest pain, **A15**
Donor hearts with coronary artery disease – an under utilised resource in cardiac transplantation?, **A49**
Elective coronary angiography by non-medical angiographers. A comparison with cardiology specialist registrars, **A68**
Non-invasive imaging of the coronary arteries by computed tomography (CT) for excluding significant coronary stenoses in patients requiring aortic valve replacement, **A47**
Radiation dose in coronary multislice CT: a comparison with conventional diagnostic angiography, **A16**
- coronary artery**, Effect of nitrate administration on wave speed propagation in the human coronary artery, **A66**
- coronary artery bypass grafting**, Relationship between mild renal dysfunction and outcomes after coronary artery bypass grafting, **A19**
The adipocyte derived peptide hormone leptin is an endothelial independent vasodilator in humans: implications for vascular homeostasis, **A28**
- coronary artery disease**, Donor hearts with coronary artery disease – an under utilised resource in cardiac transplantation?, **A49**
Enhanced external counterpulsation (EECP) improves angina control and exercise tolerance in patients with chronic stable refractory angina, **A68**
Erythropoietin delays endothelial cell senescence by preserving telomere length, **A14**
Prognostic value of myocardial contrast echocardiography in patients presenting to hospital with acute chest pain and negative troponin, **A31**
The Genetic Risk of Acute Coronary Events (GRACE) Study cohort - demography and description of a large UK discordant-sibling collection, **A41**
The immediate and 2-year outcomes of enhanced external counterpulsation (EECP) in the treatment of chronic refractory angina—a UK perspective, **A68**
- coronary calcium**, Plasma osteoprotegerin levels predict the extent of subclinical coronary atherosclerosis and adverse cardiac events in asymptomatic type 2 diabetic subjects, **A17**
Prognostic value of coronary artery calcium screening in asymptomatic patients with type 2 diabetes, **A16**
- coronary disease**, Relationship between mild renal dysfunction and outcomes after coronary artery bypass grafting, **A19**
- coronary flow reserve**, Altered coronary microvascular function and peripheral endothelial dysfunction in young patients with systemic lupus erythematosus, **A28**
- coronary intervention**, Clinical presentation with in-stent restenosis is not always benign, **A39**
Introduction of a risk scoring system for percutaneous coronary intervention (PCI) procedures, **A63**
- corticotrophin releasing hormone**, The cardiovascular effects of intravenous corticotrophin releasing hormone (CRH) in healthy volunteers, **A66**
- cost effectiveness**, UK Primary Angioplasty Cost Effectiveness Study (UK-PACES) 30 day outcome data, **A27**
- creatine kinase**, Troponin T (TnT) and Creatine Kinase MB (CK-MB) release following Percutaneous Coronary Intervention (PCI): Are they equivalent?, **A24**
- Cryocath**, Experience with 75 consecutive cryoablation procedures in a single UK tertiary centre, **A10**
- CT**, Non-invasive imaging of the coronary arteries by computed tomography (CT) for excluding significant coronary stenoses in patients requiring aortic valve replacement, **A47**
- cytokine**, Association between the -174G/C polymorphism of the Interleukin-6 gene and carotid atherosclerosis: family study and meta-analysis, **A15**
P38-MAPK mediates the early negative inotropic effect of tumour necrosis factor- α (TNF). Evidence of synergy between a direct negative inotropic effect and coronary constriction, **A57**
- death**, Demographic and temporal trends in out-of-hospital sudden cardiac death in Belfast, **A63**
- defibrillation**, A novel rectangular biphasic waveform from a radiofrequency defibrillator compared with a conventional waveform for the transvenous cardioversion of chronic atrial fibrillation in patients, **A9**
Dofetilide significantly reduces acute and chronic defibrillation thresholds in patients with an inadequate defibrillation safety margin, **A35**
- defibrillator**, Safety and acceptability of conscious sedation for subpectoral implantation of implantable cardioverter defibrillators (ICDs), **A54**
- delayed preconditioning**, Evidence for the presence of a second window of preconditioning in humans during serial balloon occlusion at angioplasty on consecutive days, **A40**
- diabetes**, Fasting plasma glucose fails to detect new diabetes or abnormal glucose tolerance (GI) in patients admitted with acute myocardial infarction (MI), **A47**
Inflammation and acute hyperglycaemia in diabetics with non-ST elevation acute coronary syndromes (ACS). Analyses from OPUS TIMI 16 and TACTICS TIMI 18, **A60**
Real life impact of sirolimus eluting stent implantation in treating diabetic patients: a 6 month follow up report from the international e-CYPHER registry, **A5**
Reversal of vascular dysfunction in subjects at risk of type 2 diabetes mellitus (DM), **A53**

- diabetes mellitus**, Elevated serum glucose predicts mortality in patients admitted to hospital with a new diagnosis of heart failure, **A7**
Is clinically overt diabetes mellitus a cause or a consequence of heart failure?, **A58**
Thrombogenicity of flowing blood is increased in patients with type 2 diabetes mellitus (T2DM) despite treatment with aspirin, **A61**
- diabetic**, BNP detects silent ischaemia on exercise testing in type II diabetic patients, **A45**
- diagnosis**, Screening for suspected heart failure with N terminal pro-B type natriuretic peptide (NT proBNP) in primary care: money well spent?, **A56**
Suspected heart failure in primary care - the utility of n-terminal pro b-type natriuretic peptide (NT PROBNP) as a pre-screening test for secondary care referral: a real life study, **A22**
Value of heart fatty acid binding protein (hFABP) for very early detection of acute myocardial infarction (AMI), **A60**
- diastolic heart failure**, An analysis of the apparent frequency of diastolic heart failure in a secondary care setting, using different published definitions, **A22**
- dilated cardiomyopathy**, Left ventricular remodelling in dilated cardiomyopathy (DCM) with and without evidence of late gadolinium enhancement-cardiovascular magnetic resonance (LGE-CMR), **A71**
- dissection**, Presentation and outcome of Marfan's syndrome patients with type B dissection and thoracoabdominal aortic aneurysm, **A19**
- district hospital**, Audit: management of patients with heart failure in a district general hospital, **A12**
Two years of primary coronary intervention for ST elevation myocardial infarction in a district general hospital, **A59**
- dobutamine stress**, Stress Induced QT prolongation does not predict ischaemia in patients with stable Coronary Artery Disease, **A46**
- dobutamine stress echo**, Dobutamine stress echocardiography (DSE) and the resting but not exercise electrocardiograph (ECG) predict severe coronary artery disease (CAD) in end stage renal failure, **A68**
- dofetilide**, Dofetilide significantly reduces acute and chronic defibrillation thresholds in patients with an inadequate defibrillation safety margin, **A35**
- doppler echocardiography**, Cardiac dyssynchrony as measured by total isovolumic time may predispose to central sleep apnoea in chronic heart failure, **A21**
- drug eluting stent**, Low incidence of revascularisation of bare metal stents in the era of drug eluting stents: a single tertiary centre experience, **A5**
Multivessel PCI procedures with implantation of Sirolimus eluting stents: a report from the e-CYPHER registry, **A6**
Percutaneous coronary intervention with drug eluting stents for bifurcation lesions: 6 month results, **A6**
Real life impact of sirolimus eluting stent implantation in treating diabetic patients: a 6 month follow up report from the international e-CYPHER registry, **A5**
Real world use of sirolimus eluting stents in saphenous vein graft disease: data from the e-CYPHER registry, **A5**
Sirolimus eluting stent and unprotected left main stenosis: the multicenter e-CYPHER registry, **A5**
Comparison of outcomes in real world use of Sirolimus eluting stents with the randomised controlled clinical trials: can the results be reproduced?, **A6**
Duration of clopidogrel therapy following percutaneous coronary intervention (PCI) with bare metal (BMS) and drug eluting stents (DES). A questionnaire survey in the UK; do we need to be NICE?, **A64**
- drug eluting stents**, In-patient, 30-day MACE and MACE at 1–6 months in 73 consecutive bifurcation lesions treated with taxus stents and the 'CRUSH-T Stent' Technique, **A38**
Retrospective study to investigate the use of Drug Eluting Stents (DES) for the treatment of In-Stent Restenosis - 'The Real world' experience, **A38**
What is the appropriate ratio between coronary artery bypass graft (CABG) and percutaneous intervention (PCI) in the drug eluting stent era?, **A63**
- dystrophin**, Role of the PMCA-Syntrophin-nNOS complex in cardiomyopathy, **A45**
- early invasive management**, Use of regional guidelines and rapid access to cardiac catheter laboratories facilitate early invasive management in patients with non-ST segment elevation acute coronary syndromes (NSTEMI/ACS), **A62**
- echocardiography**, 3D contraction front mapping: novel, high resolution imaging of mechanical contraction of the left ventricle, **A30**
A randomised comparison of stentless versus stented valves for aortic stenosis, **A18**
An analysis of the apparent frequency of diastolic heart failure in a secondary care setting, using different published definitions, **A22**
Can real-time 3D echocardiography predict reverse left ventricular remodelling post cardiac resynchronisation therapy?, **A31**
Can reperfusion be predicted by acute thickening of the at risk myocardium in STEMI? Full pressure v pressure limiting reperfusion, **A32**
The impact of the introduction of NT-proBNP into primary care on secondary care referral rates, **A57**
- edge to edge repair**, Edge-to-edge repair for Barlow's disease: early and late clinical and echocardiographic results, **A49**
- efficacy**, Catheter ablation of atrial fibrillation: primary and long term efficacy for three approaches in a single centre, **A10**
- effusion**, Variability in pericardiocentesis practice among cardiologists in the UK—need for guidelines, **A72**
- Eisenmenger's syndrome**, The effect of Bosentan on exercise capacity in patients with Eisenmenger's syndrome, **A35**
- elderly**, Anti-anginal and anti-ischaemic efficacy of ivabradine—a selective and specific sinus node If current inhibitor—compared to atenolol in elderly patients with chronic stable angina, **A69**
Association between serum thyrotrophin concentrations and occurrence of atrial fibrillation in a large community-based population of elderly subjects (The Birmingham Elderly Thyroid Study), **A40**
Clinical benefits of low LDL targets in elderly patients with coronary disease: An analysis from PROVE IT-TIMI-22, **A46**
- electrocardiogram**, Dobutamine stress echocardiography (DSE) and the resting but not exercise electrocardiograph (ECG) predict severe coronary artery disease (CAD) in end stage renal failure, **A68**
- electrophysiology**, Catheter location trial: a prospective randomised comparison of carto, NavX and fluoroscopy for catheter ablation, **A37**
- endogenous fibrinolysis**, Combustion derived nanoparticulate impairs vascular function and endogenous fibrinolysis in man: an explanation for the increased cardiovascular mortality associated with air pollution, **A2**
- endothelial damage**, Increased numbers of circulating endothelial cells predict adverse cardiovascular events following an acute coronary syndrome, **A27**
- endothelial dysfunction**, Altered coronary microvascular function and peripheral endothelial dysfunction in young patients with systemic lupus erythematosus, **A28**
Reversal of vascular dysfunction in subjects at risk of type 2 diabetes mellitus (DM), **A53**
- endothelial function**, Combustion derived nanoparticulate impairs vascular function and endogenous fibrinolysis in man: an explanation for the increased cardiovascular mortality associated with air pollution, **A2**
- endothelin**, The effect of Bosentan on exercise capacity in patients with Eisenmenger's syndrome, **A35**
- endothelium**, Atorvastatin improves endothelial function and decreases the expression of proinflammatory cytokines and adhesion molecules in patients with heart failure, **A20**
Insulin resistance is a substrate for accelerated endothelial dysfunction in middle age—studies in mice heterozygous for knockout of the insulin receptor gene (IRKO), **A28**
Investigation of P2X7 dependent mechanisms of Interleukin-1 beta (IL-1 β) release from human endothelial cells, **A13**
NOX-2 (gp91phox) disruption prevents endothelial ischaemia-reperfusion injury in humans, **A27**
Prevention of hypoxia-induced pulmonary hypertension and vascular remodelling by targeting endothelial tetrahydrobiopterin-dependent nitric oxide synthase regulation, **A4**
- enhanced external counterpulsation**, Enhanced external counterpulsation (EECP) improves angina control and exercise tolerance in patients with chronic stable refractory angina, **A68**
The immediate and 2-year outcomes of enhanced external counterpulsation (EECP) in the treatment of chronic refractory angina—a UK perspective, **A68**
- eNOS**, Endothelial tetrahydrobiopterin (BH4) regulates eNOS coupling in vivo: functional effects of BH4-eNOS stoichiometry in endothelial targeted transgenic mice, **A50**

- epicardial**, Percutaneous transthoracic epicardial ablation (PTEA) for failed endocardial ablations: promising results of a new technique, **A54**
- epidemiology**, The hospital burden of suspected acute coronary syndromes (ACS): recent trends, **A60**
The prevalence/incidence paradox of atrial fibrillation in heart failure, **A57**
- ERAF**, Determinants of early recurrence of AF in pacemaker AF suppression study patients, **A9**
- erythrocytes**, Cholesterol content of erythrocyte membranes may contribute to the development of acute coronary syndrome. A new paradigm regarding coronary artery disease instability?, **A45**
- ethnicity**, Declining case fatality rates for acute myocardial infarction in south Asians and whites in the last 15 years, **A41**
Ethnic differences in the prevalence and aetiology of left ventricular systolic dysfunction (LVD) in high-risk community subjects, **A58**
Ethnicity and variation in prognosis for patients newly hospitalised for heart failure, **A8**
- Euro Heart Failure Survey**, Predictors of short term mortality in heart failure: EuroHeart Failure Survey, **A7**
- evidence-based care**, Using a coronary heart disease (CHD) register to support evidence-based cardiac care in general practice: lessons from the Scottish national CHD demonstration project, **A61**
- exercise**, Is exercise training beneficial in adults with congenital heart disease?, **A33**
- exercise capacity**, Plasma N-terminal pro-brain natriuretic peptide predicts exercise capacity in degenerative mitral valve disease, **A49**
- exercise testing**, Myocardial perfusion scintigraphy predicts functional outcome following percutaneous coronary intervention, **A17**
Pregnant cardiac patients: development of a generic test to evaluate the functional adequacy of diseased hearts to cope with pregnancy and labour, **A33**
- exercise tolerance test**, BNP detects silent ischaemia on exercise testing in type II diabetic patients, **A45**
- extrapulmonary sites**, Does increased area of pulmonary vein isolation decrease the number of non pulmonary initiators of atrial fibrillation?, **A36**
- fabry disease**, Disease severity and abnormal collagen turn over in anderson fabry disease (AFD), **A70**
- failed lysis**, The REACT (REscue Angioplasty v Conservative treatment or repeat Thrombolysis) Trial: impact of age upon efficacy of treatment, **A24**
The REACT (REscue Angioplasty v Conservative treatment or repeat Thrombolysis) Trial: results according to actual treatment received, **A22**
- failed thrombolysis**, Factors influencing left ventricular function following successful rescue percutaneous coronary intervention for failed thrombolysis in acute myocardial infarction, **A24**
- false alert**, False cardiac arrest (FCA) calls—the right time to turn away?, **A63**
- fibrinolysis**, Circadian clock genes cause activation of the human PAI-1 gene promoter with 4G/5G allelic preference, **A51**
- fibrosis**, Aldosterone mediates angiotensin II induced interstitial cardiac fibrosis via a Nox2 containing NADPH oxidase, **A3**
- flow mediated dilatation**, Circulating endothelial cells, von Willebrand factor, and flow mediated dilatation as indices of endothelial damage/dysfunction in patients with acute coronary syndromes, **A14**
- follow up**, Mortality of patients in England with left ventricular systolic dysfunction and heart failure due to other causes, **A8**
- fontan**, Latest results of endocardial arrhythmia ablation following the modified Fontan, **A34**
- fontan circulation**, Quadriceps and respiratory muscle strength in patients with a Fontan circulation, **A34**
- frequency analysis**, Surface atrial frequency analysis in patients with atrial fibrillation: assessing the effect of two standard linear ablation lesions sited in the left atrium, **A10**
- functional capacity**, Testosterone therapy in men with heart failure: a double blind placebo controlled trial, **A20**
- future directions**, A UK centre experience of the efficacy of implantable cardiac defibrillators (ICDs) for the prevention of sudden cardiac death in patients with hypertrophic cardiomyopathy (HCM), **A70**
- gap junction**, Atrial fibrillation-induced Connexin 43 redistribution in atria of the Goat, **A43**
- Gd-CMR**, Clinical benefit from cardiac resynchronisation therapy (CRT) depends on myocardial viability rather than on QRS duration, **A52**
- gene expression**, Over-expression of myocyte stress 1 protein in vitro switches on the fetal gene programme and anti-apoptotic pathways, **A44**
- gene therapy**, Beta-blockers inhibit transgene expression under transcriptional regulation of the major immediate-early cytomegalovirus enhancer/promoter in vivo, **A29**
Genetic alteration of marrow derived stromal cells with HIF-1 α /VP16 enhances collateral forming potential in vitro and in vivo, **A1**
Intracoronary delivery of a secreted transforming growth factor- β type II receptor from a novel collagen-membrane bound stent inhibits in-stent neointimal hyperplasia, **A50**
- genetics**, A rare variant of the leptin (LEP) gene has large effects on blood pressure (BP) and carotid intima-medial thickness (CIMT): study of 1428 individuals in 248 families, **A64**
Association between common polymorphisms of the pro-opiomelanocortin (POMC) gene and body fat distribution: a family study, **A41**
Association between the -174G/C polymorphism of the Interleukin-6 gene and carotid atherosclerosis: family study and meta-analysis, **A15**
Genetic variation at the locus encompassing 11-beta hydroxylase and aldosterone synthase accounts for heritability in cortisol precursor (11-deoxycortisol) urinary metabolite excretion, **A65**
Identification of cardiac abnormalities in the Sonic Hedgehog mouse mutant using optical projection tomography, **A34**
Role of the planar cell polarity pathway in the development of the outflow tract, **A34**
The Genetic Risk of Acute Coronary Events (GRACE) Study cohort - demography and description of a large UK discordant-sibling collection, **A41**
- glucose tolerance test**, Fasting plasma glucose fails to detect new diabetes or abnormal glucose tolerance (GI) in patients admitted with acute myocardial infarction (MI), **A47**
- growth factors**, Erythropoietin delays endothelial cell senescence by preserving telomere length, **A14**
- H-FABP**, H-FABP: a strong predictor of mortality after 'new' definition myocardial infarction, **A46**
- haemoglobin**, Haemoglobin is related to prognostic factors but is not an independent survival predictor in patients referred for cardiac transplantation, **A48**
Indicators of increased risk of mortality following first hospital admission with heart failure: the value of routine clinical variables, **A8**
- halo mapping**, Ablation of difficult right sided accessory pathways (AP) aided by simple mapping of the tricuspid VALVE annulus (TVA) using a halo catheter, **A55**
- heart**, The true incidence of primary graft dysfunction leading to death in heart transplantation: a validated audit in 1820 patients, **A48**
- heart disease**, Pregnant cardiac patients: development of a generic test to evaluate the functional adequacy of diseased hearts to cope with pregnancy and labour, **A33**
- heart failure**, Actions of aldosterone blockade in patients with mild-moderately severe heart failure, **A20**
Anaemia predisposes to heart failure in acute coronary syndromes independently of the severity of myocardial injury, **A7**
Atorvastatin improves endothelial function and decreases the expression of proinflammatory cytokines and adhesion molecules in patients with heart failure, **A20**
Audit: management of patients with heart failure in a district general hospital, **A12**
Cardiac dyssynchrony as measured by total isovolumic time may predispose to central sleep apnoea in chronic heart failure, **A21**
Clinical benefit from cardiac resynchronisation therapy (CRT) depends on myocardial viability rather than on QRS duration, **A52**
Clinical predictors of 6-minute walk test (6-MWT) performance in patients with chronic heart failure (CHF), **A58**
Effects of bradykinin (BK) on venous capacitance in health and treated chronic heart failure (CHF), **A65**
Effects of optimal pharmacological treatment on left ventricular remodelling, systolic and diastolic function in patients with chronic heart failure, **A20**
Effects of physical training on anxiety and depression in chronic heart failure, **A12**

- Ethnic differences in the prevalence and aetiology of left ventricular systolic dysfunction (LVD) in high-risk community subjects, **A58**
- Increased QT dispersion as a predictor of mortality following biventricular pacing, **A54**
- Interrelation between NT-pro BNP, QRS width, and severity of heart failure: EuroHeart Failure Survey, **A6**
- Is clinically overt diabetes mellitus a cause or a consequence of heart failure?, **A58**
- Mortality of patients in England with left ventricular systolic dysfunction and heart failure due to other causes, **A8**
- Predictors of short term mortality in heart failure: EuroHeart Failure Survey, **A7**
- Role of real-time transthoracic 3D echocardiography in the assessment of mechanical asynchrony, **A31**
- Screening for suspected heart failure with N terminal pro-B type natriuretic peptide (NT proBNP) in primary care: money well spent?, **A56**
- Suspected heart failure in primary care - the utility of n-terminal pro b-type natriuretic peptide (NT PROBNP) as a pre-screening test for secondary care referral: a real life study, **A22**
- Testosterone therapy in men with heart failure: a double blind placebo controlled trial, **A20**
- The prevalence/incidence paradox of atrial fibrillation in heart failure, **A57**
- heart rate variability**, Cardiovascular effects of exposure to air pollutants—human exposure studies demonstrating an adverse influence on heart rate variability (HRV), **A67**
- Changes in heart rate variability measures due to cardiac rehabilitation, **A11**
- The cardiovascular effects of intravenous corticotrophin releasing hormone (CRH) in healthy volunteers, **A66**
- heart transplantation**, Ultrasonic strain and strain rate imaging: a new non-invasive technique to identify acute rejection after heart transplantation, **A47**
- home based**, Randomised controlled trial comparing the clinical effectiveness of home based and hospital based rehabilitation after acute myocardial infarction: the Cornwall Heart Attack Rehabilitation Management Study, **A3**
- home care**, Home-based versus supervised centre-based cardiac rehabilitation in a multi-ethnic population: 6 month results of the Birmingham Rehabilitation Uptake Maximisation Study (BRUM), **A53**
- homocysteine**, Homocysteine predicts cardiovascular risk independently of the metabolic syndrome, **A42**
- hormones**, Testosterone therapy in men with heart failure: a double blind placebo controlled trial, **A20**
- hospital based**, Randomised controlled trial comparing the clinical effectiveness of home based and hospital based rehabilitation after acute myocardial infarction: the Cornwall Heart Attack Rehabilitation Management Study, **A3**
- human**, Characterisation of the electrical wavefronts seen in human permanent AF, **A37**
- High frequency re-entry circuits driving AF: evidence from global left atrial mapping in humans, **A2**
- hyperglycaemia**, Elevated serum glucose predicts mortality in patients admitted to hospital with a new diagnosis of heart failure, **A7**
- hypertension**, A rare variant of the leptin (LEP) gene has large effects on blood pressure (BP) and carotid intima-medial thickness (CIMT): study of 1428 individuals in 248 families, **A64**
- Autonomic remodelling in hypertension following cardiac neural nitric oxide synthase (nNOS) gene transfer, **A65**
- Genetic variation at the locus encompassing 11-beta hydroxylase and aldosterone synthase accounts for heritability in cortisol precursor (11-deoxycortisol) urinary metabolite excretion, **A65**
- Protection against the development of obesity, insulin resistance, and hypertension in mice by overexpression of insulin-like growth factor binding protein-2 (IGFBP-2), **A51**
- hypertrophic cardiomyopathy**, A UK centre experience of the efficacy of implantable cardiac defibrillators (ICDs) for the prevention of sudden cardiac death in patients with hypertrophic cardiomyopathy (HCM), **A70**
- Delineating the role of adenosine monophosphate (AMP)-activated protein kinase (AMPK) γ 2 subunit in hypertrophic cardiomyopathy (HCM), **A69**
- Electrocardiography and cardiac magnetic resonance imaging (MRI) before and after alcohol septal myocardial ablation (ASA) in hypertrophic obstructive cardiomyopathy, **A70**
- hypertrophy**, A randomised comparison of stentless versus stented valves for aortic stenosis, **A18**
- FoxO transcription factor expressions and activity are regulated during the development of cardiomyocyte hypertrophy, **A44**
- Myostatin regulates cardiomyocyte growth through modulation of Akt signalling in vitro and in vivo, **A1**
- Novel mutations in cardiac MYPBC3 cause early onset malignant hypertrophic cardiomyopathy (HCM), **A69**
- Over-expression of myocyte stress 1 protein in vitro switches on the fetal gene programme and anti-apoptotic pathways, **A44**
- TGFbeta Typell receptor interacts with TGFbeta Activated Kinase (TAK1): implications for controlling cardiac hypertrophy, **A43**
- The effect of smoking and other lifestyle factors on the left ventricular growth response to exercise in 312 young healthy men assessed using cardiovascular magnetic resonance, **A21**
- The tumour suppressor RASSF1A attenuates cardiomyocyte hypertrophy in response to phenylephrine, **A44**
- hypoplastic left heart syndrome**, Risk factors for sudden death in infants having undergone the Norwood operation for hypoplastic left heart syndrome (HLHS), **A53**
- ICD/CRT/CRTD devices**, Usage of implantable defibrillators and biventricular pacemakers in cardiac transplant candidates: a survey of UK transplant centres, **A48**
- imaging**, Measurement of coronary stent expansion using StentBoost™ image enhancement software: a comparison with intravascular ultrasound, **A39**
- implantable cardiac defibrillator**, A UK centre experience of the efficacy of implantable cardiac defibrillators (ICDs) for the prevention of sudden cardiac death in patients with hypertrophic cardiomyopathy (HCM), **A70**
- in stent restenosis**, Retrospective study to investigate the use of Drug Eluting Stents (DES) for the treatment of In-Stent Restenosis – ‘The Real world’ experience, **A38**
- inducible myocardial ischaemia**, QT peak prolongation may reflect inducible myocardial ischaemia, **A47**
- inflammation**, A membrane-bound vaccinia viral protein ‘Mem35K’ blocks CC-chemokine induced cell migration and CC-chemokine activity in vitro, **A30**
- Atorvastatin improves endothelial function and decreases the expression of proinflammatory cytokines and adhesion molecules in patients with heart failure, **A20**
- Gene transfer of a broad spectrum CC-chemokine inhibitor reduces macrophage recruitment and smooth muscle content in vein graft atherosclerosis in apolipoprotein E-knockout mice, **A13**
- Inflammation and acute hyperglycaemia in diabetics with non-ST elevation acute coronary syndromes (ACS). Analyses from OPUS TIMI 16 and TACTICS TIMI 18, **A60**
- Inflammatory genotypes predisposing to premature coronary artery disease (CAD) in a large discordant sibship collection and utility as risk predictors, **A67**
- Tribbles-1, a mitogen activated protein kinase (MAPK) scaffold regulates inflammation in vascular smooth muscle cells (VSMC) and atherosclerosis, **A15**
- inotropic reserve**, Effect of dobutamine stress on left ventricular filling in ischaemic cardiomyopathy: pathophysiology and prognostic implications, **A21**
- instability**, Cholesterol content of erythrocyte membranes may contribute to the development of acute coronary syndrome. A new paradigm regarding coronary artery disease instability?, **A45**
- insulin**, Insulin resistance is a substrate for accelerated endothelial dysfunction in middle age—studies in mice heterozygous for knockout of the insulin receptor gene (IRKO), **A28**
- insulin resistance**, Protection against the development of obesity, insulin resistance, and hypertension in mice by overexpression of insulin-like growth factor binding protein-2 (IGFBP-2), **A51**
- interleukin-10**, Circulating systemic levels of interleukin-6 and interleukin-10. Do they indicate the presence of a vulnerable plaque?, **A15**
- interleukin-1**, Investigation of P2X7 dependent mechanisms of Interleukin-1 beta (IL-1 β) release from human endothelial cells, **A13**
- interleukin-6**, Circulating systemic levels of interleukin-6 and interleukin-10. Do they indicate the presence of a vulnerable plaque?, **A15**
- intracoronary nitrates**, Effect of nitrate administration on wave speed propagation in the human coronary artery, **A66**

- intraventricular dyssynchrony**, 3D contraction front mapping: novel, high resolution imaging of mechanical contraction of the left ventricle, **A30**
- ischaemia**, Stress Induced QT prolongation does not predict Ischaemia in patients with stable Coronary Artery Disease, **A46**
- ischaemia-reperfusion injury**, NOX-2 (gp91phox) disruption prevents endothelial ischaemia-reperfusion injury in humans, **A27**
- ivabradine**, Anti-anginal and anti-ischaemic efficacy of ivabradine—a selective and specific sinus node If current inhibitor—compared to atenolol in elderly patients with chronic stable angina, **A69**
- kinase**, Myostatin regulates cardiomyocyte growth through modulation of Akt signalling in vitro and in vivo, **A1**
- L-type calcium current**, Autonomic remodelling in hypertension following cardiac neural nitric oxide synthase (nNOS) gene transfer, **A65**
- latent diabetes**, Is clinically overt diabetes mellitus a cause or a consequence of heart failure?, **A58**
- LDL**, Clinical benefits of low LDL targets in elderly patients with coronary disease: An analysis from PROVE IT-TIMI-22, **A46**
- left bundle branch block**, Left bundle branch block (LBBB) myocardial infarction (MI)—a study of thrombolysis in a district general hospital, **A59**
- left main stem**, Sirolimus eluting stent and unprotected left main stenosis: the multicenter e-CYPHER registry, **A5**
- Left ventricular function**, Can real-time 3D echocardiography predict reverse left ventricular remodelling post cardiac resynchronisation therapy?, **A31**
Improved haemodynamic profile and left ventricular function following aortic valve replacement with a stentless rather than stented bioprosthesis: a randomised controlled trial, **A18**
Severe left ventricular hypertrophy (LVH) indicates the failure of myocardial adaptation in aortic stenosis (AS), **A71**
- left ventricular hypertrophy**, Severe left ventricular hypertrophy (LVH) indicates the failure of myocardial adaptation in aortic stenosis (AS), **A71**
- length of stay**, The hospital burden of suspected acute coronary syndromes (ACS): recent trends, **A60**
- leptin**, A rare variant of the leptin (LEP) gene has large effects on blood pressure (BP) and carotid intima-medial thickness (CIMT): study of 1428 individuals in 248 families, **A64**
The adipocyte derived peptide hormone leptin is an endothelial independent vasodilator in humans: implications for vascular homeostasis, **A28**
- long QT peak**, QT peak prolongation may reflect inducible myocardial ischaemia, **A47**
- LV remodelling**, Improved accuracy of low power contrast echocardiography for the assessment of left ventricular remodeling compared with unenhanced harmonic echocardiography following acute myocardial infarction: comparison with cardiac magnetic resonance imaging, **A32**
Left ventricular remodelling in dilated cardiomyopathy (DCM) with and without evidence of late gadolinium enhancement-cardiovascular magnetic resonance (LGE-CMR), **A71**
- LV systolic function**, 3D contraction front mapping: novel, high resolution imaging of mechanical contraction of the left ventricle, **A30**
An analysis of the apparent frequency of diastolic heart failure in a secondary care setting, using different published definitions, **A22**
- macrophages**, Calcific crystals directly induce the release of osteopontin from human macrophages via pi3-kinase, ERK1/2 map kinase, and NFκB signalling pathways: implications for atherogenesis and calcific vascular disease, **A14**
- magnetic resonance imaging**, Effects of optimal pharmacological treatment on left ventricular remodelling, systolic and diastolic function in patients with chronic heart failure, **A20**
Isolated left ventricular noncompaction: two year experience with cardiovascular magnetic resonance, **A17**
Left ventricular non-compaction (LVNC): insights from cardiovascular magnetic resonance (CMR) imaging, **A71**
New generation three-dimensional echocardiography for left ventricular volumetric and functional measurements: comparison with cardiac magnetic resonance, **A30**
Transmural extent of infarction on contrast enhanced magnetic resonance imaging (CE-MRI) predicts recovery of contractile function in patients with first myocardial infarction treated with thrombolysis, **A67**
- mapping**, Characterisation of the electrical wavefronts seen in human permanent AF, **A37**
Characteristics of global electrical restitution determine ventricular electrical stability in man, **A37**
High frequency re-entry circuits driving AF: evidence from global left atrial mapping in humans, **A2**
- Marfan**, An operation for the Marfanoid aorta which spares the aortic valve and the vascular intima: Robisceck's concept realised by a rapid prototyping engineering solution, **A18**
- Marfan's Syndrome**, Presentation and outcome of Marfan's syndrome patients with type B dissection and thoracoabdominal aortic aneurysm, **A19**
- mechanical asynchrony**, Role of real-time transthoracic 3D echocardiography in the assessment of mechanical asynchrony, **A31**
- mechanical heart valves**, Anticoagulation control and outcome in patients with second generation mechanical heart valves, **A19**
- mechanisms**, Interactions between focal triggers and left atrial substrate in the initiation and maintenance of atrial fibrillation: a study using dominant frequency analysis of non-contact mapping data, **A36**
QT peak prolongation may reflect inducible myocardial ischaemia, **A47**
- menu based**, Physical activity behaviour in users of a menu based cardiac rehabilitation programme: a longitudinal survey, **A11**
- metabolic syndrome**, Homocysteine predicts cardiovascular risk independently of the metabolic syndrome, **A42**
- MINAP**, Understanding variation: application of statistical process control to the myocardial infarction national audit project database for 2003–2004, **A41**
- mismatch**, Patient prosthesis mismatch in aortic valve replacement—size does not matter, **A19**
- mitral valve disease**, Plasma N-terminal pro-brain natriuretic peptide predicts exercise capacity in degenerative mitral valve disease, **A49**
- mitral valve surgery**, Edge-to-edge repair for Barlow's disease: early and late clinical and echocardiographic results, **A49**
- MMP-2**, Multimarker approach to risk stratification in 'new' definition myocardial infarction, **A45**
- mortality**, Impact of secondary prevention medication on 30 day mortality for patients admitted to English and Welsh hospitals with acute coronary syndromes, **A25**
Mortality of patients in England with left ventricular systolic dysfunction and heart failure due to other causes, **A8**
Predictors of short term mortality in heart failure: EuroHeart Failure Survey, **A7**
The relationship between troponin levels and 90 day mortality for patients with ST segment infarction; a paradox, **A26**
The true incidence of primary graft dysfunction leading to death in heart transplantation: a validated audit in 1820 patients, **A48**
- mouse models**, Identification of cardiac abnormalities in the Sonic Hedgehog mouse mutant using optical projection tomography, **A34**
Role of the planar cell polarity pathway in the development of the outflow tract, **A34**
- MRI**, Real-time integrated X-ray and magnetic resonance imaging (XMR)—A novel strategy to optimise precise catheter placement during pulmonary vein isolation, **A37**
- multi vessel disease**, Multivessel PCI procedures with implantation of Sirolimus eluting stents: a report from the e-CYPHER registry, **A6**
- multislice computed tomography**, Diagnostic accuracy of non-invasive coronary angiography with multislice computed tomography in patients presenting with acute chest pain, **A15**
Radiation dose in coronary multislice CT: a comparison with conventional diagnostic angiography, **A16**
- murine**, A new murine model of arterial stenting, **A51**
- MYBPC3**, Novel mutations in cardiac MYBPC3 cause early onset malignant hypertrophic cardiomyopathy (HCM), **A69**
- myocardial contrast echocardiography**, Prediction of transmural extent of acute myocardial infarction—comparison between myocardial contrast echocardiography and radionuclide perfusion imaging, **A32**
- myocardial infarction**, Comparison of first and second acute myocardial infarction: recent trends in incidence and case fatality, **A4**
Direct 24-hour ambulance triage and transfer of patients with acute myocardial infarction (AMI) to a cardiac intervention centre with no accident and emergency department, **A59**

- H-FABP: a strong predictor of mortality after 'new' definition myocardial infarction, **A46**
- Left bundle branch block (LBBB) myocardial infarction (MI)—a study of thrombolysis in a district general hospital, **A59**
- Logistics and economics of interhospital transfer v direct admission for primary angioplasty, **A22**
- Multimarker approach to risk stratification in 'new' definition myocardial infarction, **A45**
- Optimising target vessel reperfusion times with primary percutaneous intervention for acute myocardial infarction, **A23**
- Pre hospital ECG diagnosis by ambulance crews reduces time to reperfusion in ST segment elevation myocardial infarction (STEMI) patients treated with primary angioplasty (PA), **A23**
- Selective gene deletion of the neuronal nitric oxide synthase (nNOS) accelerates pathological left ventricular remodelling and functional deterioration after myocardial infarction (MI), **A51**
- The Genetic Risk of Acute Coronary Events (GRACE) Study cohort - demography and description of a large UK discordant-sibling collection, **A41**
- Transmural extent of infarction on contrast enhanced magnetic resonance imaging (CE-MRI) predicts recovery of contractile function in patients with first myocardial infarction treated with thrombolysis, **A67**
- UK Primary Angioplasty Cost Effectiveness Study (UK-PACES) 30 day outcome data, **A27**
- Value of heart fatty acid binding protein (hFABP) for very early detection of acute myocardial infarction (AMI), **A60**
- myocardial perfusion scintigraphy**, Myocardial perfusion scintigraphy predicts functional outcome following percutaneous coronary intervention, **A17**
- myocardial protection**, Evidence for the presence of a second window of preconditioning in humans during serial balloon occlusion at angioplasty on consecutive days, **A40**
- myocardial relaxation**, Cardiac nNOS regulates myocardial relaxation by stimulating phospholamban phosphorylation, **A3**
- NAD(P)H oxidase**, A myocardial nox2 containing NAD(P)H oxidase contributes to oxidative stress in human atrial fibrillation, **A44**
- NADPH oxidase**, NOX-2 (gp91 phox) disruption prevents endothelial ischaemia-reperfusion injury in humans, **A27**
- national service framework**, Rapid access chest pain clinics in the post national service framework era: are protocol driven nurse led clinics safe? A retrospective audit, **A12**
- national statistics**, The growth of cardiac rehabilitation in the UK since 1998, **A10**
- natriuretic peptide**, Screening for suspected heart failure with N terminal pro-B type natriuretic peptide (NT proBNP) in primary care: money well spent?, **A56**
- Suspected heart failure in primary care - the utility of n-terminal pro b-type natriuretic peptide (NT PROBNP) as a pre-screening test for secondary care referral: a real life study, **A22**
- neointima**, A new murine model of arterial stenting, **A51**
- nephrotoxicity**, Calcineurin inhibitor (CI) withdrawal and replacement with sirolimus (SIR) in CTR with renal dysfunction, **A49**
- nitric oxide**, Cardiac nNOS regulates myocardial relaxation by stimulating phospholamban phosphorylation, **A3**
- Nitric oxide is not required for vasculogenesis or angiogenesis in zebrafish embryos, **A28**
- Nitric oxide mediates the effects of vagus nerve stimulation on ventricular fibrillation and electrical restitution in the isolated innervated heart, **A55**
- Prevention of hypoxia-induced pulmonary hypertension and vascular remodelling by targeting endothelial tetrahydrobiopterin-dependent nitric oxide synthase regulation, **A4**
- Selective gene deletion of the neuronal nitric oxide synthase (nNOS) accelerates pathological left ventricular remodelling and functional deterioration after myocardial infarction (MI), **A51**
- nitric-oxide synthase**, Role of the PMCA-Syntrophin-nNOS complex in cardiomyopathy, **A45**
- Tetrahydrobiopterin regulates vagal control of heart rate, **A57**
- nNOS**, Autonomic remodelling in hypertension following cardiac neural nitric oxide synthase (nNOS) gene transfer, **A65**
- non-cardiac disease**, A rapid access cardiology service effectively diagnoses non-cardiac disease and low risk among patients presenting with chest pain, breathlessness, and palpitations, **A13**
- non-compaction**, Isolated left ventricular noncompaction: two year experience with cardiovascular magnetic resonance, **A17**
- Left ventricular non-compaction (LVNC): insights from cardiovascular magnetic resonance (CMR) imaging, **A71**
- non-medical angiographer**, Elective coronary angiography by non-medical angiographers. A comparison with cardiology specialist registrars, **A68**
- non-ST segment elevation acute coronary syndromes**, Use of regional guidelines and rapid access to cardiac catheter laboratories facilitate early invasive management in patients with non-ST segment elevation acute coronary syndromes (NSTEMI), **A62**
- noncontact mapping**, Interactions between focal triggers and left atrial substrate in the initiation and maintenance of atrial fibrillation: a study using dominant frequency analysis of non-contact mapping data, **A36**
- norwood**, Risk factors for sudden death in infants having undergone the Norwood operation for hypoplastic left heart syndrome (HLHS), **A53**
- NT-proBNP**, The impact of the introduction of NT-proBNP into primary care on secondary care referral rates, **A57**
- NTpro-BNP**, Plasma N-terminal pro-brain natriuretic peptide predicts exercise capacity in degenerative mitral valve disease, **A49**
- nurse led**, Rapid access chest pain clinics in the post national service framework era: are protocol driven nurse led clinics safe? A retrospective audit, **A12**
- obesity**, Association between common polymorphisms of the pro-opiomelanocortin (POMC) gene and body fat distribution: a family study, **A41**
- Protection against the development of obesity, insulin resistance, and hypertension in mice by overexpression of insulin-like growth factor binding protein-2 (IGFBP-2), **A51**
- one year outcomes**, A retrospective analysis of 1500 patients undergoing PCI for unstable angina—troponin status at presentation does not predict 1 year outcome, **A25**
- osteopontin**, Calcific crystals directly induce the release of osteopontin from human macrophages via pi3-kinase, ERK1/2 map kinase, and NFκB signalling pathways: implications for atherogenesis and calcific vascular disease, **A14**
- osteoprotegerin**, Plasma osteoprotegerin levels predict the extent of subclinical coronary atherosclerosis and adverse cardiac events in asymptomatic type 2 diabetic subjects, **A17**
- out of hours**, Impact of a 24-hour primary PCI service for ST elevation myocardial infarction (STEMI) on consultants' work patterns, **A27**
- outcome data PCI**, Sirolimus eluting stent and unprotected left main stenosis: the multicenter e-CYPHER registry, **A5**
- Comparison of outcomes in real world use of Sirolimus eluting stents with the randomised controlled clinical trials: can the results be reproduced?, **A6**
- Real life impact of sirolimus eluting stent implantation in treating diabetic patients: a 6 month follow up report from the international e-CYPHER registry, **A5**
- Real world use of sirolimus eluting stents in saphenous vein graft disease: data from the e-CYPHER registry, **A5**
- outcome**, Latest results of endocardial arrhythmia ablation following the modified Fontan, **A34**
- outcome PCI data**, Multivessel PCI procedures with implantation of Sirolimus eluting stents: a report from the e-CYPHER registry, **A6**
- outpatient**, The impact of the introduction of NT-proBNP into primary care on secondary care referral rates, **A57**
- P2X7**, Investigation of P2X7 dependent mechanisms of Interleukin-1 beta (IL-1β) release from human endothelial cells, **A13**
- P2Y1**, Molecular basis for platelet hyper-reactivity to adenosine diphosphate (ADP) associated with the platelet P2Y1 receptor 1622 G allele, **A60**
- pacemaker**, Determinants of early recurrence of AF in pacemaker AF suppression study patients, **A9**
- PAD**, Initial results from the national defibrillator programme (NDP), **A63**
- PAI-1**, Circadian clock genes cause activation of the human PAI-1 gene promoter with 4G/5G allelic preference, **A51**
- palpitation**, Cardiac physiologist led arrhythmia clinic substantially reduces Consultant clinic workload, **A56**
- paramedics**, Delivery of pre-hospital thrombolysis by paramedics, **A59**
- patent foramen ovale**, The role of paradoxical embolism and thrombophilia in ischaemic stroke in young adults, **A33**
- PCI**, Duration of clopidogrel therapy following percutaneous coronary intervention (PCI) with bare metal (BMS) and drug eluting stents (DES). A questionnaire survey in the UK; do we need to be NICE?, **A64**

- Retrospective study to investigate the use of Drug Eluting Stents (DES) for the treatment of In-Stent Restenosis – ‘The Real world’ experience, **A38**
- What is the appropriate ratio between coronary artery bypass graft (CABG) and percutaneous intervention (PCI) in the drug eluting stent era?, **A63**
- percutaneous**, Percutaneous transthoracic epicardial ablation (PTEA) for failed endocardial ablations: promising results of a new technique, **A54**
- percutaneous coronary intervention**, Myocardial perfusion scintigraphy predicts functional outcome following percutaneous coronary intervention, **A17**
- Percutaneous coronary intervention with drug eluting stents for bifurcation lesions: 6 month results, **A6**
- Stent expansion: ‘‘A failure to deliver’’. A combination of delivery balloon constraint and acute stent recoil reduces predicted stent dimension by 27% irrespective of reference vessel size, **A40**
- Successful thrombus extraction with the Rescue Thrombus Management System™ during acute percutaneous coronary intervention (PCI): Does not necessarily restore optimal myocardial tissue perfusion, **A24**
- perfusion**, Prognostic power of normal myocardial perfusion scintigraphy with 99m-Tc agents: the DGH experience, **A16**
- peri-operative**, Peri-operative beta-blockade for patients undergoing infra-renal vascular surgery (POBBLE): results of a randomised double blind controlled trial, **A69**
- pericardial**, Variability in pericardiocentesis practice among cardiologists in the UK—need for guidelines, **A72**
- pericardial window**, Is a pericardial window more effective than conservative treatment in reducing the incidence of post-operative atrial fibrillation (AF) following coronary artery bypass grafting (CABG)?, **A50**
- pericardiocentesis**, Variability in pericardiocentesis practice among cardiologists in the UK—need for guidelines, **A72**
- physical activity**, Physical activity behaviour in users of a menu based cardiac rehabilitation programme: a longitudinal survey, **A11**
- physiotherapy**, Effects of physical training on anxiety and depression in chronic heart failure, **A12**
- platelet aggregation**, Molecular basis for platelet hyper-reactivity to adenosine diphosphate (ADP) associated with the platelet P2Y1 receptor 1622 G allele, **A60**
- PMCA**, Role of the PMCA-Syntrophin-nNOS complex in cardiomyopathy, **A45**
- polymorphism**, Inflammatory genotypes predisposing to premature coronary artery disease (CAD) in a large discordant sibship collection and utility as risk predictors, **A67**
- post-operative**, Does pre-operative atrial cell electrophysiology predict post-operative atrial fibrillation?, **A43**
- pre-excitation syndrome**, Delineating the role of adenosine monophosphate (AMP)-activated protein kinase (AMPK) γ 2 subunit in hypertrophic cardiomyopathy (HCM), **A69**
- pre-hospital**, Delivery of pre-hospital thrombolysis by paramedics, **A59**
- predictors**, Clinical predictors of 6-minute walk test (6-MWT) performance in patients with chronic heart failure (CHF), **A58**
- pregnancy**, Pregnant cardiac patients: development of a generic test to evaluate the functional adequacy of diseased hearts to cope with pregnancy and labour, **A33**
- prehospital diagnosis**, Pre hospital ECG diagnosis by ambulance crews reduces time to reperfusion in ST segment elevation myocardial infarction (STEMI) patients treated with primary angioplasty (PA), **A23**
- prescribing**, Prevalence, incidence, primary care burden, and medical treatment of angina in Scotland: age, sex, and socio-economic disparities, **A40**
- prevalence**, Ethnic differences in the prevalence and aetiology of left ventricular systolic dysfunction (LVD) in high-risk community subjects, **A58**
- prevention**, Using a coronary heart disease (CHD) register to support evidence-based cardiac care in general practice: lessons from the Scottish national CHD demonstration project, **A61**
- primary angioplasty**, Direct 24-hour ambulance triage and transfer of patients with acute myocardial infarction (AMI) to a cardiac intervention centre with no accident and emergency department, **A59**
- Impact of a 24-hour primary PCI service for ST elevation myocardial infarction (STEMI) on consultants’ work patterns, **A27**
- Logistics and economics of interhospital transfer v direct admission for primary angioplasty, **A22**
- Optimising target vessel reperfusion times with primary percutaneous intervention for acute myocardial infarction, **A23**
- Pre hospital ECG diagnosis by ambulance crews reduces time to reperfusion in ST segment elevation myocardial infarction (STEMI) patients treated with primary angioplasty (PA), **A23**
- UK Primary Angioplasty Cost Effectiveness Study (UK-PACES) 30 day outcome data, **A27**
- primary care**, Prevalence, incidence, primary care burden, and medical treatment of angina in Scotland: age, sex, and socio-economic disparities, **A40**
- primary PCI**, Two years of primary coronary intervention for ST elevation myocardial infarction in a district general hospital, **A59**
- primary percutaneous coronary intervention**, Impact of change of protocol for the management of patients with ST elevation myocardial infarction: an 18 month UK tertiary cardiac centre experience, **A26**
- principal component analysis**, Surface atrial frequency analysis in patients with atrial fibrillation: assessing the effect of two standard linear ablation lesions sited in the left atrium, **A10**
- prognosis**, Contemporary prognosis of angina and non-cardiac chest pain at the interface between primary and secondary care: multicentre outcome analysis of 8802 patients attending rapid access chest pain clinic, **A12**
- Effect of dobutamine stress on left ventricular filling in ischaemic cardiomyopathy: pathophysiology and prognostic implications, **A21**
- Elevated serum glucose predicts mortality in patients admitted to hospital with a new diagnosis of heart failure, **A7**
- Ethnicity and variation in prognosis for patients newly hospitalised for heart failure, **A8**
- Haemoglobin is related to prognostic factors but is not an independent survival predictor in patients referred for cardiac transplantation, **A48**
- Indicators of increased risk of mortality following first hospital admission with heart failure: the value of routine clinical variables, **A8**
- Prognostic value of myocardial contrast echocardiography in patients presenting to hospital with acute chest pain and negative troponin, **A31**
- Prognostic value of normal stress echocardiography in patients with suspected Non-ST elevation acute coronary syndrome and negative troponin, **A33**
- prosthetic valve**, Improved haemodynamic profile and left ventricular function following aortic valve replacement with a stentless rather than stented bioprosthesis: a randomised controlled trial, **A18**
- provision**, The growth of cardiac rehabilitation in the UK since 1998, **A10**
- psychological factors**, Effects of physical training on anxiety and depression in chronic heart failure, **A12**
- psychological rehabilitation**, Psychological rehabilitation for cardiac patients: systematic review and meta-analysis, **A11**
- pulmonary hypertension**, Improved survival of systemic sclerosis-associated pulmonary arterial hypertension (SSc-PAH) in the endothelin antagonist era, **A64**
- Prevention of hypoxia-induced pulmonary hypertension and vascular remodelling by targeting endothelial tetrahydrobiopterin-dependent nitric oxide synthase regulation, **A4**
- Prognostic value of brain natriuretic peptide (BNP) in patients with pulmonary arterial hypertension (PAH), **A58**
- Risk factors for sudden death in infants having undergone the Norwood operation for hypoplastic left heart syndrome (HLHS), **A53**
- pulmonary vein**, Ablation of epicardial autonomic ganglionated plexi during minimally invasive surgical ablation of atrial fibrillation, **A9**
- QRS width**, Interrelation between NT-pro BNP, QRS width, and severity of heart failure: EuroHeart Failure Survey, **A6**
- QT dispersion**, Increased QT dispersion as a predictor of mortality following biventricular pacing, **A54**
- Effects of amiloride and spironolactone on QT in stroke survivors—a randomised placebo controlled double-blind cross over study, **A65**
- Stress Induced QT prolongation does not predict Ischaemia in patients with stable Coronary Artery Disease, **A46**
- quadriceps**, Quadriceps and respiratory muscle strength in patients with a Fontan circulation, **A34**

- radial access day case PCI**, Evaluation of Novel dose of C7E3Fab (abciximab) on periprocedural release of Troponin I and Thrombin generation marker (PF1+2) in Radial access Coronary Stenting as Elective Day case procedure 'The ENFORCED study', **A38**
- radial artery**, Transradial coronary intervention without pre-screening for a dual palmar blood supply, **A39**
- radiation dosimetry**, Radiation dose in coronary multislice CT: a comparison with conventional diagnostic angiography, **A16**
- radiofrequency**, Experience with 75 consecutive cryoablation procedures in a single UK tertiary centre, **A10**
- radiofrequency ablation**, Can atrial fibrillation with a coarse ECG appearance be treated by catheter ablation of the tricuspid valve-inferior vena cava isthmus? Preliminary results of a multicentre randomised controlled trial, **A9**
Left atrial radiofrequency ablation during mitral valve surgery for continuous atrial fibrillation: results of a prospective randomised clinical trial, **A1**
- randomised controlled trial**, Home-based versus supervised centre-based cardiac rehabilitation in a multi-ethnic population: 6 month results of the Birmingham Rehabilitation Uptake Maximisation Study (BRUM), **A53**
- rapid access chest pain clinic**, Rapid access chest pain clinics in the post national service framework era: are protocol driven nurse led clinics safe? A retrospective audit, **A12**
Contemporary prognosis of angina and non-cardiac chest pain at the interface between primary and secondary care: multicentre outcome analysis of 8802 patients attending rapid access chest pain clinic, **A12**
- rapid cardiology**, A rapid access cardiology service effectively diagnoses non-cardiac disease and low risk among patients presenting with chest pain, breathlessness, and palpitations, **A13**
- RASSF1**, The tumour suppressor RASSF1A attenuates cardiomyocyte hypertrophy in response to phenylephrine, **A44**
- REACT trial**, The REACT (REscue Angioplasty v Conservative treatment or repeat Thrombolysis) Trial: impact of age upon efficacy of treatment, **A24**
- real world experience**, Comparison of outcomes in real world use of Sirolimus eluting stents with the randomised controlled clinical trials: can the results be reproduced?, **A6**
- rescue angioplasty**, The REACT (REscue Angioplasty v Conservative treatment or repeat Thrombolysis) Trial: impact of age upon efficacy of treatment, **A24**
- regional guidelines**, Use of regional guidelines and rapid access to cardiac catheter laboratories facilitate early invasive management in patients with non-ST segment elevation acute coronary syndromes (NSTEACS), **A62**
- regional transfer**, Rapid reduction of inter hospital Acute Coronary Syndrome (ACS) transfer waiting times; the role of a Regional Transfer Unit (RTU), **A25**
- rehabilitation**, Randomised controlled trial comparing the clinical effectiveness of home based and hospital based rehabilitation after acute myocardial infarction: the Cornwall Heart Attack Rehabilitation Management Study, **A3**
- rejection**, Ultrasonic strain and strain rate imaging: a new non-invasive technique to identify acute rejection after heart transplantation, **A47**
- renal dysfunction**, Relationship between mild renal dysfunction and outcomes after coronary artery bypass grafting, **A19**
- renal failure**, Dobutamine stress echocardiography (DSE) and the resting but not exercise electrocardiograph (ECG) predict severe coronary artery disease (CAD) in end stage renal failure, **A68**
- reperfusion**, Can reperfusion be predicted by acute thickening of the at risk myocardium in stemi? Full pressure v pressure limiting reperfusion, **A32**
- repolarization**, Characteristics of global electrical restitution determine ventricular electrical stability in man, **A37**
- rescue device**, Successful thrombus extraction with the Rescue Thrombus Management System™ during acute percutaneous coronary intervention (PCI): Does not necessarily restore optimal myocardial tissue perfusion, **A24**
- rescue PCI**, Impact of change of protocol for the management of patients with ST elevation myocardial infarction: an 18 month UK tertiary cardiac centre experience, **A26**
The REACT (REscue Angioplasty v Conservative treatment or repeat Thrombolysis) Trial: results according to actual treatment received, **A22**
- rescue percutaneous coronary intervention**, Factors influencing left ventricular function following successful rescue percutaneous coronary intervention for failed thrombolysis in acute myocardial infarction, **A24**
- resistance**, Insulin resistance is a substrate for accelerated endothelial dysfunction in middle age—studies in mice heterozygous for knockout of the insulin receptor gene (IRKO), **A28**
- respiratory muscles**, Quadriceps and respiratory muscle strength in patients with a Fontan circulation, **A34**
- restenosis**, Clinical presentation with in-stent restenosis is not always benign, **A39**
Intracoronary delivery of a secreted transforming growth factor- β type II receptor from a novel collagen-membrane bound stent inhibits in-stent neointimal hyperplasia, **A50**
The role of angiogenic growth factors in intimal hyperplasia, **A1**
- restitution**, Characteristics of global electrical restitution determine ventricular electrical stability in man, **A37**
- restrictive filling**, Effect of dobutamine stress on left ventricular filling in ischaemic cardiomyopathy: pathophysiology and prognostic implications, **A21**
- resuscitation**, False cardiac arrest (FCA) calls—the right time to turn away?, **A63**
Initial results from the national defibrillator programme (NDP), **A63**
- Resynchronisation therapy**, Can real-time 3D echocardiography predict reverse left ventricular remodelling post cardiac resynchronisation therapy?, **A31**
Resynchronization therapy in adults with congenital heart disease, **A35**
- revascularisation**, Low incidence of revascularisation of bare metal stents in the era of drug eluting stents: a single tertiary centre experience, **A5**
- risk factors**, Prospective analysis of the association of mean leucocyte telomere length with risk of coronary heart disease and interaction with statin treatment, **A14**
- risk score**, Indicators of increased risk of mortality following first hospital admission with heart failure: the value of routine clinical variables, **A8**
- risk scoring**, Introduction of a risk scoring system for percutaneous coronary intervention (PCI) procedures, **A63**
- risk stratification**, H-FABP: a strong predictor of mortality after 'new' definition myocardial infarction, **A46**
- saphenous vein grafts**, Real world use of sirolimus eluting stents in saphenous vein graft disease: data from the e-CYPHER registry, **A5**
- scintigraphy**, Prognostic power of normal myocardial perfusion scintigraphy with 99m-Tc agents: the DGH experience, **A16**
- secondary prevention**, Impact of secondary prevention medication on 30 day mortality for patients admitted to English and Welsh hospitals with acute coronary syndromes, **A25**
- sedation**, Safety and acceptability of conscious sedation for sub-pectoral implantation of implantable cardioverter defibrillators (ICDs), **A54**
The safety of DC cardioversion under sedation: long-term experience, **A36**
- signalling**, Myostatin regulates cardiomyocyte growth through modulation of Akt signalling in vitro and in vivo, **A1**
- signalling pathway**, P38-MAPK mediates the early negative inotropic effect of tumour necrosis factor- α (TNF). Evidence of synergy between a direct negative inotropic effect and coronary constriction, **A57**
FoxO transcription factor expressions and activity are regulated during the development of cardiomyocyte hypertrophy, **A44**
- sirolimus**, Calcineurin inhibitor (CI) withdrawal and replacement with sirolimus (SIR) in CTR with renal dysfunction, **A49**
- six-minute walk test**, Clinical predictors of 6-minute walk test (6-MWT) performance in patients with chronic heart failure (CHF), **A58**
- sleep apnoea**, Cardiac dysynchrony as measured by total isovolumic time may predispose to central sleep apnoea in chronic heart failure, **A21**
- smoking**, The effect of smoking and other lifestyle factors on the left ventricular growth response to exercise in 312 young healthy men assessed using cardiovascular magnetic resonance, **A21**
- smooth muscle cells**, Beta-blockers inhibit transgene expression under transcriptional regulation of the major immediate-early cytomegalovirus enhancer/promoter in vivo, **A29**
Tribbles-1, a mitogen activated protein kinase (MAPK) scaffold regulates inflammation in vascular smooth muscle cells (VSMC) and atherosclerosis, **A15**

- South Asian**, Ethnicity and variation in prognosis for patients newly hospitalised for heart failure, **A8**
Declining case fatality rates for acute myocardial infarction in south Asians and whites in the last 15 years, **A41**
Increased arterial stiffness in healthy young South Asian men, **A66**
- SPECT**, Prediction of transmural of acute myocardial infarction—comparison between myocardial contrast echocardiography and radionuclide perfusion imaging, **A32**
- spirinolactone**, Actions of aldosterone blockade in patients with mild-moderately severe heart failure, **A20**
- ST elevation infarction**, The relationship between troponin levels and 90 day mortality for patients with ST segment infarction; a paradox, **A26**
- stable angina**, Anti-anginal and anti-ischaemic efficacy of ivabradine—a selective and specific sinus node If current inhibitor—compared to atenolol in elderly patients with chronic stable angina, **A69**
- statin**, Reversal of vascular dysfunction in subjects at risk of type 2 diabetes mellitus (DM), **A53**
- statins**, Clinical benefits of low LDL targets in elderly patients with coronary disease: An analysis from PROVE IT-TIMI-22, **A46**
Prospective analysis of the association of mean leucocyte telomere length with risk of coronary heart disease and interaction with statin treatment, **A14**
- statistical process control**, Understanding variation: application of statistical process control to the myocardial infarction national audit project database for 2003–2004, **A41**
- stem cells**, Genetic alteration of marrow derived stromal cells with HIF-1 α /VP16 enhances collateral forming potential in vitro and in vivo, **A1**
- STEMI**, Can reperfusion be predicted by acute thickening of the at risk myocardium in stemi? Full pressure v pressure limiting reperfusion, **A32**
Impact of primary angioplasty (PPCI) programme on the management of patients presenting with ST-elevation myocardial infarction (STEMI) in a North West London borough, **A53**
- stent**, A new murine model of arterial stenting, **A51**
Clinical presentation with in-stent restenosis is not always benign, **A39**
Measurement of coronary stent expansion using StentBoost™ image enhancement software: a comparison with intravascular ultrasound, **A39**
Intracoronary delivery of a secreted transforming growth factor- β type II receptor from a novel collagen-membrane bound stent inhibits in-stent neointimal hyperplasia, **A50**
Stent expansion: “A failure to deliver”. A combination of delivery balloon constraint and acute stent recoil reduces predicted stent dimension by 27% irrespective of reference vessel size, **A40**
- strain**, Ultrasonic strain and strain rate imaging: a new non-invasive technique to identify acute rejection after heart transplantation, **A47**
- strain rate imaging**, Ultrasonic strain and strain rate imaging: a new non-invasive technique to identify acute rejection after heart transplantation, **A47**
- stress echocardiography**, Prognostic value of normal stress echocardiography in patients with suspected Non-ST elevation acute coronary syndrome and negative troponin, **A33**
- stroke**, The role of paradoxical embolism and thrombophilia in ischaemic stroke in young adults, **A33**
- sub-pectoral**, Safety and acceptability of conscious sedation for sub-pectoral implantation of implantable cardioverter defibrillators (ICDs), **A54**
- substrate**, The influence of atrial conduction, refractoriness, and volume on outcomes following pulmonary vein isolation (PVI), **A54**
- sudden**, Demographic and temporal trends in out-of-hospital sudden cardiac death in Belfast, **A63**
- superoxide**, A myocardial nox2 containing NAD(P)H oxidase contributes to oxidative stress in human atrial fibrillation, **A44**
Endothelial tetrahydrobiopterin (BH4) regulates eNOS coupling in vivo: functional effects of BH4-eNOS stoichiometry in endothelial targeted transgenic mice, **A50**
- surgery**, An operation for the Marfanoid aorta which spares the aortic valve and the vascular intima: Robiscek’s concept realised by a rapid prototyping engineering solution, **A18**
- survival**, Comparison of first and second acute myocardial infarction: recent trends in incidence and case fatality, **A4**
Improved survival of systemic sclerosis-associated pulmonary arterial hypertension (SSc-PAH) in the endothelin antagonist era, **A64**
Patient prosthesis mismatch in aortic valve replacement—size does not matter, **A19**
Prognostic power of normal myocardial perfusion scintigraphy with ^{99m}Tc agents: the DGH experience, **A16**
- systematic review**, Psychological rehabilitation for cardiac patients: systematic review and meta-analysis, **A11**
- systemic lupus erythematosus**, Altered coronary microvascular function and peripheral endothelial dysfunction in young patients with systemic lupus erythematosus, **A28**
- systemic sclerosis**, Improved survival of systemic sclerosis-associated pulmonary arterial hypertension (SSc-PAH) in the endothelin antagonist era, **A64**
Prognostic value of brain natriuretic peptide (BNP) in patients with pulmonary arterial hypertension (PAH), **A58**
- TAK1**, TGF β 2 Type I receptor interacts with TGF β 2 Activated Kinase (TAK1): implications for controlling cardiac hypertrophy, **A43**
- telomeres**, Prospective analysis of the association of mean leucocyte telomere length with risk of coronary heart disease and interaction with statin treatment, **A14**
- testosterone**, Testosterone inhibits extracellular calcium entry via voltage-gated calcium channels in A7r5 vascular smooth muscle cells at physiological concentrations, **A43**
- tetrahydrobiopterin**, Endothelial tetrahydrobiopterin (BH4) regulates eNOS coupling in vivo: functional effects of BH4-eNOS stoichiometry in endothelial targeted transgenic mice, **A50**
Tetrahydrobiopterin regulates vagal control of heart rate, **A57**
- TGF β 2**, TGF β 2 Type I receptor interacts with TGF β 2 Activated Kinase (TAK1): implications for controlling cardiac hypertrophy, **A43**
- thoracoabdominal aortic aneurysm**, Presentation and outcome of Marfan’s syndrome patients with type B dissection and thoracoabdominal aortic aneurysm, **A19**
- three-dimensional**, Can real-time 3D echocardiography predict reverse left ventricular remodelling post cardiac resynchronisation therapy?, **A31**
3D contraction front mapping: novel, high resolution imaging of mechanical contraction of the left ventricle, **A30**
- 3D echo**, New generation three-dimensional echocardiography for left ventricular volumetric and functional measurements: comparison with cardiac magnetic resonance, **A30**
Role of real-time transthoracic 3D echocardiography in the assessment of mechanical asynchrony, **A31**
Selective gene deletion of the neuronal nitric oxide synthase (nNOS) accelerates pathological left ventricular remodelling and functional deterioration after myocardial infarction (MI), **A51**
- threshold**, Dofetilide significantly reduces acute and chronic defibrillation thresholds in patients with an inadequate defibrillation safety margin, **A35**
- thrombolysis**, Delivery of pre-hospital thrombolysis by paramedics, **A59**
Impact of change of protocol for the management of patients with ST elevation myocardial infarction: an 18 month UK tertiary cardiac centre experience, **A26**
Impact of primary angioplasty (PPCI) programme on the management of patients presenting with ST-elevation myocardial infarction (STEMI) in a North West London borough, **A53**
Left bundle branch block (LBBB) myocardial infarction (MI)—a study of thrombolysis in a district general hospital, **A59**
Transmural extent of infarction on contrast enhanced magnetic resonance imaging (CE-MRI) predicts recovery of contractile function in patients with first myocardial infarction treated with thrombolysis, **A67**
- thrombus**, Successful thrombus extraction with the Rescue Thrombus Management System™ during acute percutaneous coronary intervention (PCI): Does not necessarily restore optimal myocardial tissue perfusion, **A24**
Thrombogenicity of flowing blood is increased in patients with type 2 diabetes mellitus (T2DM) despite treatment with aspirin, **A61**
- thyroid**, Association between serum thyrotrophin concentrations and occurrence of atrial fibrillation in a large community-based population of elderly subjects (The Birmingham Elderly Thyroid Study), **A40**
- TIMI frame count**, A prospective, randomised, controlled trial to study the effect of intra coronary injection of verapamil and adenosine on coronary blood flow during urgent percutaneous coronary intervention, **A23**
- TnI**, Multimarker approach to risk stratification in ‘new’ definition myocardial infarction, **A45**

- transcranial Doppler**, The role of paradoxical embolism and thrombophilia in ischaemic stroke in young adults, **A33**
- transfer**, How long do patients wait with acute coronary syndromes for inter-hospital transfer in the UK? Results from the coronary heart disease collaborative national audit 2004, **A25**
- transplantation**, The true incidence of primary graft dysfunction leading to death in heart transplantation: a validated audit in 1820 patients, **A48**
- transvenous**, A novel rectangular biphasic waveform from a radiofrequency defibrillator compared with a conventional waveform for the transvenous cardioversion of chronic atrial fibrillation in patients, **A9**
- trends**, Comparison of first and second acute myocardial infarction: recent trends in incidence and case fatality, **A4**
- tribbles**, Tribbles-1, a mitogen activated protein kinase (MAPK) scaffold regulates inflammation in vascular smooth muscle cells (VSMC) and atherosclerosis, **A15**
- tricuspid valve-inferior vena cava isthmus**, Can atrial fibrillation with a coarse ECG appearance be treated by catheter ablation of the tricuspid valve-inferior vena cava isthmus? Preliminary results of a multicentre randomised controlled trial, **A9**
- triple cardiac marker**, Is it safe to discharge patients from accident and emergency using a rapid point of care triple cardiac marker test to rule out acute coronary syndrome in patients presenting with chest pain?, **A26**
- troponin**, A retrospective analysis of 1500 patients undergoing PCI for unstable angina—troponin status at presentation does not predict 1 year outcome, **A25**
Clopidogrel preloading and myocardial necrosis after PCI, **A38**
Evaluation of Novel dose of C7E3Fab (abciximab) On periprocedural release of Troponin I and Thrombin generation marker (PF1+2) in Radial access Coronary Stenting as Elective Day case procedure 'The ENFORCED study', **A38**
Patients with non-ST elevation acute coronary syndromes associated with myonecrosis have a similar rate of death or MI as ST elevation MI at 6 months: An audit of all ACS admissions in a UK District General Hospital over 1 year, **A42**
The relationship between troponin levels and 90 day mortality for patients with ST segment infarction; a paradox, **A26**
Troponin T (TnT) and Creatine Kinase MB (CK-MB) release following Percutaneous Coronary Intervention (PCI): Are they equivalent?, **A24**
- type 2 diabetes**, Plasma osteoprotegerin levels predict the extent of subclinical coronary atherosclerosis and adverse cardiac events in asymptomatic type 2 diabetic subjects, **A17**
Prognostic value of coronary artery calcium screening in asymptomatic patients with type 2 diabetes, **A16**
- UK transplant centres**, Usage of implantable defibrillators and biventricular pacemakers in cardiac transplant candidates: a survey of UK transplant centres, **A48**
- ultrasound**, Measurement of coronary stent expansion using StentBoost™ image enhancement software: a comparison with intravascular ultrasound, **A39**
Rapid pulmonary vein (PV) electrical isolation using a novel focused ultrasound balloon catheter in patients with atrial fibrillation (AF), **A55**
- unstable angina**, A retrospective analysis of 1500 patients undergoing PCI for unstable angina—troponin status at presentation does not predict 1 year outcome, **A25**
- vagus nerve**, Nitric oxide mediates the effects of vagus nerve stimulation on ventricular fibrillation and electrical restitution in the isolated innervated heart, **A55**
- variation**, Understanding variation: application of statistical process control to the myocardial infarction national audit project database for 2003–2004, **A41**
- vascular smooth muscle cell**, Inhibitory Smads regulate pathological cellular differentiation in atherosclerosis, **A29**
- vasculogenesis**, Nitric oxide is not required for vasculogenesis or angiogenesis in zebrafish embryos, **A28**
- vasodilatation**, The adipocyte derived peptide hormone leptin is an endothelial independent vasodilator in humans: implications for vascular homeostasis, **A28**
- vein graft**, Gene transfer of a broad spectrum CC-chemokine inhibitor reduces macrophage recruitment and smooth muscle content in vein graft atherosclerosis in apolipoprotein E-knockout mice, **A13**
- venous tone**, Effects of bradykinin (BK) on venous capacitance in health and treated chronic heart failure (CHF), **A65**
- ventricular**, The effect of smoking and other lifestyle factors on the left ventricular growth response to exercise in 312 young healthy men assessed using cardiovascular magnetic resonance, **A21**
- ventricular dysfunction**, Resynchronization therapy in adults with congenital heart disease, **A35**
- ventricular fibrillation**, Nitric oxide mediates the effects of vagus nerve stimulation on ventricular fibrillation and electrical restitution in the isolated innervated heart, **A55**
- ventricular function**, New generation three-dimensional echocardiography for left ventricular volumetric and functional measurements: comparison with cardiac magnetic resonance, **A30**
- ventricular remodelling**, Effects of optimal pharmacological treatment on left ventricular remodelling, systolic and diastolic function in patients with chronic heart failure, **A20**
- verapamil**, A prospective, randomised, controlled trial to study the effect of intra coronary injection of verapamil and adenosine on coronary blood flow during urgent percutaneous coronary intervention, **A23**
- von Willebrand factor**, Circulating endothelial cells, von Willebrand factor, and flow mediated dilatation as indices of endothelial damage/dysfunction in patients with acute coronary syndromes, **A14**
Increased numbers of circulating endothelial cells predict adverse cardiovascular events following an acute coronary syndrome, **A27**
- vulnerable plaque**, Circulating systemic levels of interleukin-6 and interleukin-10. Do they indicate the presence of a vulnerable plaque?, **A15**
- waiting list**, Rapid reduction of inter hospital Acute Coronary Syndrome (ACS) transfer waiting times; the role of a Regional Transfer Unit (RTU), **A25**
- wave speed**, Effect of nitrate administration on wave speed propagation in the human coronary artery, **A66**
- wide area isolation**, Does increased area of pulmonary vein isolation decrease the number of non pulmonary initiators of atrial fibrillation?, **A36**