

26. **Blackledge HM**, Newton J, Squire IB. Prognosis for South Asian and white patients newly admitted to hospital with heart failure in the United Kingdom: historical cohort study. *BMJ* 2003;**327**:526–31.
27. **Wang M**, Yip GW, Wang AY, *et al*. Peak early diastolic mitral annulus velocity by tissue Doppler imaging adds independent and incremental prognostic value. *J Am Coll Cardiol* 2003;**41**:820–6.
28. **Nagueh SF**, Middleton KJ, Kopelen HA, *et al*. Doppler tissue imaging: a noninvasive technique for evaluation of left ventricular relaxation and estimation of filling pressures. *J Am Coll Cardiol* 1997;**30**:1527–33.
29. **Ommen SR**, Nishimura RA, Appleton CP, *et al*. Clinical utility of Doppler echocardiography and tissue Doppler imaging in the estimation of left ventricular filling pressures: a comparative simultaneous Doppler-catheterization study. *Circulation* 2000;**102**:1788–94.
30. **Mullens W**, Borowski AG, Curtin RJ, *et al*. Tissue Doppler imaging in the estimation of intracardiac filling pressure in decompensated patients with advanced systolic heart failure. *Circulation* 2009;**119**:62–70.
31. **Geske JB**, Sorajja P, Nishimura RA, *et al*. Evaluation of left ventricular filling pressures by Doppler echocardiography in patients with hypertrophic cardiomyopathy: correlation with direct left atrial pressure measurement at cardiac catheterization. *Circulation* 2007;**116**:2702–8.
32. **Yamamoto T**, Oki T, Yamada H, *et al*. Prognostic value of the atrial systolic mitral annular motion velocity in patients with left ventricular systolic dysfunction. *J Am Soc Echocardiogr* 2003;**16**:333–9.
33. **Rosen BD**, Edvardsson T, Lai S, *et al*. Left ventricular concentric remodeling is associated with decreased global and regional systolic function: the Multi-Ethnic Study of Atherosclerosis. *Circulation* 2005;**112**:984–91.
34. **de Simone G**, Kitzman DW, Chinali M, *et al*. Left ventricular concentric geometry is associated with impaired relaxation in hypertension: the HyperGEN study. *Eur Heart J* 2005;**26**:1039–45.
35. **Levy D**, Garrison RJ, Savage DD, *et al*. Left ventricular mass and incidence of coronary heart disease in an elderly cohort. The Framingham Heart Study. *Ann Intern Med* 1989;**110**:101–7.
36. **Kumaran K**, Fall CH, Martyn CN, *et al*. Left ventricular mass and arterial compliance: relation to coronary heart disease and its risk factors in South Indian adults. *Int J Cardiol* 2002;**83**:1–9.
37. **Gardin JM**, Henry WL, Savage DD, *et al*. Echocardiographic measurements in normal subjects: evaluation of an adult population without clinically apparent heart disease. *J Clin Ultrasound* 1979;**7**:439–47.
38. **Tripodkiadis F**, Tentolouris K, Androulakis A, *et al*. Left atrial mechanical function in the healthy elderly: new insights from a combined assessment of changes in atrial volume and transmitral flow velocity. *J Am Soc Echocardiogr* 1995;**8**:801–9.
39. **Thomas L**, Levett K, Boyd A, *et al*. Compensatory changes in atrial volumes with normal aging: is atrial enlargement inevitable? *J Am Coll Cardiol* 2002;**40**:1630–5.

Correction

Khattab AA, Hamm CW, Senges S, *et al*. Sirolimus-eluting stent treatment for unprotected versus protected left main coronary artery disease in widespread clinical routine: 6-month and 3-year clinical follow-up results from the prospective multicentre German Cypher Registry. *Heart* 2007;**93**:1251–5. This article was printed with an incorrect DOI. The DOI should be 10.1136/hrt.2006.104703.

Heart 2010;**96**:471. doi:10.1136/hrt.2007.104703corr2