e0563 IMPLANTATION OF LEFT VENTRICULAR EPICARDIAL LEAD GUIDED BY OVERLAY REF FOR CARDIAC RESYNCHRONISATION THERAPY

doi:10.1136/hrt.2010.208967.563

Tang Kai, Zhao Dongdong, Xu Yawei. ¹Shanghai 10th People's Hospital, Tongji University School of Medicine

Objective This study was to report our initial experiences with cardiac resynchronisation therapy (CRT) guided by Overlay Ref for the treatment of heart failure.

Methods Eleven patients with heart failure were enrolled in this study, with ($24\% \pm 11\%$) of mean left ventricular ejection fraction (LVEF) and ($78 \text{ mm} \pm 13 \text{ mm}$) of left ventricular end-diastolic dimension (LVEDD). Using Overlay Ref technique, a reference image (inverted) of coronary sinus (CS) and its branches was faded into the live fluoroscopic image. The guidewire (originally designed for percutaneous transluminal coronary angioplasty) could be placed to the target branch of CS guided by Overlay Ref image. Then the left ventricular epicardial lead was positioned to the desired spot through the guidewire.

Results All patients were successfully implanted with LV leads guided by Overlay Ref. Exposure time to X-ray was ($19.7 \text{ min} \pm 14.2 \text{ min}$). And the total duration of procedure was $99 \text{ min} \pm 42 \text{ min}$. 2 weeks after the implantation procedure, ultracardiography (UCG) test showed that LVEF of this patient was raised to ($28\% \pm 9\%$).

Conclusion Overlay Ref technique could facilitate the procedure of LV lead implantation for CRT.

e0564 EFFICACY AND SAFETY OF IMPLANTABLE CARDIOVERTER DEFIBRILLATOR AVOIDING ROUTINE DEFIBRILLATION THRESHOLD TESTING

doi:10.1136/hrt.2010.208967.564

¹Liu Qiming, ¹Zhou Shenghua, ¹Qi Shushan, ²Zeng Gaofeng, ³Ma Xiaofeng, ⁴Huang He. ¹Department of Cardiology, Second Xiangya Hospital, Central South University, Changsha; ²Department of Cardiology, Second Affiliated Hospital, Nanhua University, Henyang Hunan; ³Department of Cardiology, Nanhua Hospital, Nanhua University, Henyang Hunan; ⁴Department of Cardiology, Xiangtan Central Hospital, Xiangtan Hunan

Objective To evaluate the efficacy and safety of implantable cardioverter defibrillator (ICD) and cardiac resynchronisation therapy-defibrillators (CRT-D) avoiding defibrillation threshold (DFT) testing when treating ventricular tachycardia (VT) or ventricular fibrillation (VF).

Methods We analysed a continuous database of the 21 patients who had avoided DFT during ICD implantation from Oct. 1999 to Aug. 2008. Follow-up data were completed and analysed in the 21 patients with ICD implantation.

Results ICDs were implanted successfully in 17 patients with VT or VF, and CRT-D were implanted successfully in 4 myocardial infarction patients with severe heart failure who avoided DFT during ICD or CRT-D implantation. Eight patients accepted DFT 1 week later, VT or VF was not induced in 3 patients (37.5%). During the mean follow-up of 1–7 (4.2 ± 1.9) yrs, malignant ventricular arrhythmia was recorded in 16 patients. Among them, 89 episodes were successfully terminated by defibrillation (100%), 120 VT events were terminated by the first run of antitachycardia pacing (51.1%) and 22 by low energy cardioversion (59.2%). All patients took antiarrhythmia drugs after ICD or CRT-D implantation. No patient died from malignant ventricular arrhythmia during the follow-up.

Conclusion No application of routine DFT may avoid complications associated with DFT during ICD or CRT-D implantation. ICD or CRT-D implantation may effectively treat fatal ventricular tachyarrhythmias and prevent sudden cardiac death.

e0565 ANALYSIS OF INAPPROPRIATE THERAPY IN THE PATIENTS IMPLANTABLE CARDIOVERTER DEFIBRILLATORS

doi:10.1136/hrt.2010.208967.565

Wang Hui, Qu baiming, Wu Lixuan. Department of Cardiology Zhejiang Provincial People's Hospital Hangzhou

Objective Observed the efficacy of the patients with implantable cardioverter defibrillators (ICD). Sought to identify the reasons of inappropriate shocks, to minimise the risk of inappropriate shocks.

Methods Followed up nineteen patients with ICD. Stored ICD electrograms from all shocks episodes were analysed to distinguish appropriate and inappropriate shocks according to history of tachycardia. Any ICD therapy not delivered for VT or VF was deemed inappropriate shocks therapy. It can be considered as inappropriate that ICD firing for sinus tachycardia, supraventricular tachycardia, atrial fibrillation, atrial flutter, and nonsustained ventricular tachycardia.

Result There is no shock episode in ten patients (52.63%). Appropriate shocks occurred in 6 patients (31.58%) Inappropriate shocks occurred in three patients (15.79%). Inappropriate shock episodes constituted 4 of 11 total shock episodes. We presented three cases of inappropriate shocks due to atrial fibrillation, sinus tachycardia and high frequency noise.

Conclusions Inappropriate shocks occurred commonly in our cases. Varied reason induced the inappropriate shocks. Programming the device parameter and choosing proper discriminators can minimised rate of inappropriate shocks.

e0566 CORRELATION OF ECG FINDINGS WITH SYMPTOMS OF PALPITATIONS USING NOVEL MULTI-LEAD MOBILE PHONE ECG

doi:10.1136/hrt.2010.208967.566

Michael CL Lim. Singapore Medical Specialists Centre

Objective The aim of this study is to correlate the ECG findings with symptoms of palpitations using a novel multi-lead mobile phone ECG.

Methods A total of 39 consecutive patients with symptoms of palpitations were enrolled for a period of 3–6 months. Standard 12 lead baseline ECGs were performed and patients were divided into 4 main categories of ventricular conduction defects; 1) sinus tachycardia, 2) supraventricular tachycardia, 3) ventricular ectopics and 4) non-sustained ventricular tachycardia. Patients were each given a mobile phone ECG (EPI Life) to record their ECGs during symptoms. ECG recordings were correlated with symptoms including palpitations.

Results Of the 130 ECG recordings assessed, 33% showed sinus tachycardia, 25% supraventricular tachycardia, 41% ventricular ectopics and 1% non-sustained ventricular tachycardia. All patients were symptomatic during ECG recordings.

Conclusions Mobile phone ECG (EPI Life) is a useful diagnostic tool for early detection of arrhythmias and prevention of morbidities and mortalities.

Clinical and Research Medicine: Arrhythmias and Intervention Therapy

e0567 EARLY DIAGNOSIS AND RESCUE PERICARDIOCENTESIS FOR ACUTE CARDIAC TAMPONADE DURING RADIOFREQUENCY ABLATION FOR ARRHYTHMIAS, IS FLUOROSCOPY ENOUGH?

doi:10.1136/hrt.2010.208967.567

¹Huang Xinmiao, ¹Hu Jianqiang, ²Zhou Fei, ¹Qin Yongwen, ¹Cao Jiang, ¹Zhou Bingyan. ¹Changhai Hospital, Second Military Medical University; ²Department of Cardiovascular Diseases, No. 117 Hospital of PLA, Hangzhou, China

Background With the number of complex catheter ablation procedures increasing, procedure-related acute cardiac tamponade is