GW23-e1874

SIMVASTATIN PREVENT ATHEROSCLEROSIS DEVELOPMENT THROUGH INHIBITING PROTEIN EXPRESSION OF ADHESION MOLECULES

doi:10.1136/heartjnl-2012-302920b.37

Li xiaoyan, Lu ying, Zhang ping, Qu caihong. *The Third Affiliated Hospital of Sun Yat-sen University*

Objectives To explore the effects of simvastatin on protein expressions of intercellular adhesion molecule 1 (ICAM-1), vascular cell adhesion molecule 1 (VCAM-1) and endothelial leukocyte adhesion molecule 1 (E-selectin) of cholesterol diet rabbits and the underlying mechanisms of the beneficial effects of simvastatin on atherosclerosis (AS).

Methods Twenty-four male rabbits were randomly divided into three groups: normal diet groups, cholesterol diet groups and the simvastatin groups which received both cholesterol diet and simvastatin 5 mg/kg/d intragastrically. ELISA was used to detect plasma protein levels of ICAM-1, VCAM-1 and E-selectin of rabbits at 0, 8 and 16 weeks, respectively. After 16 weeks, rabbits were executed and the artery wall was harvested for the pathologic and morphologic observations. Western blot and RT-PCR were used to

determine ICAM-1, VCAM-1 and E-selectin level of rabbits artery wall.

Results Compared with normal diet groups, the expression of ICAM-1, VCAM-1 and E-selectin in cholesterol diet groups plasma and artery wall was obviously increased both in protein and mRNA level. ($n=6$; $p<0.01$). However, simvastatin could inhibit the formation of atherosclerotic plaques obviously; suppress plasma and artery wall protein expression and mRNA level of ICAM-1, VCAM-1 and E-selectin ($n=6$; $p<0.01$).

Conclusions Simvastatin could ameliorates atherosclerosis (AS) through inhibiting expression level of plasma and artery wall adhesion molecule.