

ORIGINAL ARTICLE

Osteoprotegerin and cardiovascular mortality in patients with non-ST elevation acute coronary syndromes

Ragnhild Røysland,^{1,2} Marc P Bonaca,^{3,4} Torbjørn Omland,^{1,2} Marc Sabatine,^{3,4} Sabina A Murphy,³ Benjamin M Scirica,^{3,4} Mette Bjerre,^{5,6} Allan Flyvbjerg,^{5,6} Eugene Braunwald,^{3,4} David A Morrow^{3,4}

¹Division of Medicine, Akershus University Hospital, Lørenskog, Norway

²K.G. Jebsen Cardiac Research Centre and Centre for Heart Failure Research, University of Oslo, Oslo, Norway

³TIMI Study Group, Brigham and Women's Hospital, Boston, Massachusetts, USA

⁴Cardiovascular Division, Brigham and Women's Hospital, Boston, Massachusetts, USA

⁵Department of Endocrinology and Internal Medicine, Aarhus University Hospital, Aarhus, Denmark

⁶The Medical Research Laboratories, Department of Clinical Medicine, Faculty of Health Sciences, Aarhus University, Aarhus, Denmark

Correspondence to

Professor T Omland, Division of Medicine, Akershus University Hospital, Sykehusveien 27, 1478 Lørenskog, Norway; torbjorn.omland@medisin.uio.no

Accepted 17 January 2012

ABSTRACT

Objective To assess the relationship between osteoprotegerin (OPG) and cardiovascular death, and the pathobiological mechanisms contributing to the association, in acute coronary syndromes (ACS).

Design Prospective observational.

Setting Biomarker substudy of MERLIN-TIMI 36, a randomised, placebo controlled trial of ranolazine in non-ST elevation (NSTEMI)-ACS.

Patients 4463 patients with NSTEMI-ACS.

Interventions Ranolazine or placebo.

Main outcome measures Incidence of cardiovascular death (CV death); additionally, heart failure (HF), cardiac arrhythmias, in-hospital ischaemia, severe recurrent ischaemia or recurrent myocardial infarction (MI).

Results During a median follow-up of 341 days, 208 patients died of cardiovascular causes. The OPG baseline concentration was strongly associated with both 30 day and 1 year incidence of CV death. After adjustment for conventional risk markers, OPG concentrations (log transformed) remained a significant predictor of CV death by 30 days (HR (95% CI) 2.32 (1.30 to 4.17); $p=0.005$) and by 1 year (HR 1.85 (1.33 to 2.59); $p<0.001$). Baseline levels of OPG were also an independent predictor of new or worsening HF at 30 days (HR 2.25 (1.38 to 3.69); $p=0.001$) and 1 year (HR 1.81 (1.26 to 2.58) $p=0.001$). By univariable analysis, higher OPG was associated with both early ischaemic and arrhythmic events. Although OPG levels were associated with recurrent MI within 12 months, this association was attenuated and no longer significant after multivariable adjustment.

Conclusions OPG is independently associated with 30 day and 1 year risk of cardiovascular mortality and HF development after NSTEMI-ACS. As no independent relationship between OPG levels and recurrent ischaemia or MI was observed, myocardial dysfunction may be a more important stimulus for OPG production than ischaemia in ACS.

INTRODUCTION

The biomarker osteoprotegerin (OPG) has been linked to coronary atherosclerosis, coronary plaque stability and inflammation.^{1–3} OPG is a member of the tumour necrosis factor receptor superfamily and was first identified as a regulator of bone resorption.⁴ By binding the receptor activator of

nuclear factor κ B ligand (RANKL), acting as a decoy receptor to competitively inhibit RANKL interaction with its receptor, RANK, OPG inhibits osteoclastogenesis. OPG is expressed in the vascular system in both endothelial and smooth muscle cells and is also present in atherosclerotic plaques and in early atherosclerotic lesions.⁵ Mice deficient in OPG develop severe osteoporosis as well as medial calcification of the aorta and renal arteries.⁶ In human studies, OPG levels have been associated with traditional cardiovascular risk factors such as increasing age, decreased kidney function and duration of diabetes.^{7–9} It has also been associated with prevalence and severity of peripheral vascular disease¹⁰ and prognosis in cerebrovascular disease.¹¹ In coronary artery disease (CAD), OPG has been found to be associated with the severity¹² and number of coronary artery plaques.¹ In addition, OPG levels are associated with coronary artery calcification on CT,¹ as well as the risk of developing cardiovascular disease in the general population.¹³

OPG is inversely related to indices of left ventricular function in the general population¹⁴ and has been associated with the development and progression of heart failure. Accordingly, it has previously been shown that both experimental and clinical heart failure is associated with increased expression of the OPG/RANKL/RANK axis.¹⁵ Moreover, in patients with acute myocardial infarction (AMI), OPG levels reflect final infarct size¹⁶ and predict outcome in patients presenting with post-infarction heart failure (HF).¹⁷ In a study of 897 patients with acute coronary syndrome (ACS), we have previously observed that OPG levels obtained early after admission for ACS were associated with long term all-cause mortality.¹⁸

To evaluate the relationship between OPG and cardiovascular outcomes and to further explore the underlying pathobiological mechanisms contributing to the prognostic value of OPG in ACS, we analysed baseline blood samples from 4463 participants in the MERLIN (Metabolic Efficiency with Ranolazine for Less Ischaemia in NSTEMI-ACS)-TIMI 36 trial, a large, well characterised, contemporary population of patients with non-ST elevation (NSTEMI)-ACS. In contrast with prior work, we investigated the association between OPG and cardiovascular mortality, and assessed whether the increased risk of death associated with higher levels



This paper is freely available online under the BMJ Journals unlocked scheme, see <http://heart.bmj.com/site/about/unlocked.xhtml>

of OPG could be attributed to one or more of the main determinants of mortality after ACS—that is, electrical instability, left ventricular dysfunction or recurrent ischaemia. Accordingly, we assessed associations between OPG and the incidence of cardiac arrhythmias, HF development and recurrent ischaemic events.

METHODS

Study population

This is a substudy of the MERLIN-TIMI 36 trial, and the design, entry criteria and main results have been previously described.^{19,20} In brief, the MERLIN-TIMI 36 trial was a randomised, controlled, multicentre study of 6560 patients with NSTEMI-ACS treated with ranolazine or placebo. Eligible patients had at least 10 min of ischaemic symptoms at rest and presented with one of the following: elevated biomarkers of myonecrosis, ST segment depression ≥ 0.1 mV, a history of diabetes mellitus or an intermediate to high (≥ 3) Thrombolysis in Myocardial Infarction (TIMI) risk score. Exclusion criteria included persistent ST segment elevation, end stage renal disease requiring dialysis, cardiogenic shock or a life expectancy < 12 months. The primary efficacy endpoint was a composite of cardiovascular death, AMI or recurrent ischaemia during the study period. The major safety endpoints were death from any cause and symptomatic documented arrhythmia.

In this biomarker substudy, we measured OPG levels at enrolment in all available plasma samples ($n=4463$). We investigated OPG with a primary interest in the risk of cardiovascular death. In addition, we examined the potential mechanisms of adverse cardiovascular outcomes by assessing recurrent ischaemia, MI or development of new or worsening HF during follow-up, as well as development of symptomatic documented arrhythmias on continuous ECG recordings from the first 7 days of follow-up. Endpoints were adjudicated by a blinded clinical events committee. The protocol was approved by the relevant institutional review boards, and informed consent was obtained from all patients prior to study commencement.

Blood sampling procedures and biochemical assays

Samples of venous blood were obtained at randomisation in ethylenediamine tetraacetic acid anticoagulated plastic tubes, and plasma was isolated within 60 min of sample acquisition. Plasma samples were stored in plastic cryovials at -20°C or colder at the enrolling site until shipped to the TIMI Biomarker Core Laboratory (Boston, Massachusetts, USA) where they were maintained at -80°C or colder. The samples were shipped to Aarhus University Hospital, Aarhus, Denmark, for analysis.

Plasma levels of OPG were quantified by an inhouse time resolved immunofluorometric assay, using commercially available monoclonal antibodies (DY085E from R&D Systems, Abingdon, UK). The method has been described previously.²¹ The limit of detection was 15 ng/l. A plasma control was included in every OPG assay. The intra- and inter-assay variations (%) were below 5% and 9%, respectively, at concentration of 1000 ng/l. In a population based sample of 216 subjects (39% male, mean age 43 (SD 11) years) without a history of cardiovascular disease, hypertension or diabetes, the median OPG concentration was 1164 ng/l (5th percentile 733 ng/l, 95th percentile 2211 ng/l). C reactive protein (CRP) was measured by an established high sensitivity (hs) nephelometric method (Siemens). The assay range for hs-CRP was 0.05–10 mg/l, with a limit of detection of 0.03 mg/l and coefficient of variation of 5.1%, 2.2% and 2.5% at hs-CRP concentrations of 0.17, 1.16 and 1.88 mg/l, respectively. Levels of cardiac troponin I (cTnI) were

measured using the TnI ultra assay (Siemens Healthcare Diagnostics Inc, Deerfield, Illinois, USA) and B type natriuretic peptide (BNP) was measured using the ADVIA Centaur (Siemens Healthcare Diagnostics Inc, Deerfield, Illinois, USA).

The assay for cTnI has a lower level of detection of 6 ng/l, a 99th percentile value of 40 ng/l and a total imprecision of 10% at a concentration of 30 ng/l.²² For BNP, the decision limit was 80 ng/l.^{23,24} The assay range for BNP was 0.5–5000 ng/l with a coefficient of variation of 3.4%, 2.9% and 2.4% at BNP concentrations of 48, 461 and 1768 ng/l, respectively. All biomarker testing was performed by personnel blinded to clinical outcomes and treatment allocation.

Statistical analysis

Plasma concentrations of OPG are reported using the median and IQR of values. Cut-offs for OPG categories correspond to the 33rd and 67th percentiles. Categorical baseline characteristics were compared using χ^2 tests for categorical variables and are presented as proportions; continuous variables were compared using the Wilcoxon rank sum test. Multivariate logistic regression analysis was performed to identify independent predictors of increased circulating OPG levels at baseline. Cox proportional hazard regression was performed to assess associations between OPG levels and the outcome measures at the 30 day and 1 year follow-up periods.

Both unadjusted and covariate adjusted analyses were performed. In the covariate adjusted analysis, all elements of the TIMI risk score for NSTEMI-ACS, including cTnI ultra, as well as history of chronic HF, creatinine clearance < 60 ml/min, gender, BMI, BNP and CRP were included. OPG was entered both as a continuous (log transformed) variable and as a categorical variable (ie, divided in tertiles). The discriminative value of adding OPG to a predefined risk model was evaluated by calculating the category-less net reclassification improvement.²⁵

Analysis was performed using STATA V.10.1 (StataCorp). A *p* value (two tailed) < 0.05 was considered to indicate statistical significance.

RESULTS

Baseline characteristics

Median time from onset of ischaemic symptoms to randomisation was 24 h (IQR 13–34 h). Median value of OPG was 1632 ng/l (IQR 1206–2242 ng/l), and cut-off levels between tertiles were 1345 ng/l and 2000 ng/l, respectively. There was a very weak, but statistically significant, association between time from symptom onset and OPG levels at baseline ($r=0.03$; $p=0.022$). Characteristics of the patients according to OPG tertile 3 versus tertiles 1 and 2 at baseline are shown in table 1. Patients with higher OPG levels were more likely to be older, to be female, to have an index diagnosis of NSTEMI, a left ventricular ejection fraction $\leq 50\%$, a BNP value > 80 ng/l and a positive cTnI (table 2). A history of diabetes, hypertension, chronic HF and impaired renal function, expressed as estimated creatinine clearance < 60 ml/min, were more frequent in the highest versus the two lower OPG tertiles. In addition, in patients without known diabetes, a fasting glucose > 7 mmol/l (126 mg/100 ml) or non-fasting glucose ≥ 11 mmol/l (200 mg/100 ml) was more frequent in the highest versus the two lower OPG tertiles ($p=0.009$). In contrast, a history of dyslipidaemia, a family history of CAD and to be a current smoker were more frequent in the two lower OPG tertiles. The frequency of prior angina, MI, percutaneous coronary intervention or coronary artery bypass graft did not differ between the third versus the first and second OPG tertiles. By multivariable

logistic regression analysis, age ≥ 75 years, a history of diabetes and estimated creatinine clearance < 60 ml/min were the variables most closely associated with higher baseline OPG levels (table 3).

Relationship to cardiovascular death

During a median follow-up time of 341 days, 208 patients died of cardiovascular causes. The concentration of OPG at baseline was strongly associated with cardiovascular mortality at both 30 days and 1 year (figure 1). By univariable analysis, the HR associated with 1 SD increase in logarithmically transformed OPG levels at baseline was 4.67 (95% CI 2.85 to 7.67; $p < 0.001$) at 30 days and 3.85 (95% CI 2.92 to 5.08; $p < 0.001$) after 1 year (table 4). After adjustment for conventional risk markers, including TIMI risk score covariates, a history of chronic HF, creatinine clearance < 60 ml/min, gender, BMI, BNP, CRP and OPG concentrations remained significantly associated with cardiovascular mortality after 30 days (HR (95% CI) 2.32 (1.30 to 4.17); $p = 0.005$) and 1 year (HR 1.85 (1.33 to 2.59); $p < 0.001$). Comparing the third versus the first and second tertiles yielded an adjusted HR of 2.14 (1.21 to 3.80; $p = 0.009$) for cardiovascular mortality at 30 days and 1.78 (1.31 to 2.43; $p < 0.001$) at 1 year. By net reclassification analysis, adding OPG tertile 3 versus tertiles 1 and 2 to a risk marker model including TIMI risk score

covariates, BMI, BNP and CRP resulted in significantly improved reclassification of patients ($p < 0.0001$).

Exploratory univariable analyses were performed to assess the relationships between OPG and subcategories of cardiovascular death. OPG was associated with the incidence of sudden death ($n = 85$, HR 2.19 (1.43 to 3.36)), the incidence of fatal HF secondary to coronary atherosclerotic disease ($n = 38$, HR 7.20 (3.41 to 15.22)) as well as with the incidence of other coronary atherosclerotic deaths ($n = 113$, HR 2.51 (1.73 to 3.63)).

Relationship to HF hospitalisation

One hundred seventy-seven patients were hospitalised for HF during the first year after ACS. By multivariable analysis, a 1 SD increase in logarithmically transformed OPG levels at baseline was associated with increased risk of HF hospitalisation (HR 1.81 (1.26 to 2.58) $p = 0.001$) (table 4). Comparing the third versus the first and second tertiles yielded an HR of 3.04 (2.26 to 4.11; $p = 0.001$) for HF hospitalisation. By multivariable analysis, OPG in the third tertile was associated with increased risk of hospitalisation for HF within 30 days (HR 1.77 (1.11 to 2.82); $p = 0.017$) and 1 year (HR 1.70 (1.22 to 2.36); $p = 0.002$). In a subgroup analysis of patients with BNP < 80 ng/l, higher OPG levels were still significantly associated with hospitalisation for HF ($p = 0.0059$). Adding OPG tertile 3 versus tertiles 1 and 2 to

Table 1 Baseline characteristics

	OPG tertile 3 (%)	n	N	OPG tertiles 1 and 2 (%)	n	N	p Value
Randomisation group (ranolazine)	48.4	719	1485	49.8	1484	2978	0.37
Age ≥ 75 years	30.1	447	1485	9.9	296	2978	< 0.001
Gender (male)	59.1	878	1485	67.7	2017	2978	< 0.001
Race (Caucasian)	96.7	1436	1485	96.8	2882	2978	0.89
BMI group							
BMI < 25 kg/m ²	24.6	361	1465	18.5	545	2943	< 0.001
BMI 25– < 30 kg/m ²	41.0	600	1465	43.6	1283	2943	
BMI ≥ 30 kg/m ²	34.4	504	1465	37.9	1115	2943	
Index diagnosis							
Unstable angina	41.1	611	1485	53.1	1582	2978	< 0.001
NSTEMI	56.7	842	1485	44.1	1312	2978	
Other	2.2	32	1485	2.8	84	2978	
Increased glucose*	7.9	66	836	5.4	113	2112	0.009
History of diabetes	42.4	629	1485	27.8	827	2978	< 0.001
History of hypertension	77.5	1145	1478	73.1	2165	2960	0.0018
History of dyslipidaemia	63.5	847	1334	69.9	1899	2715	< 0.001
Current smoker	19.6	291	1485	27.9	829	2976	< 0.001
History of CHF	23.3	346	1485	19.5	582	2978	0.0036
Family history of CAD	33.3	443	1330	42.9	1205	2812	< 0.001
Prior angina	58.9	847	1438	59.8	1755	2933	0.55
Prior myocardial infarction	35.5	521	1466	36.0	1062	2954	0.79
Prior PCI	19.7	291	1475	20.8	618	2965	0.39
Prior CABG	13.1	194	1483	11.2	332	2977	0.060
Creatinine clearance < 60 ml/min	33.2	490	1478	13.7	407	2969	< 0.001
TIMI risk score							< 0.001
0–2	17.2	255	1485	28.2	839	2978	
3–4	53.7	798	1485	53.6	1597	2978	
≥ 5	29.1	432	1485	18.2	542	2978	
Aspirin†	95.6	1420	1485	96.8	2882	2978	0.05
Clopidogrel	65.3	970	1485	62.2	1852	2978	0.041
Beta blocker	89.7	1332	1485	90.4	2692	2978	0.46
ACE-I or ARB	81.3	1208	1485	76.8	2286	2978	0.0005
Statin	78.2	1162	1485	79.0	2352	2978	0.57

*Fasting glucose > 7.0 mmol/l (126 mg/100 ml) or non-fasting glucose ≥ 11.1 mmol/l (200 mg/100 ml).

†During hospitalisation or at discharge.

ACE-I, ACE inhibitor; ARB, angiotensin receptor blocker; BMI, body mass index; CABG, coronary artery bypass graft; CAD, coronary artery disease; CHF, congestive heart failure; NSTEMI, non-ST elevation myocardial infarction; OPG, osteoprotegerin; PCI, percutaneous coronary intervention; TIMI, Thrombolysis in Myocardial Infarction.

Table 2 Osteoprotegerin and the severity of structural heart disease

	OPG tertile 3 (%)	n cases	N total	OPG tertiles 1 and 2 (%)	n cases	N total	p Value
BNP >80 ng/l	56.7	842	1484	35.6	1060	2978	<0.001
Tnl ultra positive (≥ 40 ng/l)	73.1	1069	1463	60.4	1787	2959	<0.001
Disease extent ($\geq 50\%$)*							
None	10.4	84	807	12.1	202	1669	<0.001
One VD	22.6	182	807	31.4	524	1669	
Two VD	25.5	206	807	24.0	400	1669	
Three VD or LM	41.5	335	807	32.5	543	1669	
LVEF (<50%)†	45.7	454	994	35.3	706	2000	<0.001

*Among 2476 patients with angiographic data available.

†Among 2994 patients with LVEF available.

BNP, B type natriuretic peptide; LM, left main; LVEF, left ventricular ejection fraction; OPG, osteoprotegerin; Tnl, troponin I; VD, vessel disease.

a risk marker model that included BNP and hs-CRP resulted in significantly improved reclassification of patients ($p < 0.0001$).

Relation to coronary anatomy, in-hospital ischaemia and in-hospital cardiac arrhythmias

Among patients undergoing coronary angiography ($n=2476$), patients with elevated OPG were more likely to have multivessel (≥ 2) CAD and disease involving the left anterior descending artery (table 2). Within the first 7 days of follow-up, 892 patients had significant ischaemic events and 3413 patients had significant arrhythmia on Holter recordings. Both ischaemic and significant arrhythmic events were significantly more frequent in the upper OPG versus the two lower tertiles during the first 7 days of follow-up but these relationships were attenuated after adjustment for patient age. The number of patients with significant ventricular tachycardia (ie, ≥ 8 beats) did not differ according to OPG tertile.

Relationship to recurrent ischaemia and myocardial infarction at 30 days and 1 year

In contrast with the findings in-hospital, the risk of experiencing recurrent ischaemia, defined as severe recurrent ischaemic events leading to hospitalisation or revascularization and worsening angina requiring intensification of medical therapy, at 30 days and 1 year was not associated with OPG levels. However, 86 patients in the first OPG tertile, 115 in the second and 135 patients in the third tertile experienced an AMI during the 12 months after the index event. Thus, by univariable analysis, OPG was associated with the 1 year incidence of recurrent MI (HR 1.67 (1.33 to 2.09); $p=0.001$). In a multivariable model, however, this association was attenuated and no longer significant (table 4).

Table 3 Independent predictors of higher circulating osteoprotegerin (tertile 3) at baseline

Variable	OR (95% CI)	p Value
Age (>75 years)	2.18 (1.76 to 2.72)	<0.001
Creatinine clearance <60 ml/min	1.80 (1.46 to 2.23)	<0.001
Baseline BNP (>80 ng/l)	1.60 (1.36 to 1.89)	<0.001
History of hypercholesterolaemia (yes)	0.70 (0.59 to 0.82)	<0.001
History of diabetes (yes)	2.04 (1.73 to 2.41)	<0.001
History of heart failure (yes)	1.46 (1.19 to 1.80)	<0.001
Gender (males)	0.76 (0.64 to 0.89)	0.001
Region: Eastern Europe (referent North America)	0.74 (0.59 to 0.93)	0.010
Region: Western Europe (referent North America)	1.33 (1.08 to 1.63)	0.006
Index event NSTEMI (referent UA)	1.34 (1.10 to 1.62)	0.003
Family history of CAD	0.83 (0.70 to 0.97)	0.021
Tnl ultra positive (≥ 40 ng/l)	1.24 (1.01 to 1.53)	0.042

BNP, B type natriuretic peptide; CAD, coronary artery disease; NSTEMI, non-ST elevation myocardial infarction; Tnl, troponin I; UA, unstable angina.

Relationship with treatment allocation or ejection fraction determination

No interaction between OPG levels and treatment allocation was found. Moreover, for the endpoints of cardiovascular mortality or HF, no interaction was observed between OPG levels and access to left ventricular ejection fraction determination.

DISCUSSION

In this large study of well characterised patients with NSTEMI-ACS, the main finding was that OPG is robustly and independently associated with the 30 day and 1 year risk of cardiovascular mortality and hospitalisation for HF. These associations translated into a significant reclassification of the risk of both cardiovascular death and HF hospitalisations. Although OPG was related to the angiographic severity of CAD and ischaemia on Holter recording during the first 7 days, no association with the incidence of recurrent AMI or of cardiac arrhythmias was observed after multivariable adjustment. These findings point towards additional mechanistic assessment that will help elucidate whether OPG related pathways are a potential therapeutic target in the prevention and management of cardiovascular disease.

Clinical and research implications

We have previously shown an association between OPG levels and all-cause mortality among 897 patients across the spectrum


**Figure 1** Kaplan–Meier curves of cardiovascular mortality by tertiles of osteoprotegerin (OPG) at baseline.

Table 4 Associations between osteoprotegerin concentrations (continuous) and events during follow-up

	30 day outcomes				1 year outcomes			
	Unadjusted		Adjusted		Unadjusted		Adjusted	
	HR (95% CI)	p Value	HR (95% CI)	p Value	HR (95% CI)	p Value	HR (95% CI)	p Value
CV death	4.67 (2.85 to 7.67)	<0.001	2.32 (1.30 to 4.17)	0.005	3.85 (2.92 to 5.08)	<0.001	1.85 (1.33 to 2.59)	<0.001
Severe recurrent ischemia	1.23 (0.90 to 1.69)	0.19	0.93 (0.65 to 1.33)	0.69	1.16 (0.95 to 1.40)	0.14	1.04 (0.83 to 1.29)	0.73
MI	1.57 (1.13 to 2.17)	0.006	1.11 (0.76 to 1.61)	0.59	1.67 (1.33 to 2.09)	<0.001	1.13 (0.87 to 1.46)	0.37
Hospitalisation for New or Worsening HF	4.70 (3.10 to 7.12)	<0.001	2.25 (1.38 to 3.69)	0.001	4.11 (3.04 to 5.54)	<0.001	1.81 (1.26 to 2.58)	0.001
CV death or MI	2.04 (1.53 to 2.71)	<0.001	1.35 (0.97 to 1.88)	0.072	2.17 (1.80 to 2.62)	<0.001	1.37 (1.10 to 1.70)	0.005
CV death or hospitalisation for new or worsening HF	4.38 (3.07 to 6.25)	<0.001	2.14 (1.40 to 3.25)	<0.001	3.55 (2.83 to 4.47)	<0.001	1.69 (1.28 to 2.22)	<0.001

HR associated with 1 SD increase in logarithmically transformed osteoprotegerin levels at baseline.

Adjusted for Thrombolysis in Myocardial Infarction risk score covariates (including cardiac marker), history of congestive heart failure, creatinine clearance <60 ml/min, gender, body mass index (continuous), B type natriuretic peptide (log transformed) and C reactive protein (log transformed).

CV, cardiovascular; HF, heart failure; MI, myocardial infarction.

of ACS followed for 89 months.¹⁸ However, the pathobiological mechanisms underlying the association between OPG and mortality have remained elusive. Theoretically, OPG could reflect one or more of the following: atherosclerotic burden, myocardial ischaemia, arrhythmia risk or the degree of ventricular dysfunction. For instance, observations from epidemiological studies of the general population have suggested that OPG is independently associated with the extent of coronary and aortic atherosclerosis,¹ myocardial anatomy and function,¹⁴ and the risk of atrial fibrillation.²⁶ However, to what extent these mechanisms are operative and relevant to the association between OPG and survival in the setting of NSTEMI-ACS is unclear.

We have now demonstrated that OPG is independently and strongly associated with cardiovascular mortality both early after presentation (at 30 days) and in the longer term. In addition, our current large scale study, which prospectively collected data on coronary anatomy, myocardial ischaemia, cardiac arrhythmias and HF, has provided new insight relating these potential mechanisms to the association between OPG and cardiovascular mortality. The current data suggest that HF rather than myocardial ischaemia or cardiac arrhythmias is the principal factor explaining the increased risk of death associated with higher circulating OPG levels in the setting of NSTEMI-ACS. These findings extend previous observations in smaller studies of an association between OPG and long term risk of HF after ACS¹⁸ and between OPG and risk of mortality in patients with AMI complicated by HF,¹⁷ as well as of data demonstrating associations between OPG and final infarct size,¹⁶ and between OPG levels and mortality in patients with chronic HF.^{21, 27} Moreover, our findings are compatible with experimental data showing that myocardial OPG gene expression and protein content are increased in HF,¹⁵ which suggests that OPG might play a role in left ventricular remodelling and the progression of HF.

We also observed an association between the extent of atherosclerosis on coronary angiography and OPG levels. However, as the extent of atherosclerosis is also associated with the degree of ventricular dysfunction, the extent of atherosclerosis could be an epiphenomenon or mediator of the association between OPG and ventricular dysfunction. In fact, controversy exists as to whether or not OPG is related to progression of atherosclerosis. Although some epidemiological data suggest that OPG is related to the extent of coronary atherosclerosis and calcifications,¹ this finding has not been confirmed by others.¹³ Recent experimental data also suggest that OPG may play a protective role with regard to progression of atherosclerosis. For instance, mice genetically modified to not express OPG

demonstrate accelerated advanced lesion progression,²⁸ and in a mouse model, treatment with recombinant OPG was associated with stabilisation of atherosclerotic plaques by promoting smooth muscle cell and collagen accumulation.³ In human studies, however, treatment with recombinant OPG has not been associated with altered risk of cardiovascular events.^{29, 30}

Patients with ACS, in particular in those complicated by left ventricular dysfunction or HF, are at increased risk of ventricular arrhythmias and sudden cardiac death. Moreover, recent data have suggested that OPG is related to the development of new onset atrial fibrillation.²⁶ We have previously shown that OPG levels are associated with mortality in patients with HF of both ischaemic and non-ischaemic origin,²¹ and we therefore hypothesised that part of the explanation for the association between OPG levels and increased risk of mortality after NSTEMI-ACS might be that OPG levels are associated with the risk of recurrent arrhythmic events. Although the number of patients experiencing significant arrhythmic events and sudden cardiac death increased with OPG concentration, this association was attenuated after multivariable adjustment.

Strength and limitations

The strengths of the current study include the large and well characterised contemporary cohort of NSTEMI-ACS patients. Study limitations include the fact that patients with suspected NSTEMI-ACS recruited in a clinical trial may not accurately reflect or be entirely generalisable to the broad unselected population of patients with acute chest pain. Although we found evidence of added value of OPG in risk reclassification for the endpoints cardiovascular death and HF, the clinical relevance of OPG will depend on whether or not therapeutic relevance is established.

CONCLUSION

OPG is robustly associated with 30 day and 1 year risk of cardiovascular mortality and HF development but not recurrent ischaemia or MI. These findings suggest that left ventricular dysfunction, rather than myocardial ischaemia, is the more important mediator of the adverse prognosis indicated by elevation of OPG in ACS. Our findings in this well characterised patient population add to the emerging evidence supporting involvement of OPG in the pathophysiology of ACS and its consequences, and support investigation of therapies that might modify this risk, including the elucidation of OPG related pathways that might be evaluated as targets for intervention.

Contributors RR wrote the first draft of the manuscript, participated in data analysis and reviewed the manuscript. MPB and SAM participated in data analysis and review

of the manuscript. TO participated in data analysis, and assisted in writing and review of the manuscript. BMS participated in review of the manuscript. MB and AF participated in the laboratory analysis and review of the manuscript. EB participated in the design and execution of the study, and review of the manuscript. DAM participated in the design and execution of the study, data analysis and review of the manuscript. All authors have seen and approved the final version.

Funding MERLIN-TIMI 36 was supported by CV Therapeutics (now Gilead Science Inc). The company had no involvement in the study design; in the collection, analysis and interpretation of data; in the writing of the report; or in the decision to submit the paper for publication.

Competing interests TO has received speakers' honoraria and research support from Roche Diagnostics and Abbott Laboratories. MS has received grants for clinical research via the TIMI Study Group and Brigham and Women's Hospital from Nanosphere and served as a consultant for Singulex. SAM has received consultant fees from Amarin and Eli Lilly. BMS has received grants for clinical research via the TIMI Study Group and Brigham and Women's Hospital from CV Therapeutics, Novartis Pharmaceuticals Corporation, AstraZeneca Pharmaceuticals LP, Daiichi-Sankyo Inc, Merck & Co Inc, Johnson and Johnson Pharmaceutical Research & Development LLC, Bayer HealthCare Pharmaceuticals and Bristol-Myers-Squibb Company, and has served as a consultant for AstraZeneca Pharmaceuticals LP, Novartis Pharmaceuticals Corporation, CV Therapeutics, Cogentus, Shionogi and Co Ltd, Gilead Sciences Inc, Merck & Co Inc and Schering-Plough Corporation. DAM is an investigator and receives salary from the TIMI Study Group. He has received honoraria for educational presentations from Eli Lilly. He has received remuneration for consulting from Beckman-Coulter, Boehringer Ingelheim, Cardiokinetic, Critical Diagnostics, Gilead, Instrumentation Laboratory, Icaria, Menarini, Merck, OrthoClinical Diagnostics, Servier, Roche Diagnostics and Siemens, and remuneration from AstraZeneca for adjudication as a member of a Clinical Events Committee. The TIMI Study Group has received research grant support from Accumetrics, Amgen, AstraZeneca, Beckman Coulter, BG Medicine, BRAHMS, Bristol-Myers Squibb, CV Therapeutics, Daiichi Sankyo Co Ltd, Eli Lilly and Co, GlaxoSmithKline, Integrated Therapeutics, Merck and Co, Nanosphere, Novartis Pharmaceuticals, Nuvelo, Ortho-Clinical Diagnostics, Pfizer, Roche Diagnostics, Sanofi-Aventis, Sanofi-Synthelabo, Siemens and Singulex.

Ethics approval The protocol was approved by the relevant institutional review boards at all participating centres.

Provenance and peer review Not commissioned; externally peer reviewed.

REFERENCES

1. **Abedin M**, Omland T, Ueland T, *et al*. Relation of osteoprotegerin to coronary calcium and aortic plaque (from the Dallas Heart Study). *Am J Cardiol* 2007;**99**:513–18.
2. **Van CA**, Golledge J. Osteoprotegerin, vascular calcification and atherosclerosis. *Atherosclerosis* 2009;**204**:321–9.
3. **Ovchinnikova O**, Gylfe A, Bailey L, *et al*. Osteoprotegerin promotes fibrous cap formation in atherosclerotic lesions of ApoE-deficient mice—brief report. *Arterioscler Thromb Vasc Biol* 2009;**29**:1478–80.
4. **Simonet WS**, Lacey DL, Dunstan CR, *et al*. Osteoprotegerin: a novel secreted protein involved in the regulation of bone density. *Cell* 1997;**89**:309–19.
5. **Dhore CR**, Cleutjens JP, Lutgens E, *et al*. Differential expression of bone matrix regulatory proteins in human atherosclerotic plaques. *Arterioscler Thromb Vasc Biol* 2001;**21**:1998–2003.
6. **Bucay N**, Sarosi I, Dunstan CR, *et al*. Osteoprotegerin-deficient mice develop early onset osteoporosis and arterial calcification. *Genes Dev* 1998;**12**:1260–8.
7. **Kudlacek S**, Schneider B, Woloszczuk W, *et al*. Serum levels of osteoprotegerin increase with age in a healthy adult population. *Bone* 2003;**32**:681–6.
8. **Kazama JJ**, Shigematsu T, Yano K, *et al*. Increased circulating levels of osteoclastogenesis inhibitory factor (osteoprotegerin) in patients with chronic renal failure. *Am J Kidney Dis* 2002;**39**:525–32.
9. **Secchiero P**, Corallini F, Pandolfi A, *et al*. An increased osteoprotegerin serum release characterizes the early onset of diabetes mellitus and may contribute to endothelial cell dysfunction. *Am J Pathol* 2006;**169**:2236–44.
10. **Clancy P**, Oliver L, Jayalath R, *et al*. Assessment of a serum assay for quantification of abdominal aortic calcification. *Arterioscler Thromb Vasc Biol* 2006;**26**:2574–6.
11. **Jensen JK**, Ueland T, Atar D, *et al*. Osteoprotegerin concentrations and prognosis in acute ischaemic stroke. *J Intern Med* 2010;**267**:410–17.
12. **Jono S**, Ikari Y, Shioi A, *et al*. Serum osteoprotegerin levels are associated with the presence and severity of coronary artery disease. *Circulation* 2002;**106**:1192–4.
13. **Lieb W**, Gona P, Larson MG, *et al*. Biomarkers of the osteoprotegerin pathway. Clinical correlates, subclinical disease, incident cardiovascular disease, and mortality. *Arterioscler Thromb Vasc Biol* 2010;**30**:1849–54.
14. **Omland T**, Drazner MH, Ueland T, *et al*. Plasma osteoprotegerin levels in the general population: relation to indices of left ventricular structure and function. *Hypertension* 2007;**49**:1392–8.
15. **Ueland T**, Yndestad A, Oie E, *et al*. Dysregulated osteoprotegerin/RANK ligand/RANK axis in clinical and experimental heart failure. *Circulation* 2005;**111**:2461–8.
16. **Andersen GO**, Knudsen EC, Aukrust P, *et al*. Elevated serum osteoprotegerin levels measured early after acute ST-elevation myocardial infarction predict final infarct size. *Heart* 2011;**97**:460–5.
17. **Ueland T**, Jemtland R, Godang K, *et al*. Prognostic value of osteoprotegerin in heart failure after acute myocardial infarction. *J Am Coll Cardiol* 2004;**44**:1970–6.
18. **Omland T**, Ueland T, Jansson AM, *et al*. Circulating osteoprotegerin levels and long-term prognosis in patients with acute coronary syndromes. *J Am Coll Cardiol* 2008;**51**:627–33.
19. **Morrow DA**, Scirica BM, Karwatowska-Prokopczuk E, *et al*. Evaluation of a novel anti-ischemic agent in acute coronary syndromes: design and rationale for the Metabolic Efficiency with Ranolazine for Less Ischemia in Non-ST-elevation acute coronary syndromes (MERLIN)-TIMI 36 trial. *Am Heart J* 2006;**151**:1186–9.
20. **Morrow DA**, Scirica BM, Karwatowska-Prokopczuk E, *et al*. Effects of ranolazine on recurrent cardiovascular events in patients with non-ST-elevation acute coronary syndromes: the MERLIN-TIMI 36 randomized trial. *JAMA* 2007;**297**:1775–83.
21. **Roysland R**, Masson S, Omland T, *et al*. Prognostic value of osteoprotegerin in chronic heart failure: the GISSI-HF trial. *Am Heart J* 2010;**160**:286–93.
22. **Thygesen K**, Alpert JS, White HD. Universal definition of myocardial infarction. *Eur Heart J* 2007;**28**:2525–38.
23. **de Lemos JA**, Morrow DA, Bentley JH, *et al*. The prognostic value of B-type natriuretic peptide in patients with acute coronary syndromes. *N Engl J Med* 2001;**345**:1014–21.
24. **Morrow DA**, de Lemos JA, Sabatine MS, *et al*. Evaluation of B-type natriuretic peptide for risk assessment in unstable angina/non-ST-elevation myocardial infarction: B-type natriuretic peptide and prognosis in TACTICS-TIMI 18. *J Am Coll Cardiol* 2003;**41**:1264–72.
25. **Pencina MJ**, D'Agostino RB Sr, Steyerberg EW. Extensions of net reclassification improvement calculation to measure usefulness of new biomarkers. *Stat Med* 2011;**30**:11–21.
26. **Schnabel RB**, Larson MG, Yamamoto JF, *et al*. Relation of multiple inflammatory biomarkers to incident atrial fibrillation. *Am J Cardiol* 2009;**104**:92–6.
27. **Ueland T**, Dahl CP, Kjekshus J, *et al*. Osteoprotegerin predicts progression of chronic heart failure: results from CORONA. *Circ Heart Fail* 2011;**4**:145–52.
28. **Bennett BJ**, Scatena M, Kirk EA, *et al*. Osteoprotegerin inactivation accelerates advanced atherosclerotic lesion progression and calcification in older ApoE^{-/-} mice. *Arterioscler Thromb Vasc Biol* 2006;**26**:2117–24.
29. **Cummings SR**, San MJ, McClung MR, *et al*. Denosumab for prevention of fractures in postmenopausal women with osteoporosis. *N Engl J Med* 2009;**361**:756–65.
30. **Smith MR**, Egerdie B, Hernandez TN, *et al*. Denosumab in men receiving androgen-deprivation therapy for prostate cancer. *N Engl J Med* 2009;**361**:745–55.