

Table 2, Supplemental material. Hazard ratios with 95% confidence intervals, in different age groups, for revascularisation, MACE, cardiovascular death, and non cardiovascular death, in relation to presence or absence of treatment for erectile dysfunction.

Revascularisation		MACE		Cardiovascular death		Non cardiovascular death	
Age < 60 years							
Subjects in analysis ^a	No ED treatment (n=12769)	ED treatment (n=1113)	No ED treatment (n=12799)	ED treatment (n=1083)	No ED treatment (n=12609)	ED treatment (n=1273)	No ED treatment (n=12609)
Number of cases (%)	1811 (14.2)	48 (4.3)	2195 (17.1)	57 (5.3)	178 (1.4)	10 (0.8)	207 (1.6)
Incidence rate	4.60 (4.39-4.81)	1.78 (1.34-2.36)	5.71 (5.48-5.95)	2.17 (1.68-2.82)	0.40 (0.34-0.46)	0.31 (0.17-0.58)	0.46 (0.40-0.53)
Unadjusted, HR (95% CI)	Referent	0.83 (0.62-1.11)	Referent	0.80 (0.61-1.04)	Referent	0.87 (0.46-1.67)	Referent
Age-adjusted, HR (95% CI)	Referent	0.83 (0.62-1.11)	Referent	0.79 (0.60-1.03)	Referent	0.85 (0.45-1.63)	Referent
Multivariable adjusted ^b , HR (95% CI)	Referent	0.80 (0.60-1.07)	Referent	0.77 (0.59-1.00)	Referent	0.86 (0.45-1.65)	Referent
Age 60-69 years							
Subjects in analysis ^a	No ED treatment (n=13876)	ED treatment (n=1153)	No ED treatment (n=13908)	ED treatment (n=1121)	No ED treatment (n=13733)	ED treatment (n=1296)	No ED treatment (n=13733)
Number of cases (%)	1948 (14.0)	60 (5.3)	2722 (19.6)	77 (6.9)	489 (3.6)	24 (1.9)	657 (4.8)
Incidence rate	4.70 (4.49-4.91)	2.11 (1.64-2.72)	6.85 (6.59-7.11)	2.80 (2.34-3.50)	1.04 (0.95-1.14)	0.72 (0.48-1.08)	1.40 (1.29-1.51)
Unadjusted, HR (95% CI)	Referent	1.02 (0.79-1.33)	Referent	0.87 (0.69-1.09)	Referent	0.74 (0.49-1.12)	Referent
Age-adjusted, HR (95% CI)	Referent	1.02 (0.79-1.33)	Referent	0.87 (0.69-1.10)	Referent	0.76 (0.50-1.14)	Referent
							0.71 (0.49-1.02)

CI)

Multivariable adjusted ^b ,	Referent	1.05 (0.81-1.37)	Referent	0.92 (0.73-1.16)	Referent	0.91 (0.60-1.37)	Referent	0.82 (0.57-1.18)
HR (95% CI)								

Age ≥ 70 years

Subjects in analysis ^a	No ED treatment (n=13798)	ED treatment (n=436)	No ED treatment (n=13815)	ED treatment (n=419)	No ED treatment (n=13735)	ED treatment (n=499)	No ED treatment (n=13735)	ED treatment (n=499)
Number of cases (%)	1622 (11.8)	22 (5.0)	3400 (24.6)	37 (8.8)	1484 (10.8)	11 (2.2)	1726 (12.6)	18 (3.6)
Incidence rate	4.31 (4.11-4.53)	2.04 (1.34-3.10)	9.83 (9.50-10.16)	3.61 (2.62-4.98)	3.53 (3.35-3.71)	0.85 (0.47-1.54)	4.10 (3.91-4.30)	1.40 (0.88-2.22)
Unadjusted, HR (95% CI)	Referent	1.08 (0.70-1.65)	Referent	0.68 (0.49-0.94)	Referent	0.27 (0.15-0.49)	Referent	0.35 (0.22-0.56)
Age-adjusted, HR (95% CI)	Referent	1.04 (0.68-1.59)	Referent	0.70 (0.50-0.97)	Referent	0.31 (0.17-0.57)	Referent	0.39 (0.25-0.62)
Multivariable adjusted ^b (95% CI)	Referent	1.02 (0.67-1.57)	Referent	0.76 (0.55-1.05)	Referent	0.36 (0.20-0.66)	Referent	0.44 (0.27-0.69)

Abbreviations: ED, erectile dysfunction; HR, hazard ratio; CI, confidence interval; MI, myocardial infarction; MACE, major adverse cardiac event (includes hospital stay for MI, heart failure or revascularisation).
Incidence rate is given per 100 person-years.^aNumbers do not sum up to group total since subjects can be included in the No ED treatment category initially and later change to the ED treatment category. ^bAdjusted for all variables in Table 1. Incidence rates are cases per 100 person-years.