

Table 2 Statin use by age and IHD group among subjects treated for IHD (1994-2001)

			Annual rate									
			1994	1995	1996	1997	1998	1999	2000	2001		
<b>Men (age ≥35) with IHD</b>			n	14,774	16,104	17,212	18,136	19,070	19,531	17,311	12,614	
<b>Taking statins in year by age</b>	<b>35–64</b>	r/n	443/5,220	927/5,619	1,576/5,901	2,362/6,162	3,099/6,404	3,805/6,560	3,825/5,761	2,971/4,102		
		%	<b>8.5%</b>	<b>16.5%</b>	<b>26.7%</b>	<b>38.3%</b>	<b>48.4%</b>	<b>58.0%</b>	<b>66.4%</b>	<b>72.4%</b>		
	<b>65–74</b>	r/n	176/5,528	443/5,901	873/6,232	1,512/6,535	2,291/6,800	2,978/6,907	3,238/6,043	2,677/4,304		
		%	<b>3.2%</b>	<b>7.5%</b>	<b>14.0%</b>	<b>23.1%</b>	<b>33.7%</b>	<b>43.1%</b>	<b>53.6%</b>	<b>62.2%</b>		
	<b>75–84</b>	r/n	6/3,338	45/3,790	144/4,152	303/4,202	570/4,736	888/4,890	1,178/4,477	1,245/3,446		
		%	<b>0.2%</b>	<b>1.2%</b>	<b>3.5%</b>	<b>7.1%</b>	<b>12.0%</b>	<b>18.2%</b>	<b>26.3%</b>	<b>36.1%</b>		
	<b>85</b>	r/n	0/688	0/794	0/927	7/1,037	19/1,130	22/1,174	39/1,030	44/762		
		%	<b>0%</b>	<b>0%</b>	<b>0%</b>	<b>0.7%</b>	<b>1.7%</b>	<b>1.9%</b>	<b>3.8%</b>	<b>5.8%</b>		
	<b>Taking statins in yearly IHD group</b>	<b>Non-specific IHD</b>	r/n	56/1,601	119/1,703	202/1,807	318/1,872	489/1,971	673/2,080	742/1,794	543/1,236	
			%	<b>3.5%</b>	<b>7.0%</b>	<b>11.2%</b>	<b>17.0%</b>	<b>24.8%</b>	<b>32.4%</b>	<b>41.4%</b>	<b>43.9%</b>	
<b>Angina</b>		r/n	185/6,803	431/7,201	811/7,547	1,300/7,867	1,881/8,141	2,491/8,195	2,738/7,182	2,354/5,084		
		%	<b>2.7%</b>	<b>6.0%</b>	<b>10.8%</b>	<b>16.5%</b>	<b>23.1%</b>	<b>30.4%</b>	<b>38.1%</b>	<b>46.3%</b>		
<b>Myocardial infarction</b>		r/n	242/5,119	555/5,702	991/6,076	1,658/6,383	2,363/6,695	3,007/6,852	3,220/6,093	2,636/4,461		
		%	<b>4.7%</b>	<b>9.7%</b>	<b>16.3%</b>	<b>26.0%</b>	<b>35.3%</b>	<b>43.9%</b>	<b>52.9%</b>	<b>59.1%</b>		
<b>Revascularisation</b>		r/n	142/1,251	310/1,498	589/1,782	908/2,014	1,246/2,263	1,533/2,404	1,580/2,242	1,404/1,833		
		%	<b>11.4%</b>	<b>20.7%</b>	<b>33.1%</b>	<b>45.1%</b>	<b>55.1%</b>	<b>63.3%</b>	<b>70.5%</b>	<b>76.6%</b>		
<b>Women (age ≥35) with IHD</b>			n	11,771	12,728	13,660	14,234	14,823	15,107	13,152	9,483	
<b>Taking statins in year by age</b>		<b>35–64</b>	r/n	182/2,347	329/2,491	550/2,637	771/2,676	1,033/2,796	1,257/2,889	1,263/2,524	1,015/1,741	
	%		<b>7.8%</b>	<b>13.2%</b>	<b>20.9%</b>	<b>28.8%</b>	<b>36.9%</b>	<b>43.5%</b>	<b>50.0%</b>	<b>58.3%</b>		
	<b>65–74</b>	r/n	128/3,830	292/4,061	592/4,286	980/4,450	1,443/4,534	1,850/4,553	2,015/3,975	1,637/2,771		
		%	<b>3.3%</b>	<b>7.2%</b>	<b>13.8%</b>	<b>22.0%</b>	<b>31.8%</b>	<b>40.6%</b>	<b>50.7%</b>	<b>59.1%</b>		
	<b>75–84</b>	r/n	9/4,021	51/4,396	132/4,746	250/4,953	460/5,165	801/5,225	1,037/4,559	1,049/3,351		
		%	<b>0.2%</b>	<b>1.2%</b>	<b>2.8%</b>	<b>5.1%</b>	<b>8.9%</b>	<b>15.3%</b>	<b>22.8%</b>	<b>31.3%</b>		
	<b>85</b>	r/n	0/1,573	0/1,776	5/1,991	9/2,146	21/2,328	37/2,440	65/2,094	81/1,620		
		%	<b>0%</b>	<b>0%</b>	<b>0.3%</b>	<b>0.4%</b>	<b>0.9%</b>	<b>1.5%</b>	<b>3.1%</b>	<b>5.0%</b>		
	<b>Taking statins in year by IHD group</b>	<b>Non-specific IHD</b>	r/n	34/1,493	63/1,535	128/1,614	182/1,660	274/1,713	378/1,784	425/1,472	354/1,137	
			%	<b>2.3%</b>	<b>4.1%</b>	<b>7.9%</b>	<b>11.0%</b>	<b>16.0%</b>	<b>21.2%</b>	<b>28.9%</b>	<b>31.1%</b>	
<b>Angina</b>		r/n	154/7,485	331/8,103	640/8,690	1,029/8,967	1,494/9,291	2,063/9,387	2,315/8,221	2,047/5,837		
		%	<b>2.1%</b>	<b>4.1%</b>	<b>7.4%</b>	<b>11.5%</b>	<b>16.1%</b>	<b>22.0%</b>	<b>28.2%</b>	<b>35.1%</b>		
<b>Myocardial infarction</b>		r/n	93/2,547	195/2,776	368/2,997	593/3,173	905/3,338	1,149/3,396	1,258/2,937	1,053/2,094		
		%	<b>3.7%</b>	<b>7.0%</b>	<b>12.3%</b>	<b>18.7%</b>	<b>27.1%</b>	<b>33.8%</b>	<b>42.8%</b>	<b>50.3%</b>		
<b>Revascularisation</b>		r/n	38/246	83/310	143/359	206/425	284/481	360/540	382/522	328/415		
		%	<b>15.5%</b>	<b>26.8%</b>	<b>39.8%</b>	<b>48.5%</b>	<b>59.0%</b>	<b>66.7%</b>	<b>73.2%</b>	<b>79.0%</b>		