

## Web-only References

### Management of hypertension before, during, and after pregnancy (ht35444)

PR James and C Nelson-Piercy

1. Walters BNJ, Thompson ME, Lee A *et al*. Blood pressure in the puerperium. *Clin Sci* 1986;**71**:589–594.
2. Greer IA. Hypertension. In *High Risk Pregnancy*, pp 31–93. Oxford: Butterworth Heinemann, 1999.
3. Gupta M, Shennan AH, Halligan AW, *et al*. Accuracy of oscillometric blood pressure recording in pregnancy and pre-eclampsia. *Brit J Obstet Gynaecol* 1997;**104**:350–355.
4. Penny J, Halligan WF, Shennan AH, *et al*. Blood pressure measurement in severe pre-eclampsia. *Lancet* 1997;**349**:1518.
5. Ales KL, Charlson ME. The prediction of adverse outcomes in antepartum hypertension. *Clin Exp Hypertens* 1989;**8**:95–112.
6. Sibai BM, Abdella TN, Anderson GD. Pregnancy outcome in 211 patients with mild chronic hypertension. *Obstet Gynecol* 1983;**61**:571–576.
7. Sibai BM, Akl S, Fairlie F, *et al*. A protocol for managing severe pre-eclampsia in the second trimester. *Am J Obstet Gynecol* 1990;**163**:733–739.
8. Sibai BM. Diagnosis and management of chronic hypertension in pregnancy. *Obstet Gynecol* 1991;**78**:451–461.
9. Sibai BM, Caritis SN, Thom E, *et al*. Prevention of pre-eclampsia with low-dose aspirin in healthy, nulliparous pregnant women. *N Engl J Med* 1993;**329**:1213–1218.
10. Robertson WB, Khong TY, Brosens I, *et al*. The placental bed biopsy: Review from three European centers. *Am J Obstet Gynecol* 1986;**155**:401–412.
11. Meekins JW, Pijnenborg R, Hanssens M, *et al*. A study of placental bed spiral arteries and trophoblast invasion in normal and severe pre-eclamptic pregnancies. *Br J Obstet Gynaecol* 1994;**101**:669–674.
12. Matijevic R, Johnston T. In vivo assessment of failed trophoblastic invasion of the spiral arteries in pre-eclampsia. *Br J Obstet Gynaecol* 1999;**106**:78–82.
13. Bernheim J. Hypertension in pregnancy. *Nephron* 1997;**76**:254–263.
14. Walsh SW. Maternal-placental interactions of oxidative stress and antioxidants in preeclampsia. *Semin Reprod Endocrinol* 1998;**16**:93–104.
15. O'Brien E. Ambulatory blood pressure measurement in the management of hypertension. *Heart* 2003;**89**:571–576.
16. Campbell S, Diaz-Recasens J, Griffin DR, *et al*. New Doppler technique for assessing uteroplacental blood flow. *Lancet* 1983;**1**:675–677.
17. Coleman MA, McCowan LM, North RA. Mid-trimester uterine artery Doppler screening as a predictor of adverse pregnancy outcome in high risk women. *Ultrasound Obstet Gynecol* 2000;**15**:4–6.
18. Gulmezoglu AM, Hofmeyr GJ. Bed rest in hospital for suspected impaired fetal growth (Cochrane Review). In: *The cochrane library*, Issue 2, 1999. Oxford: Update software.
19. Van Buul BJ, Steegers EA, Jongsma HW, *et al*. Dietary sodium restriction in the prophylaxis of hypertensive disorders of pregnancy: effects on the intake of other nutrients. *Am J Clin Nutrition* 1995;**62**:49–57.
20. Secher NH, Olsen SF. Fish-oil and pre-eclampsia. *Brit J Obstet Gynaecol* 1990;**97**:1077–1079.

21. Impey L. Severe hypertension and fetal distress following sublingual administration of nifedipine to a patient with severe pregnancy induced hypertension at 33 weeks. *Brit J Obstet Gynaecol* 1993;**100**:959–961.
22. Waisman GD, Mayorga LM, Camera MI, *et al.* Magnesium plus nifedipine: potentiation of hypotensive effect in pre-eclampsia. *Am J Obstet Gynecol* 1988;**159**:308–309.
23. Rey E, Le Lorier J, Burgess E, *et al.* Report of Canadian Hypertension Society Consensus Conference: Pharmacologic treatment of hypertensive disorders of pregnancy. *Can Med Assoc J* 1997;**157**:1245–1254.
24. Rubin PC, Butters L, Low RA, *et al.* Clinical pharmacological studies with prazosin during pregnancy complicated by hypertension. *Brit J Clin Pharmacol* 1983;**16**:543–547.
25. Gallery EDM, Hunyor SN, Gyory AZ. Plasma volume contraction: a significant factor in both pregnancy-related hypertension (preeclampsia) and chronic hypertension in pregnancy. *Quat J Med* 1979;**192**:593–602.
26. Sibai BM, Grossman RA, Grossman HG. Effects of diuretics on plasma volume in pregnancies with long-term hypertension. *Am J Obstet Gynecol* 1984;**150**:831–835.
27. National high blood pressure education program working group report on high blood pressure in pregnancy. *Am J Obstet Gynecol* 1990;**163**:1691–1712.
28. Brown MA, Zammit VC, Lowe SA. Capillary permeability and extracellular fluid volumes in pregnancy-induced hypertension. *Clin Sci* 1989;**77**:599–604.
29. Afessa B, Green B, Delke I, *et al.* Systemic inflammatory response syndrome, organ failure, and outcome in critically ill obstetric patients in an ICU. *Chest* 2001;**120**:1271–7.
30. Burrows RF, Burrows EA. Assessing the teratogenic potential of angiotensin-converting-enzyme inhibitors in pregnancy. *Aust N Z J Obstet Gynaecol* 1998;**38**:306–311.
31. Mann RD, Mackay F, Pearce G, *et al.* Losartan: a study of pharmacovigilance data on 14522 patients. *J Hum Hyperten* 1999;**13**:551–557.
32. Barr M. Teratogen update: Angiotensin-converting enzyme inhibitors. *Teratology* 1994;**50**:399–409.
33. Saji H, Yamanaka M, Hagiwara A, *et al.* Losartan and fetal toxic effects. *Lancet* 2001;**357**:363.
34. Chua S, Redman CW. Prognosis for pre-eclampsia complicated by 5♣g or more of proteinuria in 24♣hours. *Eur J Obstet Gynecol Reprod Biol* 1992;**43**:9–12.
35. Sibai Bm, Sarinoglu C, Mercer BM. Eclampsia. VII. Pregnancy outcome after eclampsia and long-term prognosis. *Am J Obstet Gynecol* 1992;**166**:1757–1763.
36. Rasmussen S, Irgens L, Albrechtsen S, *et al.* Predicting preeclampsia in the second pregnancy from low birth weight in the first pregnancy. *Obstet Gynecol* 2000;**96**:696–700.
37. Abundis EM, Ortiz MG, Galvan AQ, *et al.* Hyperinsulinaemia in glucose tolerant women with preeclampsia. A controlled study. *Am J Hypertens* 1996;**9**:610–614.
38. Kaaja R. Insulin resistance syndrome in preeclampsia. *Semin Reprod Endocrinol* 1998;**16**:41–46.
39. Hubel CA, McLaughlin MK, Evans RW, *et al.* Fasting triglycerides, free fatty acids, and malondialdehyde are increased in preeclampsia, are positively correlated, and decrease within 48♣hours postpartum. *Am J Obstet Gynecol* 1996;**174**:975–982.

40. Soloman CG, CarrollJS, Okamura K, *et al.* Higher cholesterol and insulin levels are associated with increased risk for pregnancy-induced hypertension. *Am J Hypertens* 1999;**12**:276–282.
41. Van den Elzen HJ, Wladimiroff JW, Cohen-Overbeek TE, *et al.* Serum lipids in early pregnancy and risk of pre-eclampsia. *Brit J Obstet Gynaecol* 1996;**103**:117–122.